



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 10:45 pm BST

PDB ID : 3SB4  
Title : Crystal structure of a leucine-rich repeat protein (BT\_1240) from Bacteroides  
thetaiotaomicron VPI-5482 at 1.99 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2011-06-03  
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	<div> <div>9%</div> <div>91%</div> <div>9%</div> </div>
1	B	329	<div> <div>12%</div> <div>88%</div> <div>12%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	351	-	-	-	X
2	GOL	A	357	-	-	-	X
2	GOL	A	362	-	-	-	X
2	GOL	B	356	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5738 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypothetical leucine rich repeat protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	329	Total	C	N	O	S	Se	0	6	0
			2586	1663	424	484	4	11			
1	B	328	Total	C	N	O	S	Se	0	11	0
			2642	1700	434	491	4	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	leader sequence	UNP Q8A8C7
B	0	GLY	-	leader sequence	UNP Q8A8C7

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C<sub>8</sub>H<sub>18</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			13	8	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	193	Total	O	0	1
			194	194		
4	B	224	Total	O	0	1
			225	225		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hypothetical leucine rich repeat protein



- Molecule 1: Hypothetical leucine rich repeat protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.21Å 62.13Å 66.86Å 109.80° 93.75° 91.05°	Depositor
Resolution (Å)	27.82 – 1.99 27.82 – 1.99	Depositor EDS
% Data completeness (in resolution range)	(Not available) (27.82-1.99) 95.8 (27.82-1.99)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.06 (at 1.98Å)	Xtriage
Refinement program	BUSTER-TNT 2.8.0, BUSTER 2.8.0	Depositor
R, $R_{free}$	0.168 , 0.199 0.167 , 0.200	Depositor DCC
$R_{free}$ test set	2653 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.1	Xtriage
Anisotropy	0.244	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 51.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5738	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.49	0/2633	0.64	0/3541
1	B	0.49	0/2693	0.64	0/3620
All	All	0.49	0/5326	0.64	0/7161

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2586	0	2615	22	0
1	B	2642	0	2699	29	0
2	A	54	0	72	4	0
2	B	24	0	32	6	0
3	B	13	0	18	2	0
4	A	194	0	0	1	0
4	B	225	0	0	2	0
All	All	5738	0	5436	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:100[B]:MSE:SE	4:B:670:HOH:O	2.20	1.10
1:A:161:PRO:HB2	1:B:186:ARG:NH2	2.15	0.62
1:B:279:HIS:NE2	3:B:363:PG4:H32	2.15	0.61
1:A:83:MSE:HE3	1:A:100:MSE:SE	2.50	0.61
1:B:250:SER:OG	2:B:352:GOL:H32	2.02	0.59
1:A:199:GLU:HB3	2:A:359:GOL:H11	1.84	0.59
1:A:36:ILE:HD13	1:A:63:HIS:HB3	1.84	0.58
1:A:161:PRO:HB3	1:B:161:PRO:CG	2.34	0.57
1:B:294:ARG:HH12	2:B:360:GOL:H32	1.68	0.57
1:A:273:LEU:HD23	1:A:294:ARG:HB3	1.86	0.57
1:B:241:ILE:HG23	1:B:245[A]:MSE:HE2	1.85	0.57
1:B:257:ASN:HA	3:B:363:PG4:H31	1.87	0.57
1:B:186:ARG:NH2	4:B:556:HOH:O	2.40	0.54
1:B:260[B]:THR:HG22	1:B:283:THR:HB	1.91	0.53
1:B:266:PHE:O	1:B:292:CYS:SG	2.68	0.52
1:B:126[A]:VAL:HG23	1:B:146:LEU:HD11	1.92	0.52
1:A:71:LEU:HB3	1:A:120:LYS:HE2	1.91	0.51
1:A:147:LYS:NZ	2:A:358:GOL:H2	2.25	0.51
1:B:252:ASP:HB2	2:B:352:GOL:H31	1.92	0.51
1:B:71:LEU:HB3	1:B:120:LYS:HE2	1.92	0.51
1:A:245[B]:MSE:CE	1:A:248:LEU:HD13	2.42	0.50
1:A:83:MSE:CE	1:A:100:MSE:SE	3.09	0.50
1:A:299:LEU:HA	2:A:354:GOL:H2	1.97	0.47
1:B:45:ASN:HA	1:B:69:PRO:HD2	1.96	0.47
1:A:227:LEU:CD2	1:A:245[B]:MSE:HE1	2.46	0.46
1:B:221:PRO:HG2	2:B:350:GOL:H32	1.98	0.46
1:A:199:GLU:HB3	2:A:359:GOL:H32	1.98	0.45
1:A:39[A]:MSE:HB2	1:A:43:GLU:HG2	1.99	0.44
1:B:211:GLU:OE2	2:B:350:GOL:H12	2.17	0.44
1:B:311:GLY:CA	1:B:314[B]:MSE:HG3	2.47	0.44
1:A:45:ASN:HA	1:A:69:PRO:HD2	1.99	0.44
1:A:311:GLY:HA2	1:A:335:GLU:O	2.18	0.43
1:B:164:LEU:HD22	1:B:169[B]:THR:HG21	1.99	0.43
1:A:279:HIS:HE1	4:A:438:HOH:O	2.01	0.43
1:B:135:ILE:HG22	1:B:163:ALA:HB1	2.01	0.43
1:A:317:ASP:HB3	1:B:116:VAL:HG23	2.01	0.42
1:B:107:TYR:HB3	1:B:110:SER:O	2.20	0.42
1:A:161:PRO:HB3	1:B:161:PRO:HG2	2.01	0.42
1:B:270:LYS:HD3	2:B:360:GOL:H31	2.02	0.42
1:A:337:PHE:CE1	1:A:344:LYS:HB2	2.55	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:314[B]:MSE:SE	1:B:338:GLY:HA2	2.70	0.41
1:B:273:LEU:HD23	1:B:294:ARG:HB3	2.03	0.41
1:B:337:PHE:CE1	1:B:344:LYS:HB2	2.55	0.41
1:A:161:PRO:HB2	1:B:186:ARG:HH22	1.85	0.41
1:B:311:GLY:HA2	1:B:335:GLU:O	2.20	0.41
1:A:107:TYR:HB3	1:A:110:SER:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	332/329 (101%)	322 (97%)	9 (3%)	1 (0%)	41	37
1	B	337/329 (102%)	333 (99%)	4 (1%)	0	100	100
All	All	669/658 (102%)	655 (98%)	13 (2%)	1 (0%)	47	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	22	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/272 (102%)	276 (100%)	1 (0%)	91	93
1	B	289/272 (106%)	288 (100%)	1 (0%)	92	95
All	All	566/544 (104%)	564 (100%)	2 (0%)	91	93

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	207	MSE
1	B	349	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	A	358	-	5,5,5	0.74	0	5,5,5	0.51	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	A	353	-	5,5,5	0.41	0	5,5,5	0.34	0
2	GOL	A	359	-	5,5,5	1.29	0	5,5,5	1.04	0
2	GOL	A	357	-	5,5,5	0.59	0	5,5,5	0.59	0
3	PG4	B	363	-	12,12,12	0.63	0	11,11,11	1.00	1 (9%)
2	GOL	B	352	-	5,5,5	0.60	0	5,5,5	0.36	0
2	GOL	A	354	-	5,5,5	0.65	0	5,5,5	1.07	0
2	GOL	B	360	-	5,5,5	0.45	0	5,5,5	0.77	0
2	GOL	B	350	-	5,5,5	0.64	0	5,5,5	0.76	0
2	GOL	A	351	-	5,5,5	0.54	0	5,5,5	0.23	0
2	GOL	A	361	-	5,5,5	0.43	0	5,5,5	0.72	0
2	GOL	A	355	-	5,5,5	0.66	0	5,5,5	0.38	0
2	GOL	A	362	-	5,5,5	0.51	0	5,5,5	0.69	0
2	GOL	B	356	-	5,5,5	0.39	0	5,5,5	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	358	-	-	4/4/4/4	-
2	GOL	A	353	-	-	0/4/4/4	-
2	GOL	A	359	-	-	4/4/4/4	-
2	GOL	A	357	-	-	2/4/4/4	-
3	PG4	B	363	-	-	6/10/10/10	-
2	GOL	B	352	-	-	2/4/4/4	-
2	GOL	A	354	-	-	1/4/4/4	-
2	GOL	B	360	-	-	4/4/4/4	-
2	GOL	B	350	-	-	2/4/4/4	-
2	GOL	A	351	-	-	2/4/4/4	-
2	GOL	A	361	-	-	3/4/4/4	-
2	GOL	A	355	-	-	0/4/4/4	-
2	GOL	A	362	-	-	1/4/4/4	-
2	GOL	B	356	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	363	PG4	C3-O2-C2	-2.74	101.41	113.29

There are no chirality outliers.

All (31) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	358	GOL	O1-C1-C2-C3
2	A	357	GOL	O1-C1-C2-C3
2	B	360	GOL	O1-C1-C2-C3
2	B	360	GOL	C1-C2-C3-O3
2	A	359	GOL	O1-C1-C2-C3
2	A	359	GOL	C1-C2-C3-O3
2	A	359	GOL	O2-C2-C3-O3
2	B	350	GOL	O1-C1-C2-C3
2	A	351	GOL	O1-C1-C2-O2
2	A	351	GOL	O1-C1-C2-C3
3	B	363	PG4	O2-C3-C4-O3
2	A	358	GOL	C1-C2-C3-O3
2	B	352	GOL	C1-C2-C3-O3
2	A	354	GOL	O1-C1-C2-C3
2	A	361	GOL	C1-C2-C3-O3
3	B	363	PG4	O1-C1-C2-O2
2	A	358	GOL	O1-C1-C2-O2
2	A	358	GOL	O2-C2-C3-O3
2	A	357	GOL	O1-C1-C2-O2
2	B	360	GOL	O2-C2-C3-O3
2	A	359	GOL	O1-C1-C2-O2
2	A	361	GOL	O2-C2-C3-O3
3	B	363	PG4	O4-C7-C8-O5
3	B	363	PG4	O3-C5-C6-O4
2	B	350	GOL	O1-C1-C2-O2
2	A	361	GOL	O1-C1-C2-C3
2	A	362	GOL	O1-C1-C2-O2
2	B	352	GOL	O2-C2-C3-O3
3	B	363	PG4	C5-C6-O4-C7
2	B	360	GOL	O1-C1-C2-O2
3	B	363	PG4	C6-C5-O3-C4

There are no ring outliers.

7 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	358	GOL	1	0
2	A	359	GOL	2	0
3	B	363	PG4	2	0
2	B	352	GOL	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	354	GOL	1	0
2	B	360	GOL	2	0
2	B	350	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	321/329 (97%)	0.34	31 (9%) <b>7</b> <b>7</b>	34, 46, 65, 83	0
1	B	320/329 (97%)	0.43	38 (11%) <b>4</b> <b>4</b>	34, 44, 63, 94	0
All	All	641/658 (97%)	0.38	69 (10%) <b>5</b> <b>5</b>	34, 45, 65, 94	0

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	135	ILE	5.1
1	B	113	VAL	5.1
1	B	126[A]	VAL	4.8
1	B	104	VAL	4.7
1	B	114	ASN	4.5
1	A	0	GLY	4.3
1	B	140	PHE	3.9
1	B	128	LEU	3.7
1	A	126[B]	VAL	3.6
1	B	266	PHE	3.6
1	A	96	PHE	3.5
1	A	277	LEU	3.4
1	A	113	VAL	3.2
1	B	74	LEU	3.2
1	A	266	PHE	3.1
1	B	164	LEU	3.0
1	B	288	VAL	3.0
1	A	61	PHE	3.0
1	A	135	ILE	2.9
1	B	105	PRO	2.9
1	A	64	LEU	2.9
1	B	289	PHE	2.9
1	B	277	LEU	2.8
1	B	272	LEU	2.8

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	116	VAL	2.8
1	A	289	PHE	2.8
1	B	96	PHE	2.8
1	A	288	VAL	2.7
1	B	175	LEU	2.7
1	B	251	LEU	2.7
1	B	284	ILE	2.6
1	A	340	GLY	2.6
1	B	148	ILE	2.6
1	B	183	PHE	2.6
1	A	116	VAL	2.6
1	B	248	LEU	2.6
1	B	109	PHE	2.5
1	A	104	VAL	2.5
1	A	327	ASP	2.5
1	A	164	LEU	2.5
1	A	140	PHE	2.5
1	B	169[A]	THR	2.5
1	B	339	ASN	2.5
1	A	206	ALA	2.4
1	B	139	ALA	2.4
1	B	61	PHE	2.4
1	A	115	GLY	2.4
1	A	76	ILE	2.4
1	B	22	GLN	2.4
1	A	22	GLN	2.4
1	B	340	GLY	2.3
1	A	284	ILE	2.3
1	B	118[A]	LYS	2.3
1	A	109	PHE	2.2
1	A	183	PHE	2.2
1	B	108	ALA	2.2
1	B	290	SER	2.2
1	B	115	GLY	2.2
1	A	261	ILE	2.2
1	A	329	ILE	2.2
1	A	128	LEU	2.2
1	B	275	ILE	2.1
1	A	74	LEU	2.1
1	A	216	LYS	2.1
1	B	218	GLY	2.1
1	A	114	ASN	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	301	LEU	2.0
1	B	170	ALA	2.0
1	A	251	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	B	356	6/6	0.26	0.48	96,97,97,98	0
2	GOL	A	358	6/6	0.39	0.28	80,82,82,83	0
2	GOL	A	359	6/6	0.57	0.36	55,59,60,62	0
2	GOL	A	355	6/6	0.59	0.37	79,80,81,81	0
2	GOL	B	352	6/6	0.61	0.27	66,69,70,71	0
3	PG4	B	363	13/13	0.65	0.30	79,80,84,87	0
2	GOL	A	362	6/6	0.66	0.48	83,84,84,85	0
2	GOL	B	350	6/6	0.69	0.36	82,83,83,84	0
2	GOL	A	357	6/6	0.70	0.51	93,94,94,94	0
2	GOL	A	351	6/6	0.74	0.51	93,94,94,95	0
2	GOL	B	360	6/6	0.75	0.27	86,87,87,88	0
2	GOL	A	361	6/6	0.79	0.35	95,95,95,96	0
2	GOL	A	354	6/6	0.79	0.22	63,67,67,68	0
2	GOL	A	353	6/6	0.89	0.19	106,106,106,107	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.