



# Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2020 – 01:13 pm BST

PDB ID : 3SE0  
Title : Structural characterization of the subunit A mutant F508W of the A-ATP synthase from *Pyrococcus horikoshii*  
Authors : Tadwal, V.S.; Manimekalai, M.S.S.; Balakrishna, A.M.; Gruber, G.  
Deposited on : 2011-06-09  
Resolution : 2.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3797 (2.64-2.60)
Clashscore	141614	4168 (2.64-2.60)
Ramachandran outliers	138981	4093 (2.64-2.60)
Sidechain outliers	138945	4093 (2.64-2.60)
RSRZ outliers	127900	3731 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	588	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MPD	A	594	-	-	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4297 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called V-type ATP synthase alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	519	4109	2629	697	767	16	0	2	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	508	TRP	PHE	ENGINEERED MUTATION	UNP O57728

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

- Molecule 3 is ACETIC ACID (three-letter code: ACY) (formula: C<sub>2</sub>H<sub>4</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula:  $C_6H_{14}O_2$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		
4	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula:  $C_4H_{12}NO_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	143	Total	O	0	0
			143	143		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: V-type ATP synthase alpha chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	128.54Å 128.54Å 103.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.86 – 2.62 29.86 – 2.62	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.86-2.62) 99.8 (29.86-2.62)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.42 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, $R_{free}$	0.224 , 0.272 0.232 , 0.279	Depositor DCC
$R_{free}$ test set	1350 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	59.6	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 61.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4297	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, ACY, MPD, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/4206	0.64	2/5699 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	427	PHE	C-N-CD	-9.66	99.34	120.60
1	A	360	LEU	CA-CB-CG	5.03	126.87	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	199	ARG	Sidechain
1	A	427	PHE	Peptide

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within



the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4109	0	4155	79	0
2	A	5	0	0	0	0
3	A	8	0	6	0	0
4	A	24	0	42	8	0
5	A	8	0	12	0	0
6	A	143	0	0	7	0
All	All	4297	0	4215	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:241:THR:HG22	1:A:244:GLN:HE22	1.12	1.07
1:A:241:THR:HG22	1:A:244:GLN:NE2	1.83	0.93
1:A:68:THR:HG21	6:A:704:HOH:O	1.67	0.92
1:A:75:GLU:H	1:A:89:GLN:HE22	1.18	0.91
1:A:471:GLU:HB3	4:A:595:MPD:HM3	1.51	0.91
1:A:244:GLN:HE21	1:A:244:GLN:H	1.16	0.89
1:A:216:ARG:H	1:A:505:GLN:HE22	1.24	0.85
1:A:448:HIS:HE1	1:A:456:LYS:H	1.26	0.81
1:A:490:ALA:HB2	1:A:538:ILE:HD12	1.71	0.73
1:A:114:ARG:HE	1:A:170:ILE:HD11	1.53	0.73
1:A:199:ARG:NH1	1:A:321:ASP:OD2	2.22	0.73
1:A:490:ALA:HB2	1:A:538:ILE:CD1	2.19	0.72
1:A:356:TYR:H	1:A:357:PRO:HD2	1.54	0.69
1:A:373:ARG:HD2	6:A:603:HOH:O	1.93	0.68
1:A:249:LYS:NZ	4:A:594:MPD:HM2	2.08	0.68
1:A:241:THR:CG2	1:A:244:GLN:HE22	2.00	0.66
1:A:75:GLU:H	1:A:89:GLN:NE2	1.93	0.65
1:A:341:ARG:CG	1:A:342:GLU:H	2.08	0.65
1:A:320:ARG:HG2	1:A:386:VAL:HG23	1.77	0.65
1:A:341:ARG:HG3	1:A:342:GLU:N	2.13	0.64
1:A:448:HIS:CE1	1:A:456:LYS:H	2.12	0.64
1:A:199:ARG:HH11	1:A:318:TYR:HA	1.61	0.63
1:A:204:LYS:HD2	1:A:372:GLY:HA3	1.79	0.63
1:A:216:ARG:H	1:A:505:GLN:NE2	1.94	0.63
1:A:147:HIS:HE1	1:A:318:TYR:OH	1.84	0.61
1:A:341:ARG:HG3	1:A:342:GLU:H	1.66	0.59

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:428:PRO:HB2	6:A:620:HOH:O	2.03	0.59
1:A:554:GLU:OE2	1:A:558:ARG:NH2	2.31	0.58
1:A:242:VAL:HB	4:A:594:MPD:H13	1.85	0.57
1:A:104:ALA:HB2	6:A:729:HOH:O	2.04	0.57
1:A:249:LYS:HZ2	4:A:594:MPD:HM2	1.70	0.57
1:A:98:LYS:HE2	1:A:303:ALA:HB1	1.87	0.57
1:A:494:VAL:HG11	1:A:531:TYR:HB2	1.87	0.56
1:A:275:PHE:HE1	6:A:736:HOH:O	1.89	0.55
1:A:360:LEU:H	1:A:360:LEU:HD13	1.72	0.55
1:A:141:GLU:OE1	1:A:147:HIS:HD2	1.90	0.55
1:A:341:ARG:CG	1:A:342:GLU:N	2.71	0.54
1:A:317:GLU:O	1:A:320:ARG:HG3	2.08	0.54
1:A:467:GLN:NE2	6:A:629:HOH:O	2.30	0.54
1:A:199:ARG:NH1	1:A:318:TYR:HA	2.23	0.53
1:A:249:LYS:HZ1	4:A:594:MPD:CM	2.22	0.53
1:A:153:PRO:HG2	1:A:193:TRP:CZ3	2.44	0.52
1:A:249:LYS:NZ	4:A:594:MPD:CM	2.74	0.51
1:A:244:GLN:NE2	1:A:244:GLN:H	1.97	0.51
1:A:68:THR:HG22	1:A:69:GLY:N	2.26	0.50
1:A:236:PHE:CE2	1:A:391:ALA:HB2	2.47	0.49
1:A:490:ALA:CB	1:A:538:ILE:HD12	2.40	0.49
1:A:356:TYR:H	1:A:357:PRO:CD	2.22	0.48
1:A:65:VAL:HB	1:A:359:TYR:CE1	2.48	0.48
1:A:249:LYS:CE	4:A:594:MPD:H4	2.44	0.47
1:A:123:LYS:HG2	1:A:137:GLY:HA2	1.97	0.47
1:A:261:CYS:HB3	1:A:331:ASP:HB3	1.97	0.47
1:A:271:VAL:HB	1:A:274:GLU:HB2	1.96	0.46
1:A:91:PRO:HD2	1:A:109:ALA:HB3	1.98	0.46
1:A:143:SER:OG	1:A:289:GLU:OE2	2.27	0.46
1:A:70:ALA:HB3	1:A:103:ILE:HG12	1.98	0.46
1:A:159:ILE:HD13	1:A:177:VAL:HG22	1.98	0.45
1:A:333:THR:C	1:A:335:ARG:H	2.20	0.45
1:A:275:PHE:HB3	1:A:288:MET:HE3	2.00	0.44
1:A:468:LYS:HE3	1:A:468:LYS:HA	1.99	0.44
1:A:392:VAL:HG13	1:A:404:VAL:HG12	2.00	0.44
1:A:399:PHE:C	1:A:404:VAL:HG21	2.38	0.44
1:A:123:LYS:HD3	1:A:123:LYS:HA	1.68	0.44
1:A:244:GLN:HE21	1:A:244:GLN:N	1.99	0.44
1:A:200:PRO:HG2	1:A:377:LEU:HD11	2.00	0.43
1:A:452:ASP:HA	1:A:453:PRO:HD3	1.87	0.43
1:A:87:GLY:HA3	1:A:304:ALA:O	2.19	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:241:THR:HG23	6:A:654:HOH:O	2.19	0.42
1:A:132:GLY:O	1:A:377:LEU:HB3	2.19	0.42
1:A:486:GLU:OE2	1:A:489:ARG:NH1	2.53	0.41
1:A:65:VAL:HB	1:A:359:TYR:CZ	2.56	0.41
1:A:399:PHE:HA	1:A:404:VAL:HG11	2.03	0.41
1:A:221:PHE:CE1	1:A:503:LEU:HD11	2.56	0.41
1:A:242:VAL:HB	4:A:594:MPD:C1	2.51	0.41
1:A:216:ARG:N	1:A:505:GLN:HE22	2.04	0.41
1:A:271:VAL:HG23	1:A:275:PHE:CD2	2.56	0.41
1:A:216:ARG:HG2	1:A:520:GLN:HG2	2.03	0.41
1:A:227:GLY:HA2	1:A:386:VAL:O	2.21	0.40
1:A:90:ARG:HA	1:A:91:PRO:HD3	1.93	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	517/588 (88%)	479 (93%)	31 (6%)	7 (1%)	11	21

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	271	VAL
1	A	334	SER
1	A	428	PRO
1	A	356	TYR
1	A	71	SER
1	A	107	VAL
1	A	336	TRP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	441/492 (90%)	403 (91%)	38 (9%)	10	19

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	SER
1	A	81	LEU
1	A	89	GLN
1	A	92	LEU
1	A	93	GLU
1	A	165	GLU
1	A	167	ASP
1	A	204	LYS
1	A	208	GLU
1	A	209	VAL
1	A	240	LYS
1	A	244	GLN
1	A	261	CYS
1	A	263	GLU
1	A	268	MET
1	A	269	THR
1	A	279	LYS
1	A	285	LYS
1	A	299	ASN
1	A	320	ARG
1	A	334	SER
1	A	354	GLU
1	A	360	LEU
1	A	363	LYS
1	A	392	VAL
1	A	405	GLN
1	A	417	LEU
1	A	421	LEU
1	A	433	LEU
1	A	459	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	465	LEU
1	A	468	LYS
1	A	477	ARG
1	A	484	LEU
1	A	493	LEU
1	A	503	LEU
1	A	539	ASN
1	A	584	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	GLN
1	A	147	HIS
1	A	244	GLN
1	A	299	ASN
1	A	405	GLN
1	A	448	HIS
1	A	467	GLN
1	A	504	GLN
1	A	505	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	MPD	A	591	-	7,7,7	0.33	0	9,10,10	0.51	0
4	MPD	A	595	-	7,7,7	0.33	0	9,10,10	0.52	0
2	SO4	A	589	-	4,4,4	0.14	0	6,6,6	0.10	0
5	TRS	A	592	-	7,7,7	0.32	0	9,9,9	0.36	0
3	ACY	A	593	-	1,3,3	1.28	0	0,3,3	0.00	-
4	MPD	A	594	-	7,7,7	0.32	0	9,10,10	0.52	0
3	ACY	A	590	-	1,3,3	1.78	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	A	591	-	-	0/5/5/5	-
4	MPD	A	594	-	-	0/5/5/5	-
5	TRS	A	592	-	-	3/9/9/9	-
4	MPD	A	595	-	-	0/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	592	TRS	C2-C-C3-O3
5	A	592	TRS	N-C-C3-O3
5	A	592	TRS	C1-C-C3-O3

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	595	MPD	1	0
4	A	594	MPD	7	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	519/588 (88%)	0.37	52 (10%) 7 4	31, 57, 130, 186	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	106	GLY	9.0
1	A	351	PRO	8.5
1	A	107	VAL	8.0
1	A	96	ARG	7.9
1	A	95	ILE	7.5
1	A	352	GLY	7.5
1	A	265	GLY	6.4
1	A	268	MET	5.8
1	A	266	ASN	5.1
1	A	273	GLU	4.8
1	A	108	THR	4.5
1	A	355	GLY	4.5
1	A	103	ILE	4.4
1	A	67	GLY	4.3
1	A	65	VAL	4.3
1	A	102	PHE	4.0
1	A	64	PRO	3.8
1	A	338	GLU	3.8
1	A	344	SER	3.8
1	A	269	THR	3.7
1	A	264	ARG	3.7
1	A	263	GLU	3.6
1	A	400	SER	3.5
1	A	272	LEU	3.4
1	A	97	GLU	3.2
1	A	267	GLU	3.2
1	A	342	GLU	3.2

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	270	ASP	3.1
1	A	345	GLY	3.1
1	A	341	ARG	3.1
1	A	68	THR	3.1
1	A	104	ALA	2.9
1	A	105	ARG	2.9
1	A	94	VAL	2.9
1	A	262	GLY	2.8
1	A	311	THR	2.8
1	A	398	ASP	2.7
1	A	343	ILE	2.5
1	A	170	ILE	2.5
1	A	66	VAL	2.5
1	A	337	ALA	2.4
1	A	584	LYS	2.4
1	A	109	ALA	2.4
1	A	217	VAL	2.3
1	A	315	ILE	2.3
1	A	261	CYS	2.2
1	A	357	PRO	2.2
1	A	353	GLU	2.2
1	A	257	ILE	2.1
1	A	335	ARG	2.1
1	A	124	ALA	2.0
1	A	71	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ACY	A	590	4/4	0.67	0.37	93,93,93,94	0
4	MPD	A	595	8/8	0.73	0.35	84,85,86,86	0
4	MPD	A	591	8/8	0.81	0.33	84,85,86,86	0
5	TRS	A	592	8/8	0.83	0.47	96,97,97,97	0
4	MPD	A	594	8/8	0.89	0.27	84,85,86,86	0
2	SO4	A	589	5/5	0.90	0.25	106,107,107,107	0
3	ACY	A	593	4/4	0.91	0.21	80,80,80,81	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.