



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 10:47 pm BST

PDB ID : 3SEQ  
Title : Crystal structure of C176A mutant of glutamine-dependent NAD<sup>+</sup> synthetase from *M. tuberculosis* in complex with AMPCPP and NaAD<sup>+</sup>  
Authors : Chuenchor, W.; Gerratana, B.  
Deposited on : 2011-06-10  
Resolution : 2.73 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1271 (2.76-2.72)
Clashscore	141614	1322 (2.76-2.72)
Ramachandran outliers	138981	1297 (2.76-2.72)
Sidechain outliers	138945	1298 (2.76-2.72)
RSRZ outliers	127900	1243 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	680	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 83%, yellow 13%, green 4%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>2%</span> <span>83%</span> <span>13%</span> <span>• •</span> </div> </div>
1	B	680	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, orange 83%, yellow 13%, green 4%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>83%</span> <span>13%</span> <span>• •</span> </div> </div>
1	C	680	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 80%, yellow 15%, green 4%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>2%</span> <span>80%</span> <span>15%</span> <span>• •</span> </div> </div>
1	D	680	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 81%, yellow 14%, green 4%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>2%</span> <span>81%</span> <span>14%</span> <span>• •</span> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20820 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glutamine-dependent NAD(+) synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	661	Total	C	N	O	S	0	0	0
			5102	3234	905	948	15			
1	B	658	Total	C	N	O	S	0	0	0
			5063	3215	894	939	15			
1	C	650	Total	C	N	O	S	0	0	0
			4992	3171	885	921	15			
1	D	650	Total	C	N	O	S	0	0	0
			5002	3174	887	926	15			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP P0A5L6
A	176	ALA	CYS	engineered mutation	UNP P0A5L6
B	0	SER	-	expression tag	UNP P0A5L6
B	176	ALA	CYS	engineered mutation	UNP P0A5L6
C	0	SER	-	expression tag	UNP P0A5L6
C	176	ALA	CYS	engineered mutation	UNP P0A5L6
D	0	SER	-	expression tag	UNP P0A5L6
D	176	ALA	CYS	engineered mutation	UNP P0A5L6

- Molecule 2 is DIPHOSPHOMETHYLPHOSPHONIC ACID ADENOSYL ESTER (three-letter code: APC) (formula: C<sub>11</sub>H<sub>18</sub>N<sub>5</sub>O<sub>12</sub>P<sub>3</sub>).



Id	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 31	C 11	N 5	O 12	P 3	0	0
2	B	1	Total 31	C 11	N 5	O 12	P 3	0	0
2	C	1	Total 31	C 11	N 5	O 12	P 3	0	0
2	D	1	Total 31	C 11	N 5	O 12	P 3	0	0

- Molecule 3 is NICOTINIC ACID ADENINE DINUCLEOTIDE (three-letter code: DND) (formula:  $C_{21}H_{27}N_6O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	6	15	2		
3	B	1	Total	C	N	O	P	0	0
			44	21	6	15	2		
3	C	1	Total	C	N	O	P	0	0
			44	21	6	15	2		
3	D	1	Total	C	N	O	P	0	0
			44	21	6	15	2		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	96	Total	O	0	0
			96	96		
5	B	92	Total	O	0	0
			92	92		
5	C	73	Total	O	0	0
			73	73		
5	D	88	Total	O	0	0
			88	88		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutamine-dependent NAD(+) synthetase




- Molecule 1: Glutamine-dependent NAD(+) synthetase





- Molecule 1: Glutamine-dependent NAD(+) synthetase





● Molecule 1: Glutamine-dependent NAD(+) synthetase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	178.17Å 178.17Å 214.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.09 – 2.73 40.09 – 2.73	Depositor EDS
% Data completeness (in resolution range)	96.3 (40.09-2.73) 91.5 (40.09-2.73)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.17	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.80 (at 2.73Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, $R_{free}$	0.163 , 0.212 0.155 , 0.204	Depositor DCC
$R_{free}$ test set	4400 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.4	Xtriage
Anisotropy	1.061	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 41.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	20820	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, DND, APC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.61	0/5222	0.73	2/7099 (0.0%)
1	B	0.62	0/5183	0.72	2/7049 (0.0%)
1	C	0.61	0/5108	0.71	2/6941 (0.0%)
1	D	0.62	0/5118	0.73	2/6955 (0.0%)
All	All	0.61	0/20631	0.72	8/28044 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	D	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	294	ARG	NE-CZ-NH2	-6.44	117.08	120.30
1	C	294	ARG	NE-CZ-NH2	-6.28	117.16	120.30
1	B	294	ARG	NE-CZ-NH2	-5.82	117.39	120.30
1	D	213	ARG	NE-CZ-NH1	-5.73	117.43	120.30
1	A	69	LEU	CA-CB-CG	-5.69	102.22	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	452	VAL	Peptide
1	D	452	VAL	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5102	0	4976	69	1
1	B	5063	0	4930	69	0
1	C	4992	0	4868	75	0
1	D	5002	0	4878	74	1
2	A	31	0	14	2	0
2	B	31	0	14	2	0
2	C	31	0	14	3	0
2	D	31	0	14	1	0
3	A	44	0	25	4	0
3	B	44	0	25	1	0
3	C	44	0	25	2	0
3	D	44	0	25	2	0
4	B	6	0	8	1	0
4	D	6	0	8	0	0
5	A	96	0	0	6	0
5	B	92	0	0	2	0
5	C	73	0	0	2	0
5	D	88	0	0	4	0
All	All	20820	0	19824	276	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 276 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:98:ARG:HH11	1:D:98:ARG:HG2	1.29	0.96
1:D:633:GLN:OE1	1:D:636:ARG:NH1	2.04	0.90
1:B:98:ARG:HH11	1:B:98:ARG:HG2	1.37	0.90
1:A:98:ARG:HG2	1:A:98:ARG:HH11	1.35	0.88

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:633:GLN:OE1	1:A:636:ARG:NH1	2.06	0.88

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:GLU:OE2	1:D:297:ARG:NH2[8_555]	2.15	0.05

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	655/680 (96%)	631 (96%)	24 (4%)	0	100	100
1	B	652/680 (96%)	626 (96%)	25 (4%)	1 (0%)	47	69
1	C	642/680 (94%)	618 (96%)	23 (4%)	1 (0%)	47	69
1	D	642/680 (94%)	621 (97%)	20 (3%)	1 (0%)	47	69
All	All	2591/2720 (95%)	2496 (96%)	92 (4%)	3 (0%)	51	75

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	177	GLU
1	D	177	GLU
1	B	177	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	525/548 (96%)	506 (96%)	19 (4%)	35	55
1	B	517/548 (94%)	505 (98%)	12 (2%)	50	70
1	C	509/548 (93%)	496 (97%)	13 (3%)	46	66
1	D	512/548 (93%)	496 (97%)	16 (3%)	40	60
All	All	2063/2192 (94%)	2003 (97%)	60 (3%)	42	62

5 of 60 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	468	ARG
1	C	143	ARG
1	D	468	ARG
1	B	664	SER
1	C	257	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	514	GLN
1	B	514	GLN
1	B	515	HIS
1	C	345	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	GOL	B	682	-	5,5,5	0.40	0	5,5,5	0.17	0
3	DND	C	681	-	40,48,48	1.04	2 (5%)	47,73,73	1.28	4 (8%)
2	APC	A	680	-	27,33,33	1.33	3 (11%)	31,52,52	1.44	4 (12%)
4	GOL	D	682	-	5,5,5	0.78	0	5,5,5	0.42	0
2	APC	C	680	-	27,33,33	1.45	4 (14%)	31,52,52	1.57	8 (25%)
3	DND	D	681	-	40,48,48	1.05	2 (5%)	47,73,73	1.27	3 (6%)
3	DND	B	681	-	40,48,48	1.19	3 (7%)	47,73,73	1.21	2 (4%)
3	DND	A	681	-	40,48,48	1.11	3 (7%)	47,73,73	1.26	3 (6%)
2	APC	B	680	-	27,33,33	1.51	5 (18%)	31,52,52	1.37	3 (9%)
2	APC	D	680	-	27,33,33	1.52	3 (11%)	31,52,52	1.48	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	682	-	-	4/4/4/4	-
3	DND	C	681	-	-	8/22/62/62	0/5/5/5
2	APC	A	680	-	-	4/15/38/38	0/3/3/3
4	GOL	D	682	-	-	0/4/4/4	-
2	APC	C	680	-	-	6/15/38/38	0/3/3/3
3	DND	D	681	-	-	10/22/62/62	0/5/5/5
3	DND	B	681	-	-	8/22/62/62	0/5/5/5
3	DND	A	681	-	-	8/22/62/62	0/5/5/5
2	APC	B	680	-	-	10/15/38/38	0/3/3/3
2	APC	D	680	-	-	7/15/38/38	0/3/3/3

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	681	DND	C3N-C7N	5.45	1.52	1.47
3	D	681	DND	C3N-C7N	4.60	1.51	1.47
3	A	681	DND	C3N-C7N	4.50	1.51	1.47
2	D	680	APC	PB-O3B	4.40	1.63	1.58
2	C	680	APC	PB-O3B	4.30	1.63	1.58

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	681	DND	N3A-C2A-N1A	-5.66	119.84	128.68
3	A	681	DND	N3A-C2A-N1A	-4.98	120.89	128.68
3	C	681	DND	N3A-C2A-N1A	-4.89	121.03	128.68
3	B	681	DND	N3A-C2A-N1A	-4.77	121.22	128.68
2	A	680	APC	N3-C2-N1	-4.12	122.24	128.68

There are no chirality outliers.

5 of 65 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	682	GOL	O1-C1-C2-C3
3	C	681	DND	C5D-O5D-PN-O3P
3	C	681	DND	PN-O3P-PA-O5B
3	C	681	DND	C3D-C4D-C5D-O5D
2	A	680	APC	PB-C3A-PA-O1A


There are no ring outliers.

9 monomers are involved in 17 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	682	GOL	1	0
3	C	681	DND	2	0
2	A	680	APC	2	0
2	C	680	APC	3	0
3	D	681	DND	2	0
3	B	681	DND	1	0
3	A	681	DND	4	0
2	B	680	APC	2	0
2	D	680	APC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In


addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## Ligand APC C 680




## Ligand DND D 681









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	661/680 (97%)	-0.54	5 (0%) 86 89	13, 30, 56, 87	0
1	B	658/680 (96%)	-0.56	3 (0%) 91 93	14, 30, 55, 87	0
1	C	650/680 (95%)	-0.50	8 (1%) 79 83	17, 32, 56, 89	0
1	D	650/680 (95%)	-0.52	13 (2%) 65 72	14, 30, 55, 90	0
All	All	2619/2720 (96%)	-0.53	29 (1%) 80 85	13, 31, 56, 90	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	0	SER	5.4
1	C	0	SER	3.9
1	D	441	HIS	3.5
1	C	454	PHE	3.3
1	B	0	SER	3.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	D	682	6/6	0.67	0.35	36,68,74,81	0
2	APC	D	680	31/31	0.92	0.16	41,63,114,119	0
2	APC	C	680	31/31	0.94	0.13	37,61,103,140	0
3	DND	B	681	44/44	0.94	0.20	27,79,151,156	0
3	DND	C	681	44/44	0.94	0.23	39,73,144,154	0
3	DND	D	681	44/44	0.95	0.21	27,65,132,135	0
4	GOL	B	682	6/6	0.95	0.18	38,48,57,60	0
3	DND	A	681	44/44	0.95	0.19	31,59,121,140	0
2	APC	B	680	31/31	0.95	0.14	38,59,112,160	0
2	APC	A	680	31/31	0.95	0.14	32,59,102,161	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around APC C 680:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around DND B 681:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around DND C 681:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around DND D 681:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around DND A 681:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around APC B 680:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.