



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 21, 2022 – 02:23 PM EDT

PDB ID : 7SFE  
Title : Human DNMT1(729-1600) Bound to Zebularine-Containing 12mer dsDNA and Inhibitor GSK3830334A  
Authors : Horton, J.R.; Pathuri, S.; Cheng, X.  
Deposited on : 2021-10-03  
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.27  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	874	<div> <div>7%</div> <div>87%</div> <div>9%</div> <div>.</div> </div>
2	C	12	<div> <div>17%</div> <div>67%</div> <div>33%</div> </div>
3	D	12	<div> <div>33%</div> <div>83%</div> <div>17%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	A	1707	-	-	-	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 7338 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA (cytosine-5)-methyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	842	Total	C	N	O	S	0	10	0
			6560	4164	1144	1209	43			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	727	HIS	-	expression tag	UNP P26358
A	728	MET	-	expression tag	UNP P26358

- Molecule 2 is a DNA chain called DNA (12-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	12	Total	C	N	O	P	0	0	0
			244	116	47	70	11			

- Molecule 3 is a DNA chain called DNA (12-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	12	Total	C	N	O	P	0	0	0
			243	115	46	71	11			

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Zn	0	0
			2	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is (2R)-2-{[6-(4-aminopiperidin-1-yl)-3,5-dicyano-4-ethylpyridin-2-yl]amino}-2-phenylacetamide (three-letter code: 96I) (formula: C<sub>22</sub>H<sub>25</sub>N<sub>7</sub>O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	D	1	Total	C	N	O	0	0
			30	22	7	1		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	223	Total 223	O 223	0	0
8	C	1	Total 1	O 1	0	0
8	D	1	Total 1	O 1	0	0



- Molecule 1: DNA (cytosine-5)-methyltransferase 1





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	161.97Å 78.49Å 117.56Å 90.00° 125.83° 90.00°	Depositor
Resolution (Å)	42.13 – 2.55 42.13 – 2.55	Depositor EDS
% Data completeness (in resolution range)	97.6 (42.13-2.55) 97.6 (42.13-2.55)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.60 (at 2.54Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.201 , 0.236 0.199 , 0.234	Depositor DCC
$R_{free}$ test set	1908 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	54.1	Xtriage
Anisotropy	0.138	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 41.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7338	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 96I, ZN, 5CM, GOL, EDO, PYO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.25	0/6755	0.49	0/9185
2	C	0.52	0/250	0.83	0/382
3	D	0.51	0/250	0.83	0/382
All	All	0.28	0/7255	0.53	0/9949

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6560	0	6252	41	0
2	C	244	0	137	2	0
3	D	243	0	134	1	0
4	A	28	0	42	2	0
5	A	2	0	0	0	0
6	A	6	0	8	0	0
7	D	30	0	0	0	0
8	A	223	0	0	0	0
8	C	1	0	0	0	0
8	D	1	0	0	0	0
All	All	7338	0	6573	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1263:PHE:HB3	1:A:1317:ALA:HB3	1.75	0.69
1:A:919:SER:HB2	1:A:920:ARG:HD3	1.83	0.60
1:A:921:VAL:HG23	1:A:1003:LYS:HG2	1.84	0.59
1:A:1154:GLY:HA3	1:A:1585:ALA:HB3	1.87	0.56
1:A:1330:PRO:HD2	1:A:1356:ARG:HB2	1.86	0.56
1:A:1500:LEU:HB2	1:A:1501:PRO:HD3	1.90	0.53
1:A:922:LEU:HD23	1:A:1000:PHE:HB3	1.89	0.53
1:A:731:ILE:HG23	1:A:752:ILE:HG12	1.91	0.52
1:A:1068:LEU:HD23	1:A:1070:GLU:HG2	1.92	0.52
1:A:1233:ASN:O	1:A:1275:LYS:HD3	2.11	0.50
1:A:1385:GLU:OE2	1:A:1410[A]:ARG:NH1	2.44	0.50
1:A:731:ILE:HG12	1:A:752:ILE:HG23	1.94	0.50
1:A:1311[B]:ARG:NH2	3:D:17:DG:OP1	2.46	0.49
1:A:752:ILE:HD11	1:A:825:LEU:HG	1.95	0.49
1:A:1244:LYS:HA	1:A:1249:VAL:HG21	1.93	0.49
1:A:764:SER:HB2	1:A:774:LEU:HD12	1.94	0.48
1:A:866:GLN:HG3	1:A:867:LEU:HG	1.96	0.48
1:A:1225:PRO:HG2	1:A:1247:LEU:HD12	1.95	0.48
1:A:1252:LEU:HD21	1:A:1283:THR:HG23	1.94	0.48
1:A:1068:LEU:HD21	1:A:1075:TYR:HB2	1.95	0.48
1:A:925:SER:HB2	1:A:932:LEU:HD11	1.96	0.48
1:A:777:ALA:HB2	1:A:796:TRP:CE3	2.49	0.47
1:A:1188:THR:OG1	1:A:1207:ARG:NH1	2.44	0.47
1:A:813:GLU:OE1	4:A:1705:EDO:O2	2.34	0.45
1:A:1145:PHE:HD1	1:A:1169:MET:HE1	1.82	0.45
1:A:903:ARG:O	1:A:907:ILE:HG12	2.17	0.45
1:A:966:GLU:OE1	1:A:1468:SER:N	2.47	0.45
1:A:763:VAL:HG11	1:A:828:ILE:HD13	1.99	0.44
1:A:1382:SER:HA	4:A:1709:EDO:H11	2.00	0.43
1:A:1477:SER:OG	1:A:1484:ALA:O	2.30	0.43
1:A:1233:ASN:OD1	1:A:1233:ASN:N	2.50	0.43
1:A:1443:GLU:HA	1:A:1452:ALA:O	2.18	0.43
1:A:1224:PRO:O	1:A:1267:ASN:ND2	2.50	0.42
1:A:874:ALA:HB2	1:A:1350:PHE:CD1	2.54	0.42
1:A:776:LEU:HD22	1:A:832:VAL:HG21	2.02	0.42
1:A:744:LYS:HB3	1:A:744:LYS:HE2	1.88	0.41
1:A:1142:LEU:HD13	1:A:1165:TRP:HB2	2.02	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1039[A]:ILE:HG12	1:A:1395:TRP:CE2	2.56	0.41
2:C:2:DA:H2''	2:C:3:DG:C8	2.56	0.41
1:A:836:TYR:CE2	1:A:838:ALA:HB2	2.56	0.40
1:A:785:ASP:HB3	1:A:789:GLY:H	1.86	0.40
2:C:3:DG:H2''	2:C:4:DG:C8	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	848/874 (97%)	826 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	685/753 (91%)	676 (99%)	9 (1%)	69	80

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	885	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	920	ARG
1	A	1041	LEU
1	A	1052	ASP
1	A	1068	LEU
1	A	1136	LEU
1	A	1269	ARG
1	A	1479	VAL
1	A	1536	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PYO	D	18	3,2	14,20,21	0.52	0	16,28,31	0.95	1 (6%)
2	5CM	C	6	3,2	15,21,22	0.70	1 (6%)	19,30,33	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PYO	D	18	3,2	-	3/5/25/26	0/2/2/2
2	5CM	C	6	3,2	-	1/4/21/22	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	6	5CM	C4-N3	2.07	1.38	1.35

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	18	PYO	C4-N3-C2	2.49	115.65	114.42

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	6	5CM	O4'-C1'-N1-C6
3	D	18	PYO	O4'-C4'-C5'-O5'
3	D	18	PYO	C3'-C4'-C5'-O5'
3	D	18	PYO	C4'-C5'-O5'-P

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 2 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	EDO	A	1706	-	3,3,3	0.47	0	2,2,2	0.31	0
4	EDO	A	1707	-	3,3,3	0.47	0	2,2,2	0.33	0
4	EDO	A	1708	-	3,3,3	0.45	0	2,2,2	0.37	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	EDO	A	1709	-	3,3,3	0.46	0	2,2,2	0.32	0
7	96I	D	101	-	32,32,32	0.51	0	35,44,44	0.78	0
4	EDO	A	1705	-	3,3,3	0.45	0	2,2,2	0.37	0
6	GOL	A	1710	-	5,5,5	0.94	0	5,5,5	0.96	0
4	EDO	A	1701	-	3,3,3	0.45	0	2,2,2	0.35	0
4	EDO	A	1704	-	3,3,3	0.46	0	2,2,2	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	1706	-	-	0/1/1/1	-
4	EDO	A	1707	-	-	0/1/1/1	-
4	EDO	A	1708	-	-	0/1/1/1	-
4	EDO	A	1709	-	-	0/1/1/1	-
7	96I	D	101	-	-	0/20/32/32	0/3/3/3
4	EDO	A	1705	-	-	0/1/1/1	-
6	GOL	A	1710	-	-	2/4/4/4	-
4	EDO	A	1701	-	-	0/1/1/1	-
4	EDO	A	1704	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1710	GOL	O1-C1-C2-C3
6	A	1710	GOL	O1-C1-C2-O2


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1709	EDO	1	0
4	A	1705	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.



## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	842/874 (96%)	0.39	63 (7%) 14 17	37, 61, 122, 202	0
2	C	11/12 (91%)	0.84	2 (18%) 1 1	84, 109, 192, 229	0
3	D	11/12 (91%)	1.30	4 (36%) 0 0	101, 107, 171, 203	0
All	All	864/898 (96%)	0.41	69 (7%) 12 15	37, 63, 126, 229	0

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	737	ALA	13.1
1	A	956	VAL	10.6
1	A	960	ARG	7.7
1	A	959	PRO	7.6
1	A	746	TYR	7.0
1	A	771	SER	5.7
1	A	957	LYS	5.6
1	A	958	ARG	5.4
1	A	738	VAL	5.4
1	A	788	ASN	5.1
1	A	885	THR	5.1
1	A	747	TYR	5.0
1	A	742	GLY	4.8
1	A	955	PRO	4.6
1	A	884	PRO	4.5
1	A	887	ASP	4.3
1	A	890	PHE	4.3
1	A	954	SER	4.2
1	A	883	GLN	4.1
1	A	740	THR	4.1
3	D	24	DC	4.0
1	A	886	GLU	3.8
1	A	888	ASN	3.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	735	GLY	3.7
1	A	736	GLU	3.7
1	A	786	SER	3.6
2	C	1	DG	3.6
1	A	741	ASP	3.5
3	D	23	DT	3.5
1	A	893	CYS	3.5
1	A	1312[A]	ARG	3.3
1	A	744	LYS	3.3
1	A	1588	ILE	3.2
3	D	22	DC	3.1
1	A	787	SER	3.0
1	A	1306	VAL	2.9
1	A	1481	ALA	2.9
1	A	1584	LEU	2.9
1	A	1299	LEU	2.9
1	A	1482	GLY	2.9
1	A	882	THR	2.8
1	A	1314	ILE	2.7
1	A	1304	TYR	2.6
1	A	862	THR	2.6
1	A	1069	PRO	2.5
1	A	1580	VAL	2.5
1	A	894	VAL	2.5
3	D	21	DC	2.5
1	A	892	PHE	2.4
1	A	976	TYR	2.4
1	A	745	SER	2.4
1	A	860	GLY	2.4
1	A	1583	PRO	2.3
1	A	792	PHE	2.3
1	A	1579	ALA	2.3
1	A	1313	ALA	2.3
1	A	1264	LEU	2.2
1	A	1587	ALA	2.2
2	C	8	DC	2.2
1	A	1526	THR	2.2
1	A	904	GLN	2.1
1	A	1582	PRO	2.1
1	A	854	LEU	2.1
1	A	1556	SER	2.1
1	A	1103	ALA	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	1064	TYR	2.0
1	A	1328	PRO	2.0
1	A	953	SER	2.0
1	A	1557	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	PYO	D	18	19/20	0.81	0.19	96,110,123,140	0
2	5CM	C	6	20/21	0.94	0.15	70,76,84,88	0

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	EDO	A	1707	4/4	0.58	0.47	73,73,73,77	0
4	EDO	A	1706	4/4	0.62	0.32	70,70,73,76	0
6	GOL	A	1710	6/6	0.79	0.29	46,59,61,68	0
4	EDO	A	1701	4/4	0.88	0.10	79,81,82,87	0
4	EDO	A	1709	4/4	0.91	0.45	52,58,58,62	0
4	EDO	A	1708	4/4	0.91	0.18	51,56,58,59	0
4	EDO	A	1704	4/4	0.92	0.35	49,51,52,52	0
7	96I	D	101	30/30	0.93	0.15	77,90,97,98	0
4	EDO	A	1705	4/4	0.94	0.29	63,64,65,68	0
5	ZN	A	1702	1/1	0.97	0.06	75,75,75,75	0
5	ZN	A	1703	1/1	0.98	0.04	84,84,84,84	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.