



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 21, 2022 – 02:29 PM EDT

PDB ID : 7SFG
Title : Human DNMT1(729-1600) Bound to Zebularine-Containing 12mer dsDNA and Cofactor SAM
Authors : Horton, J.R.; Pathuri, S.; Cheng, X.
Deposited on : 2021-10-03
Resolution : 2.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.27
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.27


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1564 (2.46-2.42)
Clashscore	141614	1631 (2.46-2.42)
Ramachandran outliers	138981	1617 (2.46-2.42)
Sidechain outliers	138945	1617 (2.46-2.42)
RSRZ outliers	127900	1547 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	874	<div> <div>5%</div> <div>84%</div> <div>9%</div> <div>7%</div> </div>
2	C	12	<div> <div>100%</div> </div>
3	D	12	<div> <div>8%</div> <div>83%</div> <div>8%</div> <div>8%</div> </div>

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 7130 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA (cytosine-5)-methyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	817	Total	C	N	O	S	0	0	0
			6297	3996	1110	1149	42			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	727	HIS	-	expression tag	UNP P26358
A	728	MET	-	expression tag	UNP P26358

- Molecule 2 is a DNA chain called DNA Strand 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	12	Total	C	N	O	P	0	0	0
			244	116	47	70	11			


- Molecule 3 is a DNA chain called DNA Strand 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	12	Total	C	N	O	P	0	0	0
			243	115	46	71	11			

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Zn	0	0
			2	2		

- Molecule 5 is S-ADENOSYLMETHIONINE (three-letter code: SAM) (formula: C₁₅H₂₂N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			27	15	6	5	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	275	Total	O	0	0
			275	275		
8	C	6	Total	O	0	0
			6	6		
8	D	6	Total	O	0	0
			6	6		

- Molecule 1: DNA (cytosine-5)-methyltransferase 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	161.78Å 77.48Å 116.71Å 90.00° 125.67° 90.00°	Depositor
Resolution (Å)	41.85 – 2.43 41.85 – 2.43	Depositor EDS
% Data completeness (in resolution range)	97.7 (41.85-2.43) 97.7 (41.85-2.43)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.80 (at 2.42Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.180 , 0.223 0.179 , 0.224	Depositor DCC
R_{free} test set	2000 reflections (4.63%)	wwPDB-VP
Wilson B-factor (Å ²)	40.4	Xtriage
Anisotropy	0.174	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7130	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, PYO, GOL, SAM, 5CM, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/6461	0.56	0/8786
2	C	0.70	0/250	0.84	0/382
3	D	0.65	0/250	0.82	0/382
All	All	0.36	0/6961	0.58	0/9550

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6297	0	5959	40	0
2	C	244	0	137	0	0
3	D	243	0	133	3	0
4	A	2	0	0	0	0
5	A	27	0	22	1	0
6	A	6	0	8	0	0
7	A	24	0	36	1	0
8	A	275	0	0	3	0
8	C	6	0	0	0	0
8	D	6	0	0	1	0
All	All	7130	0	6295	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1232:MET:HG3	3:D:17:DG:H2'	1.53	0.89
1:A:1059:ARG:H	7:A:1705:EDO:H11	1.51	0.74
1:A:780:THR:HG21	1:A:893:CYS:HB2	1.76	0.68
1:A:1203:THR:HG22	1:A:1204:THR:HG23	1.80	0.63
1:A:1263:PHE:HB3	1:A:1317:ALA:HB3	1.80	0.63
1:A:1147:GLY:HA3	5:A:1703:SAM:HB1	1.84	0.60
3:D:17:DG:N7	8:D:101:HOH:O	2.33	0.57
1:A:943:PRO:HA	1:A:992:ARG:HG2	1.87	0.57
1:A:913:GLN:HG2	1:A:923:TYR:CE2	2.43	0.54
1:A:1536:GLN:NE2	8:A:1802:HOH:O	2.23	0.52
1:A:744:LYS:HA	1:A:785:ASP:HA	1.92	0.50
3:D:17:DG:H2''	3:D:18:PYO:OP1	2.11	0.50
1:A:1090:ASN:ND2	1:A:1093:SER:HB3	2.26	0.49
1:A:1448:ASP:HB3	1:A:1450:THR:H	1.78	0.49
1:A:1008:ARG:HG3	1:A:1009:PRO:HD2	1.95	0.49
1:A:1187:PHE:HB3	1:A:1189:GLU:HG3	1.94	0.48
1:A:1441:ASN:O	1:A:1454:LYS:HE3	2.12	0.48
1:A:780:THR:HB	1:A:793:HIS:HB3	1.96	0.48
1:A:777:ALA:HB2	1:A:796:TRP:CE3	2.50	0.47
1:A:767:PRO:HB2	1:A:769:ASP:O	2.15	0.47
1:A:1168:GLU:O	1:A:1188:THR:HA	2.15	0.47
1:A:995:ARG:NH2	1:A:1062:VAL:O	2.48	0.46
1:A:793:HIS:ND1	1:A:822:ASP:OD1	2.47	0.46
1:A:1171:ASP:OD2	8:A:1801:HOH:O	2.21	0.45
1:A:1464:ASN:OD1	1:A:1505:ASN:ND2	2.44	0.44
1:A:861:LYS:N	8:A:1819:HOH:O	2.50	0.44
1:A:991:TYR:OH	1:A:1356:ARG:HG2	2.18	0.44
1:A:1015:LYS:NZ	1:A:1052:ASP:OD1	2.48	0.44
1:A:874:ALA:HB2	1:A:1350:PHE:CD1	2.54	0.43
1:A:847:GLY:HA3	1:A:1256:ASP:O	2.19	0.43
1:A:1298:VAL:HG12	1:A:1341:LEU:HD22	2.00	0.43
1:A:746:TYR:HA	1:A:783:TRP:HB3	2.00	0.43
1:A:922:LEU:HD23	1:A:1000:PHE:HB3	2.01	0.42
1:A:1417:MET:CE	1:A:1547:VAL:HG21	2.49	0.42
1:A:1170:TRP:CD1	1:A:1172:PRO:HD2	2.54	0.42
1:A:914:LEU:HD13	1:A:924:TYR:HB2	2.01	0.41
1:A:1459:HIS:CD2	1:A:1477:SER:HA	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1189:GLU:OE2	1:A:1205:ASN:ND2	2.51	0.41
1:A:940:TYR:CZ	1:A:1361:PRO:HB3	2.56	0.41
1:A:1454:LYS:HE2	1:A:1456:ARG:HH12	1.86	0.41
1:A:1415:LYS:HB2	1:A:1417:MET:HE1	2.03	0.40
1:A:839:PRO:HG3	1:A:1320:PRO:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	807/874 (92%)	768 (95%)	39 (5%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	647/753 (86%)	633 (98%)	14 (2%)	52 64

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	740	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	751	CYS
1	A	759	VAL
1	A	787	SER
1	A	837	LYS
1	A	1012	THR
1	A	1045	SER
1	A	1175	GLN
1	A	1227	GLN
1	A	1240	TYR
1	A	1242	LYS
1	A	1444	VAL
1	A	1448	ASP
1	A	1479	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PYO	D	18	1,3	14,20,21	0.50	0	16,28,31	0.95	1 (6%)
2	5CM	C	6	2,3	15,21,22	0.54	0	19,30,33	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PYO	D	18	1,3	-	2/5/25/26	0/2/2/2
2	5CM	C	6	2,3	-	3/4/21/22	0/2/2/2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	18	PYO	C4-N3-C2	2.50	115.65	114.42

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	6	5CM	O4'-C1'-N1-C6
3	D	18	PYO	O4'-C1'-N1-C6
3	D	18	PYO	C2'-C1'-N1-C6
2	C	6	5CM	C3'-C4'-C5'-O5'
2	C	6	5CM	O4'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	18	PYO	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SAM	A	1703	-	21,29,29	0.65	0	18,42,42	1.05	0
7	EDO	A	1710	-	3,3,3	0.50	0	2,2,2	0.51	0
7	EDO	A	1705	-	3,3,3	0.53	0	2,2,2	0.34	0
7	EDO	A	1706	-	3,3,3	0.51	0	2,2,2	0.44	0
7	EDO	A	1707	-	3,3,3	0.55	0	2,2,2	0.03	0
7	EDO	A	1708	-	3,3,3	0.49	0	2,2,2	0.32	0
6	GOL	A	1704	-	5,5,5	1.19	1 (20%)	5,5,5	0.88	0
7	EDO	A	1709	-	3,3,3	0.57	0	2,2,2	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SAM	A	1703	-	-	4/8/33/33	0/3/3/3
7	EDO	A	1710	-	-	1/1/1/1	-
7	EDO	A	1705	-	-	0/1/1/1	-
7	EDO	A	1706	-	-	1/1/1/1	-
7	EDO	A	1707	-	-	0/1/1/1	-
7	EDO	A	1708	-	-	0/1/1/1	-
6	GOL	A	1704	-	-	2/4/4/4	-
7	EDO	A	1709	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1704	GOL	C1-C2	2.32	1.61	1.51

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1703	SAM	N-CA-CB-CG
5	A	1703	SAM	C-CA-CB-CG
5	A	1703	SAM	CB-CG-SD-CE
5	A	1703	SAM	CB-CG-SD-C5'
6	A	1704	GOL	O1-C1-C2-C3
6	A	1704	GOL	O1-C1-C2-O2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
7	A	1706	EDO	O1-C1-C2-O2
7	A	1709	EDO	O1-C1-C2-O2
7	A	1710	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1703	SAM	1	0
7	A	1705	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	817/874 (93%)	0.06	41 (5%) 28 26	24, 48, 101, 146	0
2	C	11/12 (91%)	-0.25	0 100 100	70, 83, 108, 108	0
3	D	11/12 (91%)	-0.04	1 (9%) 9 6	52, 70, 105, 107	0
All	All	839/898 (93%)	0.06	42 (5%) 28 26	24, 49, 101, 146	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1489	ALA	6.3
1	A	1600	ALA	5.1
1	A	737	ALA	4.9
1	A	746	TYR	4.7
1	A	747	TYR	4.3
1	A	1241	SER	4.2
1	A	1488	ALA	3.9
1	A	740	THR	3.8
1	A	977	SER	3.5
1	A	738	VAL	3.4
1	A	1069	PRO	3.2
1	A	1480	GLU	3.0
1	A	979	TYR	3.0
1	A	884	PRO	2.9
1	A	1240	TYR	2.9
1	A	742	GLY	2.8
1	A	788	ASN	2.7
1	A	1449	GLY	2.7
1	A	1482	GLY	2.6
1	A	978	ASP	2.6
1	A	892	PHE	2.6
3	D	24	DC	2.6
1	A	1479	VAL	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	976	TYR	2.5
1	A	1239	THR	2.5
1	A	1301	ALA	2.5
1	A	980	ILE	2.4
1	A	1485	CYS	2.4
1	A	1483	LYS	2.3
1	A	1306	VAL	2.3
1	A	1526	THR	2.3
1	A	787	SER	2.2
1	A	1460	HIS	2.2
1	A	1553	CYS	2.1
1	A	1458	THR	2.1
1	A	781	ALA	2.1
1	A	1071	CYS	2.1
1	A	1582	PRO	2.1
1	A	1583	PRO	2.1
1	A	1484	ALA	2.0
1	A	1307	ALA	2.0
1	A	849	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	5CM	C	6	20/21	0.96	0.11	62,65,74,77	0
3	PYO	D	18	19/20	0.98	0.20	36,40,43,44	0

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	EDO	A	1707	4/4	0.82	0.29	55,56,56,57	0
7	EDO	A	1709	4/4	0.82	0.22	42,48,48,49	0
6	GOL	A	1704	6/6	0.83	0.29	37,40,43,46	0
4	ZN	A	1701	1/1	0.92	0.09	78,78,78,78	0
7	EDO	A	1708	4/4	0.93	0.35	55,55,55,57	0
7	EDO	A	1705	4/4	0.93	0.13	41,43,47,52	0
7	EDO	A	1710	4/4	0.93	0.37	42,42,45,48	0
5	SAM	A	1703	27/27	0.95	0.12	31,37,42,50	0
7	EDO	A	1706	4/4	0.96	0.10	47,48,48,48	0
4	ZN	A	1702	1/1	0.97	0.06	60,60,60,60	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.