



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 06:43 pm BST

PDB ID : 3SI5
Title : Kinetochore-BUBR1 kinase complex
Authors : Blundell, T.L.; Chirgadze, D.Y.; Bolanos-Garcia, V.M.
Deposited on : 2011-06-17
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	176	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="display: flex; justify-content: space-between; align-items: center;"> <div style="width: 80%; height: 10px; background-color: green;"></div> <div style="width: 7%; height: 10px; background-color: yellow;"></div> <div style="width: 14%; height: 10px; background-color: grey;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> 80% 7% 14% </div> </div>
1	B	176	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="display: flex; justify-content: space-between; align-items: center;"> <div style="width: 80%; height: 10px; background-color: green;"></div> <div style="width: 5%; height: 10px; background-color: yellow;"></div> <div style="width: 15%; height: 10px; background-color: grey;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> 80% 5% 15% </div> </div>
2	X	24	<div> <div style="width: 21%; height: 10px; background-color: red;"></div> <div style="display: flex; justify-content: space-between; align-items: center;"> <div style="width: 63%; height: 10px; background-color: green;"></div> <div style="width: 13%; height: 10px; background-color: yellow;"></div> <div style="width: 25%; height: 10px; background-color: grey;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> 21% 63% 13% 25% </div> </div>
2	Y	24	<div> <div style="width: 4%; height: 10px; background-color: red;"></div> <div style="display: flex; justify-content: space-between; align-items: center;"> <div style="width: 50%; height: 10px; background-color: green;"></div> <div style="width: 46%; height: 10px; background-color: grey;"></div> </div> <div style="display: flex; justify-content: space-between; align-items: center;"> 4% 50% 46% </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3044 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitotic checkpoint serine/threonine-protein kinase BUB1 beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	152	Total	C	N	O	S	0	3	0
			1287	814	228	242	3			
1	B	149	Total	C	N	O	S	0	3	0
			1265	799	228	235	3			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	45	THR	-	EXPRESSION TAG	UNP O60566
A	46	GLY	-	EXPRESSION TAG	UNP O60566
A	47	SER	-	EXPRESSION TAG	UNP O60566
A	48	THR	-	EXPRESSION TAG	UNP O60566
A	49	GLY	-	EXPRESSION TAG	UNP O60566
A	50	SER	-	EXPRESSION TAG	UNP O60566
A	51	THR	-	EXPRESSION TAG	UNP O60566
A	52	GLY	-	EXPRESSION TAG	UNP O60566
A	53	SER	-	EXPRESSION TAG	UNP O60566
A	54	THR	-	EXPRESSION TAG	UNP O60566
A	55	GLY	-	EXPRESSION TAG	UNP O60566
A	56	SER	-	EXPRESSION TAG	UNP O60566
B	45	THR	-	EXPRESSION TAG	UNP O60566
B	46	GLY	-	EXPRESSION TAG	UNP O60566
B	47	SER	-	EXPRESSION TAG	UNP O60566
B	48	THR	-	EXPRESSION TAG	UNP O60566
B	49	GLY	-	EXPRESSION TAG	UNP O60566
B	50	SER	-	EXPRESSION TAG	UNP O60566
B	51	THR	-	EXPRESSION TAG	UNP O60566
B	52	GLY	-	EXPRESSION TAG	UNP O60566
B	53	SER	-	EXPRESSION TAG	UNP O60566
B	54	THR	-	EXPRESSION TAG	UNP O60566
B	55	GLY	-	EXPRESSION TAG	UNP O60566
B	56	SER	-	EXPRESSION TAG	UNP O60566

- Molecule 2 is a protein called Protein CASC5.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	X	18	Total	C	N	O	0	0	0
			143	91	26	26			
2	Y	13	Total	C	N	O	0	0	0
			99	66	15	18			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	203	GLY	-	EXPRESSION TAG	UNP Q8NG31
X	204	PRO	-	EXPRESSION TAG	UNP Q8NG31
X	205	LEU	-	EXPRESSION TAG	UNP Q8NG31
X	206	GLY	-	EXPRESSION TAG	UNP Q8NG31
X	207	SER	-	EXPRESSION TAG	UNP Q8NG31
Y	203	GLY	-	EXPRESSION TAG	UNP Q8NG31
Y	204	PRO	-	EXPRESSION TAG	UNP Q8NG31
Y	205	LEU	-	EXPRESSION TAG	UNP Q8NG31
Y	206	GLY	-	EXPRESSION TAG	UNP Q8NG31
Y	207	SER	-	EXPRESSION TAG	UNP Q8NG31

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	138	Total	O	0	0
			138	138		
3	B	101	Total	O	0	0
			101	101		
3	X	9	Total	O	0	0
			9	9		
3	Y	2	Total	O	0	0
			2	2		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Mitotic checkpoint serine/threonine-protein kinase BUB1 beta




- Molecule 1: Mitotic checkpoint serine/threonine-protein kinase BUB1 beta



- Molecule 2: Protein CASC5



- Molecule 2: Protein CASC5



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	124.40Å 40.14Å 75.39Å 90.00° 91.38° 90.00°	Depositor
Resolution (Å)	38.20 – 2.20 38.20 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (38.20-2.20) 99.9 (38.20-2.20)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.193 , 0.247 0.205 , 0.253	Depositor DCC
R_{free} test set	985 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	21.7	Xtriage
Anisotropy	0.161	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 34.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.016 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3044	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/1327	0.58	0/1792
1	B	0.55	0/1305	0.58	1/1761 (0.1%)
2	X	0.49	0/144	0.60	0/190
2	Y	0.46	0/100	0.51	0/134
All	All	0.56	0/2876	0.58	1/3877 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	73	ASP	CB-CG-OD1	5.43	123.19	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1287	0	1220	9	0
1	B	1265	0	1198	7	0
2	X	143	0	141	1	0
2	Y	99	0	85	1	0
3	A	138	0	0	2	0
3	B	101	0	0	1	0
3	X	9	0	0	0	0
3	Y	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3044	0	2644	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:160:GLU:HG3	3:A:287:HOH:O	1.89	0.72
1:B:114:ARG:NH1	3:B:234:HOH:O	2.36	0.58
1:B:82:ILE:HD13	1:B:101:LEU:HD22	1.85	0.57
1:B:82:ILE:CD1	1:B:101:LEU:HD22	2.34	0.56
1:A:202:ARG:NH1	3:A:269:HOH:O	2.40	0.51
1:A:95:GLU:CD	1:A:95:GLU:H	2.12	0.51
1:B:78:TRP:O	1:B:82:ILE:HG12	2.15	0.47
1:A:129:GLY:HA3	1:A:138:MET:HE2	1.98	0.45
1:A:123:ASN:OD1	1:A:127:LYS:HE2	2.17	0.44
1:B:99:SER:HB2	2:Y:213:ILE:HD11	1.99	0.44
1:B:162:TYR:HD1	1:B:174:ILE:CD1	2.31	0.43
1:A:58:GLN:HG2	1:A:84:TRP:NE1	2.33	0.43
1:A:75:LEU:HD21	1:A:120:ARG:HB2	2.00	0.43
1:A:95:GLU:N	1:A:95:GLU:OE1	2.34	0.42
1:A:89:TYR:CE2	1:A:95:GLU:HG2	2.54	0.42
1:B:162:TYR:CD1	1:B:174:ILE:CD1	3.03	0.42
2:X:214:ASP:HB3	2:X:217:ASP:HB2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	153/176 (87%)	153 (100%)	0	0	100	100
1	B	150/176 (85%)	146 (97%)	4 (3%)	0	100	100
2	X	16/24 (67%)	15 (94%)	0	1 (6%)	1	0
2	Y	11/24 (46%)	10 (91%)	1 (9%)	0	100	100
All	All	330/400 (82%)	324 (98%)	5 (2%)	1 (0%)	41	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	X	210	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	133/153 (87%)	133 (100%)	0	100	100
1	B	130/153 (85%)	128 (98%)	2 (2%)	65	78
2	X	15/21 (71%)	15 (100%)	0	100	100
2	Y	9/21 (43%)	9 (100%)	0	100	100
All	All	287/348 (82%)	285 (99%)	2 (1%)	84	91

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	101	LEU
1	B	106	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	GLN
1	A	144	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	153	GLN
1	A	197	GLN
1	B	87	GLN
1	B	144	ASN
1	B	197	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	152/176 (86%)	-0.30	1 (0%) 87 86	14, 20, 36, 43	3 (1%)
1	B	149/176 (84%)	-0.22	4 (2%) 54 52	14, 23, 46, 59	2 (1%)
2	X	18/24 (75%)	0.58	5 (27%) 0 0	21, 36, 57, 57	0
2	Y	13/24 (54%)	0.62	1 (7%) 13 12	50, 62, 74, 80	0
All	All	332/400 (83%)	-0.18	11 (3%) 46 44	14, 22, 54, 80	5 (1%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	54	THR	3.4
2	X	224	THR	3.1
2	X	211	ASN	3.0
1	B	94	LYS	2.7
1	B	93	GLY	2.5
2	X	225	GLY	2.4
1	B	90	PRO	2.2
2	X	210	GLU	2.1
2	X	208	SER	2.1
1	B	91	GLN	2.0
2	Y	224	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.