



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 10:03 am BST

PDB ID : 3SM8  
Title : Crystal Structure of Pseudomonas aeruginosa D-Arginine Dehydrogenase in Complex with an (N5) Flavin Adduct  
Authors : Fu, G.; Weber, I.T.  
Deposited on : 2011-06-27  
Resolution : 1.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1386 (1.12-1.04)
Clashscore	141614	1021 (1.10-1.06)
Ramachandran outliers	138981	1381 (1.12-1.04)
Sidechain outliers	138945	1379 (1.12-1.04)
RSRZ outliers	127900	1359 (1.12-1.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	381	<div> <div></div> <div>88%</div> <div>10%</div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3439 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FAD-dependent catabolic D-arginine dehydrogenase, DauA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	381	3013	1907	538	552	16	0	23	0

There are 7 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	995	HIS	-	EXPRESSION TAG	UNP Q9HXE3
A	996	HIS	-	EXPRESSION TAG	UNP Q9HXE3
A	997	HIS	-	EXPRESSION TAG	UNP Q9HXE3
A	998	HIS	-	EXPRESSION TAG	UNP Q9HXE3
A	999	HIS	-	EXPRESSION TAG	UNP Q9HXE3
A	1000	HIS	-	EXPRESSION TAG	UNP Q9HXE3
A	1001	PRO	-	EXPRESSION TAG	UNP Q9HXE3

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is [(2R,3S,4R,5R)-5-(6-amino-9H-purin-9-yl)-3,4-dihydroxytetrahydrofuran-2-yl]methyl (2R,3S,4S)-5-[7,8-dimethyl-5-(3-methylbutanoyl)-2,4-dioxo-1,3,4,5-tetrahydrobenzo[g]pteridin-10(2H)-yl]-2,3,4-trihydroxypentyl dihydrogen diphosphate (three-letter code: FNK) (formula: C<sub>32</sub>H<sub>43</sub>N<sub>9</sub>O<sub>16</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			59	32	9	16	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	349	Total	O	0	2
			349	349		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: FAD-dependent catabolic D-arginine dehydrogenase, DauA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.07Å 77.93Å 89.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.07 44.66 – 1.07	Depositor EDS
% Data completeness (in resolution range)	98.4 (10.00-1.07) 94.7 (44.66-1.07)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.41 (at 1.07Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.145 , 0.165 0.133 , 0.151	Depositor DCC
$R_{free}$ test set	9381 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	7.6	Xtriage
Anisotropy	0.334	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 52.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	3439	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FNK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.85	1/3142 (0.0%)	1.38	35/4283 (0.8%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1173	GLU	CD-OE2	-8.33	1.16	1.25

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1100[A]	ARG	CD-NE-CZ	18.58	149.62	123.60
1	A	1359	ARG	NE-CZ-NH2	13.38	126.99	120.30
1	A	1156	ARG	CD-NE-CZ	12.94	141.72	123.60
1	A	1234[A]	ASP	CB-CG-OD2	12.40	129.46	118.30
1	A	1234[B]	ASP	CB-CG-OD2	12.40	129.46	118.30
1	A	1066	ARG	NE-CZ-NH2	-11.50	114.55	120.30
1	A	1367[A]	MET	CA-CB-CG	11.06	132.10	113.30
1	A	1367[B]	MET	CA-CB-CG	11.06	132.10	113.30
1	A	1191	ARG	NE-CZ-NH1	10.39	125.50	120.30
1	A	1191	ARG	NE-CZ-NH2	-10.36	115.12	120.30
1	A	1156	ARG	NE-CZ-NH1	10.24	125.42	120.30
1	A	1160	ARG	NE-CZ-NH2	-10.23	115.19	120.30
1	A	1176	ARG	NE-CZ-NH1	10.15	125.38	120.30
1	A	1359	ARG	NE-CZ-NH1	-9.45	115.58	120.30
1	A	1160	ARG	NE-CZ-NH1	9.45	125.02	120.30
1	A	1114	ARG	CD-NE-CZ	8.30	135.22	123.60
1	A	1070	ASP	CB-CG-OD2	-8.23	110.90	118.30
1	A	1033	ARG	NE-CZ-NH2	-8.10	116.25	120.30
1	A	1119	GLU	CA-CB-CG	7.48	129.85	113.40
1	A	1372	ARG	NE-CZ-NH1	7.20	123.90	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1066	ARG	CG-CD-NE	6.89	126.27	111.80
1	A	1108	ALA	CB-CA-C	6.21	119.42	110.10
1	A	1233	ILE	O-C-N	-6.15	112.86	122.70
1	A	1305	ARG	NE-CZ-NH2	-6.13	117.24	120.30
1	A	1374	SER	N-CA-CB	-6.09	101.37	110.50
1	A	1160	ARG	CD-NE-CZ	5.93	131.91	123.60
1	A	1175	ARG	NE-CZ-NH1	5.88	123.24	120.30
1	A	1050	THR	CA-CB-CG2	-5.80	104.27	112.40
1	A	1319	ALA	O-C-N	-5.75	113.50	122.70
1	A	1059	ARG	NE-CZ-NH2	-5.61	117.49	120.30
1	A	1305	ARG	NE-CZ-NH1	5.52	123.06	120.30
1	A	1359	ARG	CD-NE-CZ	5.32	131.05	123.60
1	A	1130	ARG	NE-CZ-NH2	5.30	122.95	120.30
1	A	1234[A]	ASP	OD1-CG-OD2	-5.25	113.32	123.30
1	A	1234[B]	ASP	OD1-CG-OD2	-5.25	113.32	123.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3013	0	2952	14	0
2	A	18	0	24	1	0
3	A	59	0	41	0	0
4	A	349	0	0	3	0
All	All	3439	0	3017	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1296:ARG:O	1:A:1296:ARG:HG3	1.70	0.92

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1400:GOL:H32	2:A:1401:GOL:O3	1.86	0.76
1:A:999[B]:HIS:CE1	1:A:1188:GLY:HA3	2.40	0.56
1:A:999[B]:HIS:NE2	1:A:1188:GLY:HA3	2.21	0.56
1:A:1364:ASP:O	1:A:1367[B]:MET:HG3	2.06	0.56
1:A:998:HIS:HD2	4:A:2247:HOH:O	1.91	0.53
1:A:1230:PRO:HD3	4:A:2195:HOH:O	2.08	0.53
1:A:1225:PHE:O	1:A:1258[A]:LEU:HD22	2.09	0.52
1:A:999[B]:HIS:CD2	1:A:1188:GLY:HA3	2.46	0.50
1:A:1273:GLN:NE2	4:A:2299:HOH:O	2.45	0.48
1:A:1009[B]:ILE:CD1	1:A:1195[B]:LEU:HD11	2.44	0.47
1:A:1343:GLU:OE2	1:A:1357:HIS:HE1	1.99	0.45
1:A:1095:ASP:OD1	1:A:1097[A]:GLU:OE2	2.38	0.42
1:A:1295:ARG:HG2	1:A:1296:ARG:HG2	2.03	0.41
1:A:1296:ARG:O	1:A:1296:ARG:CG	2.53	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	398/381 (104%)	390 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	312/293 (106%)	305 (98%)	7 (2%)	52 14

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	995[A]	HIS
1	A	1006	TYR
1	A	1093	SER
1	A	1097[A]	GLU
1	A	1109	LEU
1	A	1296	ARG
1	A	1374	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	998	HIS
1	A	1357	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	A	1401	-	5,5,5	0.79	0	5,5,5	1.94	2 (40%)
2	GOL	A	1402	-	5,5,5	0.70	0	5,5,5	1.07	1 (20%)
2	GOL	A	1400	-	5,5,5	0.73	0	5,5,5	0.56	0
3	FNK	A	1500	-	54,64,64	1.30	9 (16%)	62,98,98	2.07	12 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1401	-	-	2/4/4/4	-
2	GOL	A	1402	-	-	0/4/4/4	-
2	GOL	A	1400	-	-	0/4/4/4	-
3	FNK	A	1500	-	-	1/34/58/58	0/6/6/6

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1500	FNK	C4-N3	3.91	1.39	1.33
3	A	1500	FNK	C10-N1	3.46	1.37	1.33
3	A	1500	FNK	C5X-N5	2.83	1.43	1.38
3	A	1500	FNK	C4X-N5	2.82	1.43	1.38
3	A	1500	FNK	C7M-C7	2.74	1.56	1.51
3	A	1500	FNK	O4B-C1B	2.60	1.44	1.41
3	A	1500	FNK	C9A-N10	2.56	1.42	1.38
3	A	1500	FNK	C2-N1	-2.15	1.33	1.38
3	A	1500	FNK	C2A-N3A	2.03	1.35	1.32

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1500	FNK	C4-N3-C2	9.30	122.99	115.14
3	A	1500	FNK	C1'-N10-C9A	6.28	123.23	118.29
3	A	1500	FNK	C4X-C4-N3	-5.37	116.08	123.43
3	A	1500	FNK	C4X-N5-C5X	-3.68	113.94	122.12
3	A	1500	FNK	C9A-N10-C10	-3.42	117.44	121.91
3	A	1500	FNK	C4-C4X-C10	-3.29	117.78	119.95

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1401	GOL	C3-C2-C1	-3.21	99.24	111.70
3	A	1500	FNK	O4B-C1B-C2B	-3.19	102.26	106.93
3	A	1500	FNK	CG-CB-CA	3.02	117.76	113.32
3	A	1500	FNK	O-CA-CB	-2.79	117.17	122.20
2	A	1401	GOL	O3-C3-C2	-2.50	98.20	110.20
3	A	1500	FNK	C5A-C6A-N6A	2.41	124.02	120.35
2	A	1402	GOL	C3-C2-C1	-2.21	103.10	111.70
3	A	1500	FNK	C10-C4X-N5	2.20	118.52	116.81
3	A	1500	FNK	C6-C5X-C9A	2.13	121.23	119.06

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1401	GOL	O1-C1-C2-C3
2	A	1401	GOL	O1-C1-C2-O2
3	A	1500	FNK	O4B-C4B-C5B-O5B

There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1401	GOL	1	0
2	A	1400	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	381/381 (100%)	-0.24	3 (0%) 86 82	5, 9, 30, 73	4 (1%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1375[A]	PRO	9.9
1	A	995[A]	HIS	5.7
1	A	1295	ARG	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	A	1401	6/6	0.83	0.17	28,35,46,48	0
2	GOL	A	1400	6/6	0.97	0.12	9,14,17,19	0
2	GOL	A	1402	6/6	0.98	0.09	14,15,17,18	0
3	FNK	A	1500	59/59	0.99	0.06	4,5,13,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.