



Full wwPDB X-ray Structure Validation Report ⓘ

May 28, 2020 – 08:29 pm BST

PDB ID : 1SN7
Title : KUMAMOLISIN-AS, APOENZYME
Authors : Wlodawer, A.; Li, M.; Gustchina, A.; Oda, K.; Nishino, T.
Deposited on : 2004-03-10
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	<div> <div></div> <div>75%</div> <div>18%</div> <div>5%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2813 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called kumamolisin-As.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	355	Total	C	N	O	S	0	0	1
			2526	1591	423	508	4			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	286	Total	O	0	0
			286	286		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: kumamolisin-As



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	41.81Å 44.95Å 49.11Å 113.90° 106.10° 102.30°	Depositor
Resolution (Å)	10.00 – 2.00 39.25 – 2.01	Depositor EDS
% Data completeness (in resolution range)	85.5 (10.00-2.00) 89.2 (39.25-2.01)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.70 (at 2.01Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.150 , 0.255 0.148 , 0.236	Depositor DCC
R_{free} test set	892 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	23.5	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 82.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2813	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/2587	1.51	31/3553 (0.9%)

There are no bond length outliers.

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	350	ARG	CD-NE-CZ	16.60	146.84	123.60
1	A	223	ARG	NE-CZ-NH1	14.75	127.67	120.30
1	A	223	ARG	NE-CZ-NH2	-13.76	113.42	120.30
1	A	223	ARG	CD-NE-CZ	11.33	139.47	123.60
1	A	350	ARG	NE-CZ-NH2	-10.72	114.94	120.30
1	A	350	ARG	NE-CZ-NH1	9.03	124.81	120.30
1	A	327	ARG	NE-CZ-NH1	-7.89	116.35	120.30
1	A	204	GLN	CA-CB-CG	7.27	129.39	113.40
1	A	134	ASP	CB-CG-OD2	-7.26	111.76	118.30
1	A	134	ASP	CB-CG-OD1	7.23	124.81	118.30
1	A	252	ASP	CB-CG-OD2	6.75	124.37	118.30
1	A	107	PHE	CB-CG-CD1	6.58	125.41	120.80
1	A	263	TYR	CB-CG-CD1	6.34	124.81	121.00
1	A	252	ASP	CB-CG-OD1	-6.08	112.83	118.30
1	A	194	ARG	CD-NE-CZ	6.07	132.10	123.60
1	A	248	ARG	NE-CZ-NH1	5.95	123.27	120.30
1	A	301	TYR	CB-CG-CD1	5.68	124.41	121.00
1	A	201	ARG	NE-CZ-NH1	5.58	123.09	120.30
1	A	327	ARG	NH1-CZ-NH2	5.55	125.51	119.40
1	A	290	ARG	NE-CZ-NH1	5.41	123.00	120.30
1	A	331	TYR	CB-CG-CD1	5.38	124.23	121.00
1	A	194	ARG	NE-CZ-NH2	-5.35	117.63	120.30
1	A	45	TYR	CB-CG-CD1	-5.28	117.83	121.00
1	A	107	PHE	CB-CG-CD2	-5.21	117.15	120.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	188	LEU	CA-CB-CG	5.18	127.22	115.30
1	A	117	ASP	CB-CG-OD2	5.17	122.96	118.30
1	A	263	TYR	CB-CG-CD2	-5.17	117.90	121.00
1	A	169	ASP	CB-CG-OD1	5.15	122.93	118.30
1	A	98	TYR	CB-CG-CD2	-5.13	117.92	121.00
1	A	169	ASP	CB-CG-OD2	-5.09	113.71	118.30
1	A	109	ASP	CB-CG-OD2	-5.04	113.76	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2526	0	2410	49	0
2	A	1	0	0	0	0
3	A	286	0	0	12	0
All	All	2813	0	2410	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:4:THR:HG22	1:A:261:THR:HA	1.57	0.85
1:A:40:ALA:C	1:A:40:ALA:N	2.37	0.79
1:A:40:ALA:CB	1:A:40:ALA:N	2.46	0.78
1:A:141:ILE:HD11	3:A:643:HOH:O	1.82	0.77
1:A:332:GLN:HB2	3:A:739:HOH:O	1.91	0.71
1:A:40:ALA:C	1:A:40:ALA:CB	2.62	0.68
1:A:44:GLN:HE22	1:A:266:VAL:HG11	1.62	0.65
1:A:4:THR:HG21	3:A:660:HOH:O	1.98	0.64
1:A:39:GLU:O	1:A:40:ALA:CB	2.47	0.62
1:A:347:ILE:HG21	1:A:350:ARG:HD2	1.84	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:THR:O	1:A:121:LYS:HE2	2.04	0.58
1:A:37:TYR:HE1	1:A:39:GLU:OE2	1.87	0.57
1:A:39:GLU:C	1:A:40:ALA:CB	2.75	0.55
1:A:116:HIS:HE1	3:A:635:HOH:O	1.89	0.54
1:A:9:LEU:HD23	1:A:89:LEU:CD1	2.38	0.54
1:A:146:ARG:NH2	1:A:149:LEU:HD22	2.23	0.54
1:A:228:PRO:HG2	1:A:231:GLN:HB2	1.91	0.53
1:A:315:HIS:HD2	3:A:754:HOH:O	1.92	0.52
1:A:180:PHE:CG	1:A:181:PRO:HA	2.45	0.52
1:A:111:ILE:O	1:A:115:ILE:HD12	2.10	0.52
1:A:55:GLN:NE2	1:A:94:LYS:HE2	2.24	0.52
1:A:239:SER:HB3	1:A:244:ALA:HB1	1.92	0.52
1:A:171:GLU:HB2	1:A:177:HIS:CD2	2.45	0.51
1:A:146:ARG:HH21	1:A:149:LEU:HD22	1.76	0.51
1:A:353:GLN:HA	1:A:356:LEU:HD12	1.92	0.50
1:A:4:THR:HG22	1:A:261:THR:CA	2.34	0.49
1:A:133:GLU:HA	3:A:643:HOH:O	2.12	0.47
1:A:136:TRP:HB2	3:A:643:HOH:O	2.13	0.47
1:A:219:GLY:HA3	1:A:339:PRO:O	2.15	0.47
1:A:350:ARG:NE	3:A:519:HOH:O	2.49	0.46
1:A:350:ARG:NH2	3:A:519:HOH:O	2.49	0.46
1:A:113:THR:O	1:A:117:ASP:HB2	2.16	0.45
1:A:330:ILE:HA	1:A:331:TYR:HA	1.81	0.45
1:A:39:GLU:OE1	1:A:56:VAL:HB	2.17	0.45
1:A:62:ASP:HB2	1:A:101:PRO:CD	2.47	0.45
1:A:172:GLN:H	1:A:172:GLN:HG3	1.48	0.44
1:A:201:ARG:NH2	3:A:753:HOH:O	2.50	0.44
1:A:347:ILE:CG2	1:A:350:ARG:HD2	2.47	0.44
1:A:38:ASP:HB3	3:A:772:HOH:O	2.17	0.44
1:A:115:ILE:HG12	1:A:151:ALA:HB2	2.00	0.44
1:A:9:LEU:HD23	1:A:89:LEU:HD12	1.98	0.44
1:A:145:ASN:ND2	1:A:184:SER:OG	2.51	0.43
1:A:239:SER:HB3	1:A:244:ALA:CB	2.48	0.43
1:A:12:ALA:HB1	1:A:17:PHE:CD2	2.54	0.43
1:A:45:TYR:CD2	1:A:84:GLU:HG2	2.54	0.42
1:A:233:HIS:HB3	3:A:735:HOH:O	2.20	0.41
1:A:62:ASP:HB2	1:A:101:PRO:HD2	2.02	0.41
1:A:281:ALA:HB3	1:A:282:PRO:HD3	2.00	0.41
1:A:111:ILE:HD11	1:A:127:ILE:HD13	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/364 (97%)	339 (96%)	13 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/260 (97%)	230 (91%)	23 (9%)	9	5

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	THR
1	A	24	GLN
1	A	39	GLU
1	A	44	GLN
1	A	48	SER
1	A	58	SER
1	A	89	LEU
1	A	107	PHE
1	A	115	ILE
1	A	121	LYS
1	A	145	ASN
1	A	146	ARG
1	A	168	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	172	GLN
1	A	175	LEU
1	A	188	LEU
1	A	201	ARG
1	A	213	ASP
1	A	245	SER
1	A	248	ARG
1	A	251	PRO
1	A	308	GLN
1	A	353	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	GLN
1	A	44	GLN
1	A	55	GLN
1	A	67	GLN
1	A	145	ASN
1	A	172	GLN
1	A	235	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	355/364 (97%)	-0.49	2 (0%) 89 88	14, 25, 45, 69	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	39	GLU	2.6
1	A	40	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CA	A	401	1/1	1.00	0.04	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.