



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 01:37 pm BST

PDB ID : 6SPW
Title : Structure of protein kinase CK2 catalytic subunit with the CK2beta-competitive bisubstrate inhibitor ARC3140
Authors : Niefind, K.; Schnitzler, A.
Deposited on : 2019-09-03
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	391	
2	B	7	
2	C	7	
2	D	7	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	DAS	B	2	-	-	-	X
2	DAS	C	2	-	-	-	X
2	DAS	D	2	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3321 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Casein kinase II subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	327	Total	C	N	O	S	0	3	0
			2787	1787	489	497	14			


- Molecule 2 is a protein called ARC3140.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	7	Total	C	N	O	0	0	0
			58	30	8	20			
2	C	7	Total	C	N	O	0	0	0
			58	30	8	20			
2	D	7	Total	C	N	O	0	0	0
			58	30	8	20			

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is 8-[4,5,6,7-tetrakis(iodanyl)benzimidazol-1-yl]octanoic acid (three-letter code: A0Z) (formula: C₁₅H₁₆I₄N₂O₂) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	I	N	O	0	0
			22	15	4	2	1		
4	C	1	Total	C	I	N	O	0	0
			22	15	4	2	1		
4	D	1	Total	C	I	N	O	0	0
			22	15	4	2	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	286	Total	O	0	0
			286	286		
5	B	2	Total	O	0	0
			2	2		
5	C	3	Total	O	0	0
			3	3		
5	D	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Casein kinase II subunit alpha




• Molecule 2: ARC3140



• Molecule 2: ARC3140



• Molecule 2: ARC3140



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	47.50Å 85.17Å 90.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.59 – 1.60 42.59 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (42.59-1.60) 99.8 (42.59-1.60)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 1.60Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.165 , 0.196 0.165 , 0.196	Depositor DCC
R_{free} test set	1960 reflections (3.99%)	wwPDB-VP
Wilson B-factor (Å ²)	16.4	Xtriage
Anisotropy	0.472	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 55.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3321	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A0Z, NA, DAS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	2/2862 (0.1%)	0.85	1/3869 (0.0%)
2	B	0.39	0/49	0.61	0/63
2	C	0.24	0/49	0.51	0/63
2	D	0.22	0/49	0.47	0/63
All	All	0.83	2/3009 (0.1%)	0.83	1/4058 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	22	GLU	CG-CD	5.28	1.59	1.51
1	A	209	TYR	CE2-CZ	5.25	1.45	1.38

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	312	ARG	NE-CZ-NH2	-5.54	117.53	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2787	0	2733	26	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:ASP:OD1	4:D:101:A0Z:IAX	3.03	0.47
1:A:104:PRO:HA	1:A:107:ARG:HH21	1.80	0.46
1:A:33:TRP:CH2	1:A:102:LYS:HE3	2.51	0.46
1:A:113:PHE:CD1	4:B:101:A0Z:IAZ	3.40	0.45
1:A:8:ARG:NH1	5:A:509:HOH:O	2.47	0.44
4:D:101:A0Z:CBL	4:D:101:A0Z:IAZ	3.36	0.44
1:A:27:GLU:OE1	1:A:76:LYS:HE3	2.17	0.44
1:A:19:ARG:NH2	5:A:502:HOH:O	2.47	0.43
1:A:229:LYS:HE3	5:A:668:HOH:O	2.17	0.42
1:A:196:TYR:CD2	1:A:231:PRO:HG3	2.55	0.42
1:A:194:SER:HA	2:B:3:ASP:HB2	2.01	0.42
1:A:199:GLY:HA2	1:A:216:TRP:CD1	2.55	0.41
4:B:101:A0Z:CBL	4:B:101:A0Z:IAZ	3.39	0.41
1:A:315:ALA:O	1:A:319:MET:HG3	2.21	0.41
4:C:101:A0Z:IAZ	4:C:101:A0Z:CBI	3.39	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:ARG:NH1	1:A:167:GLU:OE1[1_655]	2.05	0.15
5:A:564:HOH:O	5:A:686:HOH:O[4_566]	2.13	0.07
5:A:502:HOH:O	5:A:722:HOH:O[4_566]	2.15	0.05
5:A:744:HOH:O	5:A:751:HOH:O[4_566]	2.15	0.05

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	328/391 (84%)	320 (98%)	7 (2%)	1 (0%)	41	21
2	B	5/7 (71%)	4 (80%)	1 (20%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	5/7 (71%)	3 (60%)	2 (40%)	0	100	100
2	D	5/7 (71%)	5 (100%)	0	0	100	100
All	All	343/412 (83%)	332 (97%)	10 (3%)	1 (0%)	41	21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	193	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	304/347 (88%)	299 (98%)	5 (2%)	62	41
2	B	6/6 (100%)	6 (100%)	0	100	100
2	C	6/6 (100%)	6 (100%)	0	100	100
2	D	6/6 (100%)	4 (67%)	2 (33%)	0	0
All	All	322/365 (88%)	315 (98%)	7 (2%)	52	27

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	107	ARG
1	A	121	PHE
1	A	175	ASP
1	A	180	GLU
1	A	269	PHE
2	D	4	ASP
2	D	8	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:


Mol	Chain	Res	Type	Atoms
4	B	101	A0Z	CBI-CBL-NDB-CBA
4	B	101	A0Z	CBI-CBL-NDB-CCT
4	D	101	A0Z	CBE-CBG-CBJ-CCH
4	B	101	A0Z	CBD-CBE-CBG-CBJ
4	C	101	A0Z	CBD-CBE-CBG-CBJ
4	D	101	A0Z	CBD-CBE-CBG-CBJ
4	D	101	A0Z	CBE-CBD-CBF-CBI
4	C	101	A0Z	CBE-CBD-CBF-CBI
4	D	101	A0Z	CBF-CBD-CBE-CBG
4	C	101	A0Z	CBE-CBG-CBJ-CCH


There are no ring outliers.


3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	101	A0Z	3	0
4	B	101	A0Z	5	0
4	D	101	A0Z	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	327/391 (83%)	0.29	19 (5%) 23 20	9, 18, 46, 69	0
2	B	6/7 (85%)	8.57	6 (100%) 0 0	78, 103, 113, 123	0
2	C	6/7 (85%)	9.26	6 (100%) 0 0	117, 128, 139, 140	0
2	D	6/7 (85%)	12.50	6 (100%) 0 0	120, 137, 138, 142	0
All	All	345/412 (83%)	0.80	37 (10%) 6 5	9, 19, 69, 142	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	5	ASP	17.4
2	B	5	ASP	14.5
2	D	7	ASP	14.2
2	D	3	ASP	12.5
2	D	6	ASP	11.9
2	C	7	ASP	11.9
2	C	4	ASP	11.7
2	C	8	LYS	9.9
2	D	4	ASP	9.6
2	C	6	ASP	9.6
2	D	8	LYS	9.4
2	B	4	ASP	8.6
2	B	6	ASP	8.3
1	A	125	TYR	8.1
2	B	8	LYS	8.1
1	A	72	PRO	8.0
2	B	3	ASP	7.8
2	C	3	ASP	7.6
1	A	50	TYR	7.5
1	A	121	PHE	6.5
1	A	73	VAL	5.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	5	ASP	4.8
1	A	49	LYS	4.3
1	A	74	LYS	4.2
1	A	120	ASP	4.2
2	B	7	ASP	4.1
1	A	104	PRO	4.1
1	A	122	LYS	4.0
1	A	75	LYS	3.2
1	A	71	LYS	3.0
1	A	47	ARG	3.0
1	A	118	ASN	2.9
1	A	124	LEU	2.7
1	A	236	HIS	2.7
1	A	267	PRO	2.4
1	A	119	THR	2.2
1	A	269	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DAS	B	2	8/9	0.31	0.57	55,59,62,65	0
2	DAS	C	2	8/9	0.31	0.40	50,53,57,58	0
2	DAS	D	2	8/9	0.44	0.49	50,57,64,66	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NA	A	401	1/1	0.96	0.12	36,36,36,36	0
4	A0Z	B	101	22/23	0.99	0.12	20,32,61,61	0
4	A0Z	D	101	22/23	0.99	0.09	20,22,52,55	0
4	A0Z	C	101	22/23	0.99	0.10	20,31,55,66	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around A0Z B 101:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A0Z D 101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around A0Z C 101:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.