



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 04:38 pm BST

PDB ID : 2SSP
Title : LEUCINE-272-ALANINE URACIL-DNA GLYCOSYLASE BOUND TO
ABASIC SITE-CONTAINING DNA
Authors : Parikh, S.S.; Mol, C.D.; Slupphaug, G.; Bharati, S.; Krokan, H.E.; Tainer,
J.A.
Deposited on : 1999-04-28
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	10	
2	B	11	
3	E	223	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2460 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*CP*TP*GP*TP*(AAB)P*AP*TP*CP*TP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	10	Total	C	N	O	P	0	0	0
			190	93	26	62	9			

- Molecule 2 is a DNA chain called DNA (5'-D(*AP*AP*AP*GP*AP*TP*AP*AP*CP*AP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	11	Total	C	N	O	P	0	0	0
			227	109	50	58	10			

- Molecule 3 is a protein called PROTEIN (URACIL-DNA GLYCOSYLASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	223	Total	C	N	O	S	0	0	0
			1805	1165	318	316	6			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	23	Total	O	0	0
			23	23		
4	B	16	Total	O	0	0
			16	16		
4	E	199	Total	O	0	0
			199	199		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: DNA (5'-D(*CP*TP*GP*TP*(AAB)P*AP*TP*CP*TP*T)-3')

Chain A: 





- Molecule 2: DNA (5'-D(*AP*AP*AP*GP*AP*TP*AP*AP*CP*AP*G)-3')

Chain B: 



- Molecule 3: PROTEIN (URACIL-DNA GLYCOSYLASE)

Chain E: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	120.30 Å 48.80 Å 65.60 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.25	Depositor
% Data completeness (in resolution range)	97.7 (20.00-2.25)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.184 , 0.257	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2460	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AAB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/196	0.71	0/298
2	B	0.49	0/257	0.85	0/395
3	E	0.48	0/1866	0.68	0/2529
All	All	0.48	0/2319	0.70	0/3222

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	190	0	114	15	0
2	B	227	0	124	7	0
3	E	1805	0	1756	27	0
4	A	23	0	0	5	0
4	B	16	0	0	0	0
4	E	199	0	0	3	0
All	All	2460	0	1994	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:135:LYS:HA	3:E:135:LYS:HE2	1.40	1.03
2:B:21:DA:HO5'	2:B:21:DA:H8	1.18	0.91
2:B:30:DA:H2''	2:B:31:DG:C8	2.06	0.90
2:B:29:DC:H2''	2:B:30:DA:N7	1.91	0.85
1:A:3:DG:C8	1:A:4:DT:H72	2.12	0.84
3:E:135:LYS:HE3	3:E:232:TRP:CH2	2.21	0.74
1:A:5:AAB:H2'1	3:E:268:HIS:CE1	2.30	0.66
1:A:3:DG:H3'	4:A:625:HOH:O	1.95	0.66
3:E:189:HIS:HD2	3:E:191:ASP:H	1.47	0.62
3:E:245:TRP:HE1	3:E:283:HIS:HD2	1.49	0.60
1:A:4:DT:O2	3:E:271:PRO:HB2	2.02	0.60
1:A:1:DC:C2'	1:A:2:DT:H72	2.33	0.59
3:E:135:LYS:HE3	3:E:232:TRP:HH2	1.63	0.59
2:B:29:DC:H2''	2:B:30:DA:C8	2.39	0.57
3:E:257:ASP:HB3	3:E:260:ARG:HB3	1.87	0.56
2:B:21:DA:H8	2:B:21:DA:O5'	1.86	0.56
1:A:10:DT:C5'	4:A:647:HOH:O	2.56	0.54
1:A:10:DT:H5'	4:A:647:HOH:O	2.06	0.54
1:A:3:DG:C2'	1:A:4:DT:H72	2.38	0.53
1:A:1:DC:H2''	1:A:2:DT:C7	2.39	0.52
1:A:1:DC:H2''	1:A:2:DT:H72	1.94	0.49
1:A:5:AAB:H2'1	3:E:268:HIS:NE2	2.29	0.48
2:B:30:DA:H2''	2:B:31:DG:H8	1.73	0.48
1:A:3:DG:H2''	1:A:4:DT:C7	2.44	0.48
3:E:250:GLN:HG2	3:E:265:GLN:HE21	1.80	0.46
3:E:135:LYS:CE	3:E:135:LYS:HA	2.29	0.46
3:E:142:LEU:HB3	3:E:249:ALA:HB1	1.99	0.45
4:A:620:HOH:O	3:E:212:HIS:HD2	1.98	0.45
3:E:183:ASP:O	3:E:302:LYS:HD2	2.16	0.44
1:A:3:DG:C2'	1:A:4:DT:C7	2.97	0.43
3:E:105:LEU:HD21	3:E:222:TRP:CE3	2.54	0.43
3:E:251:LYS:NZ	4:E:407:HOH:O	2.51	0.43
3:E:274:VAL:HG23	3:E:278:PHE:HB3	2.01	0.42
3:E:135:LYS:HE3	3:E:232:TRP:CZ3	2.53	0.42
3:E:131:MET:CE	3:E:193:SER:HB3	2.50	0.42
3:E:83:GLU:CG	3:E:84:PHE:H	2.32	0.42
3:E:135:LYS:CA	3:E:135:LYS:HE2	2.30	0.41
1:A:1:DC:H2'	1:A:2:DT:H72	2.02	0.41
2:B:29:DC:C2'	2:B:30:DA:N7	2.75	0.41
3:E:171:GLU:O	3:E:175:LYS:HG3	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:228:ALA:HA	4:E:473:HOH:O	2.19	0.41
3:E:288:ASN:O	3:E:292:GLN:HG2	2.21	0.41
3:E:244:LEU:HD13	3:E:250:GLN:HA	2.02	0.41
3:E:162:ARG:HG3	4:E:375:HOH:O	2.21	0.40
3:E:281:CYS:HB2	3:E:283:HIS:CE1	2.56	0.40
1:A:10:DT:H5''	4:A:647:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	E	221/223 (99%)	208 (94%)	13 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	E	194/194 (100%)	186 (96%)	8 (4%)	30	36

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	E	82	MET
3	E	135	LYS
3	E	142	LEU
3	E	168	PRO
3	E	185	VAL
3	E	254	SER
3	E	283	HIS
3	E	291	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
3	E	124	GLN
3	E	189	HIS
3	E	236	ASN
3	E	238	ASN
3	E	265	GLN
3	E	283	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	AAB	A	5	1	9,12,13	0.62	0	8,16,19	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	AAB	A	5	1	-	0/3/17/18	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	5	AAB	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.