



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 03:41 am BST

PDB ID : 3SS1  
Title : Clostridium difficile toxin A (TcdA) glucosyltransferase domain  
Authors : Pruitt, R.N.; Chumbler, N.M.; Farrow, M.A.; Seebach, S.A.; Friedman, D.B.;  
Spiller, B.W.; Lacy, D.B.  
Deposited on : 2011-07-07  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	555	<div> <div>2%</div> <div> <div></div> <div>73%</div> <div>21%</div> <div>• •</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4637 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Toxin A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	537	Total	C	N	O	S	0	0	0
			4403	2811	728	856	8			

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	543	HIS	-	EXPRESSION TAG	UNP Q189K5
A	544	ALA	-	EXPRESSION TAG	UNP Q189K5
A	545	GLY	-	EXPRESSION TAG	UNP Q189K5
A	546	LEU	-	EXPRESSION TAG	UNP Q189K5
A	547	ARG	-	EXPRESSION TAG	UNP Q189K5
A	548	GLY	-	EXPRESSION TAG	UNP Q189K5
A	549	SER	-	EXPRESSION TAG	UNP Q189K5
A	550	HIS	-	EXPRESSION TAG	UNP Q189K5
A	551	HIS	-	EXPRESSION TAG	UNP Q189K5
A	552	HIS	-	EXPRESSION TAG	UNP Q189K5
A	553	HIS	-	EXPRESSION TAG	UNP Q189K5
A	554	HIS	-	EXPRESSION TAG	UNP Q189K5
A	555	HIS	-	EXPRESSION TAG	UNP Q189K5

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mn	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	233	Total	O	0	0
			233	233		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Toxin A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	141.91Å 141.91Å 65.95Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.30 – 2.20 48.30 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.7 (48.30-2.20) 96.3 (48.30-2.20)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.95 (at 2.20Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, $R_{free}$	0.182 , 0.227 0.177 , 0.217	Depositor DCC
$R_{free}$ test set	2014 reflections (5.23%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.6	Xtriage
Anisotropy	0.181	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 45.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.038 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4637	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.38	0/4483	0.52	0/6053

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4403	0	4401	89	1
2	A	1	0	0	0	0
3	A	233	0	0	9	1
All	All	4637	0	4401	89	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:PRO:O	3:A:720:HOH:O	1.92	0.86

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:109:ILE:HD12	1:A:109:ILE:H	1.40	0.84
1:A:519:TRP:HE1	1:A:525:SER:HG	1.30	0.80
1:A:384:GLN:HE21	1:A:509:GLN:HE21	1.35	0.74
1:A:250:LEU:HD13	1:A:274:LEU:HD21	1.67	0.73
1:A:90:THR:HG23	1:A:366:ASN:HB3	1.71	0.73
1:A:384:GLN:NE2	1:A:509:GLN:HE21	1.89	0.71
1:A:361:LYS:HB2	1:A:361:LYS:NZ	2.06	0.70
1:A:302:ARG:HH11	1:A:302:ARG:HB2	1.56	0.69
1:A:55:ILE:HD11	1:A:74:LEU:HD13	1.76	0.66
1:A:362:LEU:HD22	1:A:365:LEU:HD11	1.78	0.66
1:A:91:SER:HB2	1:A:92:PRO:HD2	1.78	0.65
1:A:292:ILE:O	3:A:751:HOH:O	2.13	0.65
1:A:195:LYS:O	3:A:756:HOH:O	2.13	0.64
1:A:336:LYS:HE3	1:A:337:LEU:HA	1.82	0.62
1:A:519:TRP:NE1	1:A:525:SER:OG	2.25	0.61
1:A:5:SER:OG	1:A:8:GLU:HG3	2.01	0.61
1:A:295:ASP:HA	1:A:298:LYS:HB2	1.83	0.60
1:A:478:TYR:HA	1:A:492:LEU:HD22	1.85	0.59
1:A:115:LYS:HE2	1:A:119:ASP:OD2	2.03	0.59
1:A:478:TYR:HD2	1:A:492:LEU:HD13	1.66	0.59
1:A:487:THR:OG1	3:A:729:HOH:O	2.17	0.59
1:A:296:LEU:HD12	1:A:297:PHE:CZ	2.39	0.58
1:A:352:LYS:HG3	1:A:357:GLU:HB3	1.85	0.58
1:A:32:GLU:O	1:A:36:LEU:HD22	2.04	0.57
1:A:261:ARG:HD2	3:A:644:HOH:O	2.03	0.57
1:A:109:ILE:H	1:A:109:ILE:CD1	2.13	0.56
1:A:167:ASN:C	1:A:167:ASN:HD22	2.08	0.56
1:A:374:ILE:HG13	1:A:386:LEU:HD23	1.88	0.56
1:A:77:ASP:OD2	3:A:746:HOH:O	2.17	0.55
1:A:31:ASP:OD2	1:A:35:LYS:HE2	2.06	0.55
1:A:302:ARG:NH1	1:A:302:ARG:HB2	2.20	0.54
1:A:311:TRP:HA	1:A:314:ILE:HD11	1.89	0.54
1:A:120:ILE:HD11	1:A:355:LYS:HG2	1.90	0.54
1:A:40:ASN:OD1	1:A:43:ASN:HB2	2.08	0.54
1:A:208:HIS:HE1	3:A:742:HOH:O	1.90	0.54
1:A:38:THR:O	1:A:44:LYS:NZ	2.41	0.53
1:A:201:ILE:HD13	1:A:259:LEU:HD21	1.90	0.53
1:A:39:ASN:ND2	1:A:89:ASN:HD21	2.06	0.53
1:A:237:ILE:HD11	1:A:242:ASN:HD21	1.73	0.53
1:A:361:LYS:HB2	1:A:361:LYS:HZ2	1.73	0.53
1:A:36:LEU:HD12	1:A:38:THR:HG23	1.91	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:310:ARG:HB2	1:A:310:ARG:NH1	2.24	0.52
1:A:509:GLN:O	1:A:514:GLU:HG2	2.09	0.52
1:A:276:LEU:HG	1:A:388:SER:HB3	1.90	0.52
1:A:216:ASP:OD2	1:A:219:VAL:HG23	2.10	0.51
1:A:189:TYR:CZ	1:A:193:ILE:HG13	2.46	0.50
1:A:193:ILE:HD13	1:A:193:ILE:O	2.11	0.50
1:A:115:LYS:HE3	1:A:115:LYS:HA	1.92	0.50
1:A:40:ASN:ND2	1:A:43:ASN:H	2.11	0.48
1:A:311:TRP:HE3	1:A:314:ILE:HD11	1.77	0.48
1:A:294:SER:OG	1:A:505:ASN:OD1	2.32	0.48
1:A:97:LEU:HD23	1:A:282:VAL:HB	1.96	0.47
1:A:237:ILE:HD11	1:A:242:ASN:ND2	2.30	0.47
1:A:306:ILE:O	1:A:306:ILE:HD12	2.15	0.47
1:A:86:LYS:O	1:A:368:SER:HB3	2.15	0.46
1:A:229:ARG:HG3	1:A:230:LYS:N	2.29	0.46
1:A:532:LYS:NZ	1:A:536:ASP:OD2	2.39	0.45
1:A:413:LEU:HG	1:A:417:ILE:HG12	1.99	0.45
1:A:369:ASP:HB2	3:A:738:HOH:O	2.16	0.45
1:A:450:ALA:N	1:A:451:PRO:CD	2.79	0.45
1:A:167:ASN:C	1:A:167:ASN:ND2	2.70	0.45
1:A:293:HIS:HE1	1:A:356:SER:O	2.00	0.45
1:A:164:GLN:HE21	1:A:164:GLN:HA	1.81	0.45
1:A:12:LEU:HA	1:A:12:LEU:HD12	1.88	0.44
1:A:313:MET:HG2	1:A:510:LEU:HB3	2.00	0.43
1:A:336:LYS:HE3	1:A:337:LEU:HG	1.99	0.43
1:A:336:LYS:HA	1:A:337:LEU:HA	1.76	0.43
1:A:361:LYS:HB2	1:A:361:LYS:HZ3	1.80	0.43
1:A:183:LYS:NZ	3:A:771:HOH:O	2.51	0.43
1:A:244:LEU:HA	1:A:244:LEU:HD12	1.73	0.43
1:A:503:PRO:HB2	1:A:506:ASN:ND2	2.32	0.43
1:A:309:ASP:O	1:A:313:MET:HG3	2.19	0.42
1:A:186:ILE:HA	1:A:186:ILE:HD13	1.83	0.42
1:A:339:GLN:HA	1:A:342:LYS:HD3	2.00	0.42
1:A:374:ILE:HD13	1:A:381:VAL:HB	2.01	0.42
1:A:527:LYS:HE2	1:A:531:GLU:OE1	2.19	0.42
1:A:323:LYS:HE2	1:A:349:ILE:HG22	2.01	0.42
1:A:103:GLY:HA2	1:A:134:ALA:O	2.21	0.41
1:A:341:LEU:HG	1:A:341:LEU:H	1.61	0.41
1:A:296:LEU:HA	1:A:296:LEU:HD22	1.91	0.41
1:A:49:LYS:HA	1:A:483:LEU:HD22	2.03	0.41
1:A:374:ILE:HG13	1:A:386:LEU:CD2	2.50	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:316:LEU:O	1:A:320:MET:HG2	2.20	0.41
1:A:518:LEU:HA	1:A:518:LEU:HD23	1.93	0.41
1:A:519:TRP:HZ2	1:A:529:GLN:OE1	2.04	0.41
1:A:336:LYS:HB3	1:A:337:LEU:HB2	2.03	0.40
1:A:338:ASP:HB3	1:A:341:LEU:HD11	2.03	0.40
1:A:33:TYR:HA	1:A:36:LEU:CD2	2.51	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:777:HOH:O	3:A:781:HOH:O[4_564]	1.93	0.27
1:A:69:ARG:NH2	1:A:157:GLU:OE1[4_564]	2.13	0.07

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	535/555 (96%)	514 (96%)	20 (4%)	1 (0%)	47 55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	332	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	496/509 (97%)	459 (92%)	37 (8%)	13	14

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LEU
1	A	7	GLU
1	A	9	LEU
1	A	12	LEU
1	A	27	LEU
1	A	36	LEU
1	A	38	THR
1	A	39	ASN
1	A	40	ASN
1	A	43	ASN
1	A	74	LEU
1	A	79	LEU
1	A	86	LYS
1	A	90	THR
1	A	93	VAL
1	A	115	LYS
1	A	128	LEU
1	A	156	LEU
1	A	161	GLN
1	A	167	ASN
1	A	193	ILE
1	A	220	LEU
1	A	244	LEU
1	A	247	GLU
1	A	276	LEU
1	A	296	LEU
1	A	302	ARG
1	A	314	ILE
1	A	340	GLN
1	A	341	LEU
1	A	347	LEU
1	A	394	LEU
1	A	436	ASN
1	A	467	LEU
1	A	492	LEU
1	A	516	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	521	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	ASN
1	A	96	ASN
1	A	125	ASN
1	A	161	GLN
1	A	164	GLN
1	A	167	ASN
1	A	208	HIS
1	A	242	ASN
1	A	248	GLN
1	A	293	HIS
1	A	384	GLN
1	A	411	GLN
1	A	506	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	537/555 (96%)	-0.52	13 (2%) 59 56	18, 40, 95, 140	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	306	ILE	6.8
1	A	347	LEU	5.9
1	A	337	LEU	3.2
1	A	297	PHE	2.9
1	A	338	ASP	2.9
1	A	341	LEU	2.9
1	A	298	LYS	2.7
1	A	345	PHE	2.6
1	A	348	ILE	2.3
1	A	301	SER	2.1
1	A	350	GLU	2.0
1	A	340	GLN	2.0
1	A	349	ILE	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MN	A	1542	1/1	0.77	0.08	99,99,99,99	0

## 6.5 Other polymers

There are no such residues in this entry.