



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 03:40 am BST

PDB ID : 3SSS
Title : CcmK1 with residues 103-113 deleted
Authors : Kimber, M.S.; Samborska, B.
Deposited on : 2011-07-08
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	110	<div> <div>2%</div> <div> <div></div> <div>80%</div> <div>15%</div> <div>• 5%</div> </div> </div>
1	B	110	<div> <div>7%</div> <div> <div></div> <div>76%</div> <div>12%</div> <div>5%</div> <div>7%</div> </div> </div>
1	C	110	<div> <div>3%</div> <div> <div></div> <div>67%</div> <div>24%</div> <div>• 6%</div> </div> </div>
1	D	110	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>12%</div> <div>• 5%</div> </div> </div>
1	E	110	<div> <div>0%</div> <div> <div></div> <div>83%</div> <div>14%</div> <div>•</div> </div> </div>
1	F	110	<div> <div>7%</div> <div> <div></div> <div>82%</div> <div>11%</div> <div>• 6%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9787 atoms, of which 4845 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbon dioxide concentrating mechanism protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	105	Total	C	H	N	O	S	0	0	0
			1607	501	812	143	149	2			
1	B	102	Total	C	H	N	O	S	0	1	0
			1580	494	801	136	147	2			
1	C	103	Total	C	H	N	O	S	0	0	0
			1572	489	797	137	147	2			
1	D	104	Total	C	H	N	O	S	0	0	0
			1590	495	805	140	148	2			
1	E	106	Total	C	H	N	O	S	0	1	0
			1652	518	833	148	151	2			
1	F	103	Total	C	H	N	O	S	0	0	0
			1572	489	797	137	147	2			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	103	LEU	-	EXPRESSION TAG	UNP Q8DKB3
A	104	GLU	-	EXPRESSION TAG	UNP Q8DKB3
A	105	HIS	-	EXPRESSION TAG	UNP Q8DKB3
A	106	HIS	-	EXPRESSION TAG	UNP Q8DKB3
A	107	HIS	-	EXPRESSION TAG	UNP Q8DKB3
A	108	HIS	-	EXPRESSION TAG	UNP Q8DKB3
A	109	HIS	-	EXPRESSION TAG	UNP Q8DKB3
A	110	HIS	-	EXPRESSION TAG	UNP Q8DKB3
B	103	LEU	-	EXPRESSION TAG	UNP Q8DKB3
B	104	GLU	-	EXPRESSION TAG	UNP Q8DKB3
B	105	HIS	-	EXPRESSION TAG	UNP Q8DKB3
B	106	HIS	-	EXPRESSION TAG	UNP Q8DKB3
B	107	HIS	-	EXPRESSION TAG	UNP Q8DKB3
B	108	HIS	-	EXPRESSION TAG	UNP Q8DKB3
B	109	HIS	-	EXPRESSION TAG	UNP Q8DKB3
B	110	HIS	-	EXPRESSION TAG	UNP Q8DKB3
C	103	LEU	-	EXPRESSION TAG	UNP Q8DKB3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	104	GLU	-	EXPRESSION TAG	UNP Q8DKB3
C	105	HIS	-	EXPRESSION TAG	UNP Q8DKB3
C	106	HIS	-	EXPRESSION TAG	UNP Q8DKB3
C	107	HIS	-	EXPRESSION TAG	UNP Q8DKB3
C	108	HIS	-	EXPRESSION TAG	UNP Q8DKB3
C	109	HIS	-	EXPRESSION TAG	UNP Q8DKB3
C	110	HIS	-	EXPRESSION TAG	UNP Q8DKB3
D	103	LEU	-	EXPRESSION TAG	UNP Q8DKB3
D	104	GLU	-	EXPRESSION TAG	UNP Q8DKB3
D	105	HIS	-	EXPRESSION TAG	UNP Q8DKB3
D	106	HIS	-	EXPRESSION TAG	UNP Q8DKB3
D	107	HIS	-	EXPRESSION TAG	UNP Q8DKB3
D	108	HIS	-	EXPRESSION TAG	UNP Q8DKB3
D	109	HIS	-	EXPRESSION TAG	UNP Q8DKB3
D	110	HIS	-	EXPRESSION TAG	UNP Q8DKB3
E	103	LEU	-	EXPRESSION TAG	UNP Q8DKB3
E	104	GLU	-	EXPRESSION TAG	UNP Q8DKB3
E	105	HIS	-	EXPRESSION TAG	UNP Q8DKB3
E	106	HIS	-	EXPRESSION TAG	UNP Q8DKB3
E	107	HIS	-	EXPRESSION TAG	UNP Q8DKB3
E	108	HIS	-	EXPRESSION TAG	UNP Q8DKB3
E	109	HIS	-	EXPRESSION TAG	UNP Q8DKB3
E	110	HIS	-	EXPRESSION TAG	UNP Q8DKB3
F	103	LEU	-	EXPRESSION TAG	UNP Q8DKB3
F	104	GLU	-	EXPRESSION TAG	UNP Q8DKB3
F	105	HIS	-	EXPRESSION TAG	UNP Q8DKB3
F	106	HIS	-	EXPRESSION TAG	UNP Q8DKB3
F	107	HIS	-	EXPRESSION TAG	UNP Q8DKB3
F	108	HIS	-	EXPRESSION TAG	UNP Q8DKB3
F	109	HIS	-	EXPRESSION TAG	UNP Q8DKB3
F	110	HIS	-	EXPRESSION TAG	UNP Q8DKB3

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total Cl 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	33	Total 33	O 33	0	0
3	B	29	Total 29	O 29	0	0
3	C	36	Total 36	O 36	0	0
3	D	44	Total 44	O 44	0	0
3	E	38	Total 38	O 38	0	0
3	F	33	Total 33	O 33	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Carbon dioxide concentrating mechanism protein





- Molecule 1: Carbon dioxide concentrating mechanism protein





- Molecule 1: Carbon dioxide concentrating mechanism protein




- Molecule 1: Carbon dioxide concentrating mechanism protein



- Molecule 1: Carbon dioxide concentrating mechanism protein



● Molecule 1: Carbon dioxide concentrating mechanism protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.40 Å 78.60 Å 110.80 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.28 – 2.05 45.28 – 1.76	Depositor EDS
% Data completeness (in resolution range)	98.4 (45.28-2.05) 81.9 (45.28-1.76)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.13 (at 1.75 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.163 , 0.211 0.160 , 0.207	Depositor DCC
R_{free} test set	2827 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	25.3	Xtriage
Anisotropy	0.176	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 41.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.019 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9787	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.76	0/806	0.81	1/1094 (0.1%)
1	B	0.82	1/792 (0.1%)	0.86	2/1075 (0.2%)
1	C	0.94	1/784 (0.1%)	0.97	2/1064 (0.2%)
1	D	0.88	0/795	0.89	2/1079 (0.2%)
1	E	0.88	0/836	0.93	1/1135 (0.1%)
1	F	0.81	0/784	0.86	1/1064 (0.1%)
All	All	0.85	2/4797 (0.0%)	0.89	9/6511 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	18	GLU	CG-CD	5.70	1.60	1.51
1	B	15	ALA	CA-CB	5.39	1.63	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	23	MET	CG-SD-CE	-9.60	84.84	100.20
1	F	23	MET	CG-SD-CE	-9.40	85.16	100.20
1	D	23	MET	CG-SD-CE	-9.01	85.78	100.20
1	C	23	MET	CG-SD-CE	-8.05	87.32	100.20
1	B	23	MET	CG-SD-CE	-6.65	89.56	100.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	86	GLU	CA-CB-CG	-5.62	101.04	113.40
1	A	23	MET	CG-SD-CE	-5.58	91.27	100.20
1	C	79	ALA	C-N-CA	-5.57	107.77	121.70
1	B	41	ARG	NE-CZ-NH1	-5.24	117.68	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	2	ALA	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	795	812	809	12	0
1	B	779	801	799	15	0
1	C	775	797	795	20	0
1	D	785	805	802	9	0
1	E	819	833	827	10	0
1	F	775	797	795	8	0
2	D	1	0	0	0	0
3	A	33	0	0	1	0
3	B	29	0	0	0	0
3	C	36	0	0	0	0
3	D	44	0	0	2	0
3	E	38	0	0	2	0
3	F	33	0	0	0	0
All	All	4942	4845	4827	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:28:ARG:HE	1:A:105:HIS:HB2	1.29	0.94
1:B:93:TYR:CE2	1:B:101:ARG:HG3	2.12	0.85
1:E:86:GLU:HG3	1:E:92:ARG:HG2	1.65	0.77
1:C:63:ASN:HD22	1:C:66:ARG:HH11	1.34	0.74
1:C:2:ALA:CB	1:C:50:VAL:H	2.01	0.74
1:E:95:GLU:OE1	1:E:95:GLU:HA	1.90	0.72
1:A:2:ALA:CB	1:A:80:ARG:HG2	2.20	0.72
1:C:2:ALA:HB1	1:C:50:VAL:H	1.55	0.70
1:A:74:SER:HB2	1:F:68:ASN:O	1.92	0.70
1:C:2:ALA:HB1	1:C:50:VAL:N	2.12	0.65
1:F:2:ALA:HB2	1:F:80:ARG:HG2	1.82	0.61
1:F:2:ALA:CB	1:F:80:ARG:HG2	2.29	0.61
1:A:28:ARG:HH11	1:A:105:HIS:CG	2.19	0.60
1:A:28:ARG:HE	1:A:105:HIS:CB	2.10	0.59
1:C:3:ILE:HG21	1:C:47:ARG:HH21	1.69	0.57
1:D:98:GLU:OE2	1:D:101:ARG:HD2	2.06	0.55
1:E:63:ASN:HD22	1:E:66:ARG:NH2	2.05	0.54
1:C:3:ILE:CG2	1:C:47:ARG:HH21	2.21	0.54
1:C:38:GLY:O	1:C:39:SER:HB3	2.08	0.53
1:C:99:GLN:NE2	1:C:99:GLN:HA	2.25	0.52
1:B:103:LEU:HG	1:B:103:LEU:O	2.11	0.51
1:B:100:PHE:CZ	1:C:87:TYR:CD2	2.98	0.51
1:D:99:GLN:O	1:D:103:LEU:HD13	2.11	0.51
1:C:19:ALA:O	1:C:23:MET:HG3	2.11	0.50
1:F:5:VAL:CG1	1:F:78:ILE:HB	2.42	0.49
1:C:85:LEU:HG	1:C:89:LEU:HD12	1.93	0.49
1:D:65:LYS:HG3	3:D:156:HOH:O	2.14	0.48
1:D:100:PHE:CZ	1:E:83:GLU:HG2	2.48	0.48
1:E:66:ARG:CZ	3:E:189:HOH:O	2.62	0.48
1:B:100:PHE:HZ	1:C:87:TYR:CE2	2.31	0.47
1:F:16:VAL:HG22	1:F:44:VAL:HG13	1.97	0.47
1:B:16:VAL:HG22	1:B:44:VAL:HG13	1.96	0.46
1:C:98:GLU:OE2	1:C:98:GLU:HA	2.15	0.46
1:A:22:ALA:HB3	1:A:60:GLY:HA2	1.98	0.46
1:A:39:SER:HA	1:B:39:SER:OG	2.15	0.46
1:B:93:TYR:CZ	1:B:101:ARG:HG3	2.51	0.45
1:E:31:LEU:HA	1:E:46:VAL:HG12	1.99	0.45
1:F:2:ALA:CB	1:F:80:ARG:CG	2.94	0.44
1:C:5:VAL:HG23	1:C:81:PRO:HG3	2.00	0.44
1:D:98:GLU:HA	1:D:98:GLU:OE1	2.17	0.44
1:D:50:VAL:O	1:D:54:GLN:HG3	2.17	0.44
1:B:100:PHE:CZ	1:C:84:ASN:HA	2.53	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:66:ARG:NH2	3:E:189:HOH:O	2.52	0.43
1:B:100:PHE:HZ	1:C:87:TYR:CD2	2.35	0.43
1:A:18:GLU:OE2	1:A:63:ASN:HB3	2.19	0.43
1:B:19:ALA:O	1:B:23:MET:HG3	2.19	0.43
1:F:5:VAL:HG12	1:F:78:ILE:HB	2.01	0.43
1:D:41:ARG:HD2	3:D:145:HOH:O	2.18	0.42
1:B:100:PHE:CE1	1:C:87:TYR:CD2	3.08	0.42
1:A:63:ASN:HD22	1:A:66:ARG:CZ	2.33	0.42
1:A:70:GLY:HA2	3:A:123:HOH:O	2.18	0.42
1:C:13:PHE:HB3	1:C:14:PRO:HD3	2.02	0.41
1:E:61:VAL:CG2	1:E:75:THR:HG21	2.50	0.41
1:F:18:GLU:OE2	1:F:63:ASN:HB3	2.20	0.41
1:A:100:PHE:CE1	1:B:84:ASN:HA	2.56	0.41
1:D:19:ALA:O	1:D:23:MET:HG3	2.21	0.41
1:C:54:GLN:HA	1:C:77:ILE:CD1	2.51	0.41
1:D:100:PHE:HZ	1:E:83:GLU:HG2	1.85	0.41
1:B:62:GLU:O	1:B:65:LYS:HB2	2.21	0.40
1:B:15:ALA:HA	1:B:67:VAL:HG21	2.03	0.40
1:E:18:GLU:OE2	1:E:63:ASN:HB3	2.22	0.40
1:A:68:ASN:O	1:B:74:SER:HB2	2.21	0.40
1:C:7:MET:HG2	1:C:45:ILE:HG12	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	103/110 (94%)	101 (98%)	2 (2%)	0	100	100
1	B	101/110 (92%)	99 (98%)	2 (2%)	0	100	100
1	C	101/110 (92%)	97 (96%)	4 (4%)	0	100	100
1	D	102/110 (93%)	100 (98%)	2 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	105/110 (96%)	102 (97%)	3 (3%)	0	100	100
1	F	101/110 (92%)	99 (98%)	2 (2%)	0	100	100
All	All	613/660 (93%)	598 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	83/88 (94%)	80 (96%)	3 (4%)	35	28
1	B	82/88 (93%)	77 (94%)	5 (6%)	18	10
1	C	81/88 (92%)	77 (95%)	4 (5%)	25	17
1	D	82/88 (93%)	80 (98%)	2 (2%)	49	42
1	E	86/88 (98%)	83 (96%)	3 (4%)	36	29
1	F	81/88 (92%)	77 (95%)	4 (5%)	25	17
All	All	495/528 (94%)	474 (96%)	21 (4%)	29	22

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	VAL
1	A	39	SER
1	A	84	ASN
1	B	16	VAL
1	B	66	ARG
1	B	84	ASN
1	B	86	GLU
1	B	101	ARG
1	C	16	VAL
1	C	34	TYR
1	C	84	ASN
1	C	92	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	16	VAL
1	D	84	ASN
1	E	16	VAL
1	E	84	ASN
1	E	105	HIS
1	F	16	VAL
1	F	84	ASN
1	F	98	GLU
1	F	103	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	63	ASN
1	A	71	GLN
1	B	71	GLN
1	C	63	ASN
1	C	71	GLN
1	C	99	GLN
1	D	71	GLN
1	D	99	GLN
1	E	63	ASN
1	F	71	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	105/110 (95%)	-0.35	2 (1%) 66 71	23, 39, 75, 92	0
1	B	102/110 (92%)	-0.11	8 (7%) 13 14	24, 40, 98, 141	0
1	C	103/110 (93%)	-0.08	3 (2%) 51 56	21, 33, 72, 107	0
1	D	104/110 (94%)	-0.37	2 (1%) 66 71	21, 30, 68, 96	0
1	E	106/110 (96%)	-0.42	1 (0%) 84 86	23, 33, 63, 96	0
1	F	103/110 (93%)	-0.15	8 (7%) 13 14	26, 38, 79, 114	0
All	All	623/660 (94%)	-0.25	24 (3%) 39 42	21, 36, 78, 141	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	103	LEU	7.2
1	F	103	LEU	5.7
1	F	102	GLU	5.0
1	B	102	GLU	3.9
1	B	98	GLU	3.6
1	D	103	LEU	3.1
1	B	100	PHE	3.1
1	B	96	ALA	3.0
1	A	3	ILE	3.0
1	A	2	ALA	2.9
1	E	108	HIS	2.9
1	F	101	ARG	2.9
1	B	104	GLU	2.8
1	B	101	ARG	2.7
1	C	3	ILE	2.7
1	F	100	PHE	2.6
1	C	103	LEU	2.5
1	F	99	GLN	2.4
1	F	98	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	99	GLN	2.2
1	F	104	GLU	2.2
1	B	97	VAL	2.1
1	F	96	ALA	2.0
1	D	102	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CL	D	111	1/1	0.92	0.14	108,108,108,108	0

6.5 Other polymers [i](#)

There are no such residues in this entry.