



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 09:22 am BST

PDB ID : 3SV4  
Title : Crystal structure of the large fragment of DNA polymerase I from *Thermus Aquaticus* in an open binary complex with dT as templating nucleobase  
Authors : Betz, K.; Diederichs, K.; Marx, A.  
Deposited on : 2011-07-12  
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	540	<div> <div>9%</div> <div>82%</div> <div>15%</div> <div>..</div> </div>
2	B	12	<div> <div>33%</div> <div>67%</div> </div>
3	C	16	<div> <div>50%</div> <div>44%</div> <div>6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	FMT	A	838	-	-	X	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9546 atoms, of which 4356 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase I, thermostable.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	533	Total	C	H	N	O	S	0	11	0
			8623	2728	4333	767	783	12			

- Molecule 2 is a DNA chain called (5'-D(\*GP\*AP\*CP\*CP\*AP\*CP\*GP\*GP\*CP\*GP\*CP\*(DOC))-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	12	Total	C	H	N	O	P	0	0	0
			252	114	12	48	67	11			

- Molecule 3 is a DNA chain called (5'-D(\*AP\*AP\*AP\*TP\*GP\*GP\*CP\*GP\*CP\*CP\*GP\*TP\*GP\*GP\*TP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	15	Total	C	N	O	P	0	0	0
			310	146	58	91	15			

- Molecule 4 is FORMIC ACID (three-letter code: FMT) (formula: CH<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C H O 4 1 1 2	0	0
4	A	1	Total C H O 4 1 1 2	0	0
4	A	1	Total C H O 5 1 2 2	0	0
4	A	1	Total C H O 5 1 2 2	0	0
4	A	1	Total C H O 5 1 2 2	0	0
4	A	1	Total C H O 5 1 2 2	0	0
4	A	1	Total C O 3 1 2	0	0
4	A	1	Total C O 3 1 2	0	0
4	B	1	Total C H O 4 1 1 2	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Mg 1 1	0	0
5	A	1	Total Mg 1 1	0	0
5	C	1	Total Mg 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	260	Total 260	O 260	0	0
6	B	25	Total 25	O 25	0	0
6	C	35	Total 35	O 35	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase I, thermostable



- Molecule 2: (5'-D(\*GP\*AP\*CP\*CP\*AP\*CP\*GP\*GP\*CP\*GP\*CP\*(DOC))-3')



- Molecule 3: (5'-D(\*AP\*AP\*AP\*TP\*GP\*GP\*CP\*GP\*CP\*CP\*GP\*TP\*GP\*GP\*TP\*C)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	108.39 Å   108.39 Å   90.47 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	46.93 – 1.99 46.93 – 1.99	Depositor EDS
% Data completeness (in resolution range)	99.5 (46.93-1.99) 99.5 (46.93-1.99)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.61 (at 2.00 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, $R_{free}$	0.179 , 0.216 0.175 , 0.209	Depositor DCC
$R_{free}$ test set	2092 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.1	Xtriage
Anisotropy	0.141	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 47.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.037 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9546	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DOC, FMT, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.22	0/4408	0.41	0/5968
2	B	0.48	0/249	1.03	1/382 (0.3%)
3	C	0.46	0/347	0.97	0/534
All	All	0.26	0/5004	0.52	1/6884 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	B	108	DG	O4'-C1'-N9	5.66	111.96	108.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4290	4333	4339	61	1
2	B	240	12	134	9	0
3	C	310	0	169	9	0
4	A	24	10	8	4	0
4	B	3	1	1	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	1	0	0	0	0
6	A	260	0	0	6	0
6	B	25	0	0	0	0
6	C	35	0	0	2	0
All	All	5190	4356	4651	77	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:732:ASP:O	6:A:849:HOH:O	1.87	0.92
4:A:838:FMT:H	3:C:211:DG:H21	1.55	0.70
1:A:559:ARG:HH11	1:A:601[B]:GLU:CG	2.08	0.66
1:A:627[B]:ASN:OD1	1:A:630:ARG:NH1	2.31	0.63
1:A:425:ARG:NH2	1:A:723:LEU:O	2.34	0.60
1:A:694:GLU:O	1:A:698:GLN:HG2	2.02	0.60
1:A:741:ARG:HA	6:A:849:HOH:O	2.03	0.59
1:A:685:PRO:HB2	1:A:687:GLU:HG3	1.83	0.59
1:A:559:ARG:HH11	1:A:601[B]:GLU:HG3	1.68	0.58
1:A:689:ALA:HA	1:A:692:PHE:CZ	2.38	0.58
1:A:646:MET:SD	1:A:662:ALA:HA	2.44	0.57
1:A:679:SER:HB2	1:A:686:TYR:HA	1.86	0.57
1:A:732:ASP:HB3	6:A:849:HOH:O	2.04	0.57
1:A:303[B]:GLU:HG2	1:A:346:LYS:HE2	1.87	0.57
3:C:214:DG:H2''	3:C:215:DT:C5'	2.35	0.56
2:B:102:DA:H2'	2:B:103:DC:C6	2.40	0.56
3:C:210:DC:H2'	3:C:211:DG:C8	2.40	0.56
1:A:639:HIS:CD2	1:A:663:LYS:HD3	2.41	0.56
1:A:682:LEU:O	1:A:684:ILE:N	2.38	0.56
1:A:675:ALA:HB2	1:A:690:GLN:HB2	1.87	0.55
1:A:679:SER:CB	1:A:686:TYR:HA	2.36	0.55
1:A:559:ARG:NH1	1:A:601[B]:GLU:HG2	2.22	0.55
2:B:103:DC:H2''	2:B:104:DC:H5'	1.90	0.54
1:A:576:SER:O	3:C:208:DG:H4'	2.08	0.54
1:A:831:LYS:O	1:A:832:GLU:C	2.46	0.54
1:A:299:TRP:CG	1:A:300:PRO:HA	2.43	0.53
1:A:740:VAL:HG12	6:A:849:HOH:O	2.08	0.53
1:A:376:LEU:HD22	1:A:420:LEU:HD22	1.90	0.53
1:A:559:ARG:NH1	1:A:601[B]:GLU:CG	2.72	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:302:PRO:HG2	1:A:328:ARG:HD3	1.92	0.52
1:A:675:ALA:CB	1:A:690:GLN:HB2	2.39	0.52
1:A:646:MET:HE1	1:A:665:ILE:HG21	1.93	0.51
3:C:214:DG:H2''	3:C:215:DT:O5'	2.11	0.51
1:A:682:LEU:HD22	1:A:682:LEU:N	2.26	0.50
1:A:775:MET:SD	1:A:802[B]:LEU:HD23	2.52	0.50
4:A:838:FMT:H	6:C:225:HOH:O	2.10	0.50
2:B:101:DG:H2'	2:B:102:DA:C8	2.47	0.50
4:A:838:FMT:C	3:C:211:DG:H21	2.24	0.49
1:A:735:ALA:HB3	6:A:849:HOH:O	2.11	0.49
1:A:526:HIS:ND1	1:A:528:ILE:HG22	2.28	0.49
1:A:616:LEU:CD2	1:A:667:PHE:CZ	2.96	0.49
1:A:664:THR:OG1	3:C:202:DA:H4'	2.13	0.49
1:A:690:GLN:HA	1:A:693:ILE:HD12	1.95	0.48
1:A:516:ALA:O	1:A:520:GLU:HG3	2.14	0.48
2:B:111:DC:H2'	2:B:112:DOC:C6	2.45	0.47
1:A:547:ASP:HB2	1:A:548:PRO:HD3	1.97	0.46
3:C:214:DG:H2''	3:C:215:DT:H5'	1.97	0.46
1:A:616:LEU:HD21	1:A:667:PHE:CZ	2.50	0.46
1:A:645:TRP:O	1:A:645:TRP:CD1	2.69	0.46
1:A:587[A]:ARG:NH1	2:B:112:DOC:OP2	2.49	0.45
1:A:492:ARG:O	1:A:496:ASP:HB2	2.16	0.45
2:B:104:DC:H2''	2:B:105:DA:H5'	1.98	0.45
1:A:519:LEU:HB2	1:A:533:LEU:HD21	1.99	0.44
1:A:303[B]:GLU:HG2	1:A:346:LYS:CE	2.46	0.44
1:A:642:THR:HG22	1:A:646:MET:CE	2.48	0.44
1:A:646:MET:CE	1:A:665:ILE:HG21	2.47	0.44
1:A:689:ALA:HA	1:A:692:PHE:CE1	2.53	0.44
1:A:780:LEU:HD11	1:A:790:GLU:HB2	2.00	0.44
2:B:111:DC:H2'	2:B:112:DOC:H6	2.00	0.44
1:A:559:ARG:HH11	1:A:601[B]:GLU:HG2	1.78	0.43
1:A:380:LEU:HD12	1:A:416:LEU:HD12	2.01	0.43
1:A:595:ARG:HD3	6:A:848:HOH:O	2.19	0.43
1:A:490:LEU:HD11	1:A:532:ILE:HD13	1.99	0.43
1:A:767:LYS:HG2	4:A:837:FMT:C	2.49	0.43
2:B:103:DC:H2''	2:B:104:DC:C5'	2.49	0.43
1:A:306:PHE:O	1:A:326:ALA:HA	2.19	0.42
1:A:631:VAL:O	1:A:636:ARG:HG2	2.20	0.42
1:A:667:PHE:O	1:A:670:LEU:HB3	2.20	0.42
2:B:104:DC:H2''	2:B:105:DA:C5'	2.50	0.41
1:A:724:PHE:CD2	1:A:759:ASP:HB3	2.54	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:204:DT:O2	6:C:5:HOH:O	2.21	0.41
1:A:541:LEU:HD12	1:A:590:LEU:HD23	2.02	0.41
1:A:588:THR:HB	1:A:589:PRO:HD2	2.02	0.41
1:A:751:MET:HB3	1:A:752:PRO:HD3	2.01	0.41
1:A:655:ASP:HB2	1:A:658:MET:CB	2.50	0.41
1:A:698:GLN:HA	1:A:698:GLN:OE1	2.21	0.40
1:A:665:ILE:HG23	1:A:666:ASN:N	2.37	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:392:ARG:NH2	1:A:602:GLU:OE2[2_545]	2.03	0.17

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	540/540 (100%)	517 (96%)	21 (4%)	2 (0%)	34 30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	586	VAL
1	A	683	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	443/441 (100%)	432 (98%)	11 (2%)	47 49

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	421	GLU
1	A	487	ARG
1	A	616	LEU
1	A	642	THR
1	A	686	TYR
1	A	687	GLU
1	A	692	PHE
1	A	713	GLU
1	A	749[A]	PHE
1	A	749[B]	PHE
1	A	832	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	DOC	B	112	3,2	14,19,20	2.57	6 (42%)	13,26,29	1.83	3 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DOC	B	112	3,2	-	0/4/18/19	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	112	DOC	C2-N3	5.36	1.48	1.38
2	B	112	DOC	C5-C4	4.81	1.52	1.41
2	B	112	DOC	C6-N1	3.52	1.40	1.35
2	B	112	DOC	C4-N4	3.14	1.44	1.35
2	B	112	DOC	O4'-C1'	-2.38	1.37	1.42
2	B	112	DOC	C4-N3	-2.04	1.32	1.35

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	112	DOC	C2-N3-C4	3.63	120.02	116.34
2	B	112	DOC	N4-C4-N3	2.86	121.01	116.49
2	B	112	DOC	O4'-C1'-C2'	2.52	109.40	106.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	112	DOC	3	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 3 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	FMT	A	835	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	836	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	834	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	833	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	838	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	837	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	1	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	A	839	-	0,2,2	0.00	-	0,1,1	0.00	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	838	FMT	3	0
4	A	837	FMT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	533/540 (98%)	0.53	51 (9%) <b>8</b> <b>7</b>	22, 35, 105, 146	0
2	B	11/12 (91%)	-0.35	0 <b>100</b> <b>100</b>	28, 34, 57, 60	0
3	C	15/16 (93%)	-0.23	0 <b>100</b> <b>100</b>	26, 36, 60, 72	0
All	All	559/568 (98%)	0.49	51 (9%) <b>9</b> <b>8</b>	22, 35, 103, 146	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	688	GLU	9.3
1	A	692	PHE	9.0
1	A	686	TYR	7.9
1	A	665	ILE	7.8
1	A	654	VAL	7.8
1	A	676	HIS	7.5
1	A	646	MET	6.7
1	A	675	ALA	6.6
1	A	644	SER	6.5
1	A	669	VAL	6.2
1	A	645	TRP	6.1
1	A	680	GLN	6.0
1	A	687	GLU	5.8
1	A	657	LEU	5.7
1	A	655	ASP	5.6
1	A	684	ILE	5.6
1	A	685	PRO	5.5
1	A	658	MET	5.4
1	A	643	ALA	5.2
1	A	678	LEU	5.2
1	A	293	ALA	5.1
1	A	683	ALA	5.0
1	A	689	ALA	4.9

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	690	GLN	4.8
1	A	642	THR	4.4
1	A	636	ARG	4.4
1	A	681	GLU	4.2
1	A	693	ILE	4.2
1	A	674	SER	4.1
1	A	664	THR	3.7
1	A	682	LEU	3.6
1	A	640	THR	3.6
1	A	736	ARG	3.4
1	A	661	ALA	3.3
1	A	698	GLN	3.3
1	A	670	LEU	3.2
1	A	695	ARG	3.2
1	A	660	ARG	3.1
1	A	641	GLU	3.1
1	A	697	PHE	2.9
1	A	696	TYR	2.8
1	A	668	GLY	2.7
1	A	656	PRO	2.7
1	A	679	SER	2.6
1	A	694	GLU	2.6
1	A	691	ALA	2.6
1	A	673	MET	2.6
1	A	677	ARG	2.5
1	A	666	ASN	2.5
1	A	749[A]	PHE	2.4
1	A	832	GLU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	DOC	B	112	18/19	0.97	0.12	24,32,38,39	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	FMT	A	1	3/3	0.72	0.20	67,69,71,86	0
5	MG	A	7	1/1	0.74	0.15	77,77,77,77	0
4	FMT	A	839	3/3	0.81	0.15	41,41,44,47	0
4	FMT	A	835	3/3	0.82	0.22	48,58,67,77	0
4	FMT	A	837	3/3	0.90	0.14	46,53,64,69	0
4	FMT	A	838	3/3	0.91	0.27	38,38,40,47	0
4	FMT	B	1	3/3	0.91	0.12	40,47,55,56	0
4	FMT	A	833	3/3	0.93	0.35	66,67,68,81	0
4	FMT	A	834	3/3	0.94	0.24	47,59,71,71	0
5	MG	C	7	1/1	0.94	0.06	60,60,60,60	0
4	FMT	A	836	3/3	0.96	0.10	44,53,58,66	0
5	MG	B	7	1/1	0.97	0.05	44,44,44,44	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.