



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 05:47 pm BST

PDB ID : 5SZD  
Title : Crystal structure of Aquifex aeolicus Hfq at 1.5Å  
Authors : Stanek, K.; Patterson, J.; Randolph, P.S.; Mura, C.  
Deposited on : 2016-08-13  
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	83	<div><div>4%</div><div><div></div><div>80%</div><div>7%</div><div>13%</div></div></div>
1	B	83	<div><div>4%</div><div><div></div><div>80%</div><div>5%</div><div>16%</div></div></div>
1	C	83	<div><div>5%</div><div><div></div><div>82%</div><div>•</div><div>14%</div></div></div>
1	D	83	<div><div>2%</div><div><div></div><div>83%</div><div></div><div>17%</div></div></div>
1	E	83	<div><div>2%</div><div><div></div><div>83%</div><div></div><div>17%</div></div></div>
1	F	83	<div><div>6%</div><div><div></div><div>86%</div><div>•</div><div>13%</div></div></div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	83	
1	H	83	
1	I	83	
1	J	83	
1	K	83	
1	L	83	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GAI	A	104	-	X	-	-
4	GAI	A	105	-	X	-	-
4	GAI	B	103	-	X	-	-
4	GAI	C	103	-	X	-	-
4	GAI	E	103	-	X	-	-
4	GAI	F	103	-	X	-	-
4	GAI	G	104	-	X	-	-
4	GAI	H	105	-	X	-	-

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8350 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-binding protein Hfq.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	72	Total	C	N	O	S	0	8	0
			650	416	118	114	2			
1	B	70	Total	C	N	O	S	0	6	0
			622	401	113	107	1			
1	C	71	Total	C	N	O	S	0	8	0
			654	424	118	110	2			
1	D	69	Total	C	N	O	S	0	4	0
			592	383	107	101	1			
1	E	69	Total	C	N	O	S	0	5	0
			615	397	111	106	1			
1	F	72	Total	C	N	O	S	0	6	0
			640	409	117	112	2			
1	G	70	Total	C	N	O	S	0	11	0
			659	421	122	115	1			
1	H	71	Total	C	N	O	S	0	7	0
			629	404	115	108	2			
1	I	71	Total	C	N	O	S	0	10	0
			660	423	121	114	2			
1	J	71	Total	C	N	O	S	0	10	0
			669	427	127	114	1			
1	K	71	Total	C	N	O	S	0	7	0
			636	408	118	109	1			
1	L	71	Total	C	N	O	S	0	8	0
			644	414	119	110	1			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP O66512
A	2	SER	-	expression tag	UNP O66512
A	3	HIS	-	expression tag	UNP O66512
B	1	GLY	-	expression tag	UNP O66512
B	2	SER	-	expression tag	UNP O66512

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	3	HIS	-	expression tag	UNP O66512
C	1	GLY	-	expression tag	UNP O66512
C	2	SER	-	expression tag	UNP O66512
C	3	HIS	-	expression tag	UNP O66512
D	1	GLY	-	expression tag	UNP O66512
D	2	SER	-	expression tag	UNP O66512
D	3	HIS	-	expression tag	UNP O66512
E	1	GLY	-	expression tag	UNP O66512
E	2	SER	-	expression tag	UNP O66512
E	3	HIS	-	expression tag	UNP O66512
F	1	GLY	-	expression tag	UNP O66512
F	2	SER	-	expression tag	UNP O66512
F	3	HIS	-	expression tag	UNP O66512
G	1	GLY	-	expression tag	UNP O66512
G	2	SER	-	expression tag	UNP O66512
G	3	HIS	-	expression tag	UNP O66512
H	1	GLY	-	expression tag	UNP O66512
H	2	SER	-	expression tag	UNP O66512
H	3	HIS	-	expression tag	UNP O66512
I	1	GLY	-	expression tag	UNP O66512
I	2	SER	-	expression tag	UNP O66512
I	3	HIS	-	expression tag	UNP O66512
J	1	GLY	-	expression tag	UNP O66512
J	2	SER	-	expression tag	UNP O66512
J	3	HIS	-	expression tag	UNP O66512
K	1	GLY	-	expression tag	UNP O66512
K	2	SER	-	expression tag	UNP O66512
K	3	HIS	-	expression tag	UNP O66512
L	1	GLY	-	expression tag	UNP O66512
L	2	SER	-	expression tag	UNP O66512
L	3	HIS	-	expression tag	UNP O66512


- Molecule 2 is (4R)-2-METHYLPENTANE-2,4-DIOL (three-letter code: MRD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			8	6	2		
2	A	1	Total	C	O	0	0
			8	6	2		
2	B	1	Total	C	O	0	0
			8	6	2		
2	C	1	Total	C	O	0	0
			8	6	2		
2	D	1	Total	C	O	0	0
			8	6	2		
2	E	1	Total	C	O	0	0
			8	6	2		
2	F	1	Total	C	O	0	0
			8	6	2		
2	G	1	Total	C	O	0	0
			8	6	2		
2	H	1	Total	C	O	0	0
			8	6	2		
2	I	1	Total	C	O	0	0
			8	6	2		
2	J	1	Total	C	O	0	0
			8	6	2		
2	K	1	Total	C	O	0	0
			8	6	2		
2	L	1	Total	C	O	0	0
			8	6	2		


- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula:

C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	6	2		
3	B	1	Total	C	O	0	0
			8	6	2		
3	C	1	Total	C	O	0	0
			8	6	2		
3	D	1	Total	C	O	0	0
			8	6	2		
3	E	1	Total	C	O	0	0
			8	6	2		
3	G	1	Total	C	O	0	0
			8	6	2		
3	H	1	Total	C	O	0	0
			8	6	2		
3	H	1	Total	C	O	0	0
			8	6	2		
3	I	1	Total	C	O	0	0
			8	6	2		
3	J	1	Total	C	O	0	0
			8	6	2		
3	K	1	Total	C	O	0	0
			8	6	2		
3	L	1	Total	C	O	0	0
			8	6	2		

- Molecule 4 is GUANIDINE (three-letter code: GAI) (formula: CH<sub>5</sub>N<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			4	1	3		
4	A	1	Total	C	N	0	0
			4	1	3		
4	B	1	Total	C	N	0	0
			4	1	3		
4	C	1	Total	C	N	0	0
			4	1	3		
4	E	1	Total	C	N	0	0
			4	1	3		
4	F	1	Total	C	N	0	0
			4	1	3		
4	G	1	Total	C	N	0	0
			4	1	3		
4	H	1	Total	C	N	0	0
			4	1	3		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	G	1	Total	Cl	0	0
			1	1		
5	J	1	Total	Cl	0	0
			1	1		
5	K	1	Total	Cl	0	0
			1	1		
5	H	1	Total	Cl	0	0
			1	1		


*Continued on next page...*



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	I	1	Total Cl 1 1	0	0
5	L	1	Total Cl 1 1	0	0
5	F	1	Total Cl 1 1	0	0

- Molecule 6 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	G	1	Total C O 7 4 3	0	0
6	J	1	Total C O 7 4 3	0	0
6	K	1	Total C O 7 4 3	0	0
6	L	1	Total C O 7 4 3	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	36	Total O 36 36	0	0
7	B	37	Total O 37 37	0	0

Continued on next page...



*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	38	Total 38	O 38	0	0
7	D	32	Total 32	O 32	0	0
7	E	35	Total 35	O 35	0	0
7	F	32	Total 32	O 32	0	0
7	G	29	Total 29	O 29	0	0
7	H	38	Total 38	O 38	0	0
7	I	35	Total 35	O 35	0	0
7	J	31	Total 31	O 31	0	0
7	K	33	Total 33	O 33	0	0
7	L	37	Total 37	O 37	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: RNA-binding protein Hfq





- Molecule 1: RNA-binding protein Hfq





- Molecule 1: RNA-binding protein Hfq





- Molecule 1: RNA-binding protein Hfq




- Molecule 1: RNA-binding protein Hfq





- Molecule 1: RNA-binding protein Hfq




- Molecule 1: RNA-binding protein Hfq





- Molecule 1: RNA-binding protein Hfq





- Molecule 1: RNA-binding protein Hfq





- Molecule 1: RNA-binding protein Hfq



- Molecule 1: RNA-binding protein Hfq



- Molecule 1: RNA-binding protein Hfq



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.46Å 66.06Å 66.10Å 60.05° 83.94° 77.17°	Depositor
Resolution (Å)	46.35 – 1.49 46.35 – 1.49	Depositor EDS
% Data completeness (in resolution range)	93.9 (46.35-1.49) 90.9 (46.35-1.49)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.23 (at 1.50Å)	Xtriage
Refinement program	PHENIX (1.10-2155_1309: ???)	Depositor
R, $R_{free}$	0.132 , 0.170 0.133 , 0.170	Depositor DCC
$R_{free}$ test set	10627 reflections (7.69%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	11.9	Xtriage
Anisotropy	0.559	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 60.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.046 for -h,-k,-k+l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	8350	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MRD, MPD, PEG, GAI, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.30	0/673	0.52	0/904
1	B	0.30	0/640	0.50	0/859
1	C	0.29	0/670	0.51	0/899
1	D	0.30	0/613	0.52	0/822
1	E	0.30	0/624	0.52	0/837
1	F	0.30	0/654	0.53	0/877
1	G	0.30	0/682	0.55	0/914
1	H	0.32	0/652	0.55	0/877
1	I	0.31	0/679	0.52	0/911
1	J	0.30	0/684	0.53	0/915
1	K	0.29	0/655	0.54	0/878
1	L	0.31	0/663	0.53	0/889
All	All	0.30	0/7889	0.53	0/10582

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	650	0	681	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	622	0	658	3	0
1	C	654	0	695	3	0
1	D	592	0	634	0	0
1	E	615	0	654	0	0
1	F	640	0	676	1	0
1	G	659	0	685	2	0
1	H	629	0	660	3	0
1	I	660	0	693	3	0
1	J	669	0	704	4	0
1	K	636	0	669	4	0
1	L	644	0	679	3	0
2	A	16	0	28	0	0
2	B	8	0	14	0	0
2	C	8	0	14	0	0
2	D	8	0	14	0	0
2	E	8	0	14	0	0
2	F	8	0	14	0	0
2	G	8	0	14	0	0
2	H	8	0	14	0	0
2	I	8	0	14	0	0
2	J	8	0	14	0	0
2	K	8	0	14	0	0
2	L	8	0	14	0	0
3	A	8	0	14	0	0
3	B	8	0	14	0	0
3	C	8	0	14	0	0
3	D	8	0	14	0	0
3	E	8	0	14	0	0
3	G	8	0	14	0	0
3	H	16	0	28	0	0
3	I	8	0	14	0	0
3	J	8	0	14	0	0
3	K	8	0	14	0	0
3	L	8	0	14	0	0
4	A	8	0	10	1	0
4	B	4	0	5	1	0
4	C	4	0	5	0	0
4	E	4	0	4	0	0
4	F	4	0	5	0	0
4	G	4	0	4	0	0
4	H	4	0	5	0	0
5	F	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	G	1	0	0	0	0
5	H	1	0	0	0	0
5	I	1	0	0	0	0
5	J	1	0	0	0	0
5	K	1	0	0	0	0
5	L	1	0	0	0	0
6	G	7	0	10	1	0
6	J	7	0	10	0	0
6	K	7	0	10	0	0
6	L	7	0	10	1	0
7	A	36	0	0	1	0
7	B	37	0	0	1	0
7	C	38	0	0	1	0
7	D	32	0	0	0	0
7	E	35	0	0	0	0
7	F	32	0	0	0	0
7	G	29	0	0	0	0
7	H	38	0	0	0	0
7	I	35	0	0	0	0
7	J	31	0	0	0	0
7	K	33	0	0	0	0
7	L	37	0	0	0	0
All	All	8350	0	8516	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:39:SER:OG	1:K:7[A]:LYS:NZ	2.26	0.68
1:C:6[B]:TYR:O	7:C:201:HOH:O	2.14	0.65
1:G:18[B]:LYS:HE3	1:G:19[B]:LYS:HE2	1.82	0.62
1:G:7[B]:LYS:NZ	1:L:40:PHE:O	2.35	0.60
1:H:21[A]:VAL:HG11	1:H:70:LEU:HD21	1.83	0.60
1:K:18[B]:LYS:HE2	1:K:19[B]:LYS:HE2	1.86	0.58
1:B:40:PHE:O	1:C:5[A]:PRO:HA	2.05	0.57
1:I:32:ARG:NH2	1:I:34:GLN:OE1	2.41	0.54
1:K:40:PHE:O	1:L:7[A]:LYS:NZ	2.37	0.53
1:B:30:GLY:HA3	4:B:103:GAI:HN32	1.74	0.52
6:G:105:PEG:H42	1:L:31:VAL:HG21	1.92	0.51

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:4[A]:MET:HB2	1:A:7:LYS:HB2	1.96	0.47
1:J:18[B]:LYS:HE3	1:J:19[B]:LYS:HE2	1.97	0.47
1:I:40:PHE:O	1:J:7[B]:LYS:NZ	2.50	0.44
1:F:22[A]:LYS:HB2	1:F:22[A]:LYS:HE3	1.78	0.44
1:H:2[B]:SER:OG	1:H:7[B]:LYS:HD2	2.18	0.44
1:H:2[A]:SER:HB2	1:H:7[A]:LYS:HD3	1.99	0.44
1:I:48[B]:GLU:OE1	1:J:69:ARG:NH1	2.38	0.43
1:B:4:MET:HB3	1:B:7:LYS:HB2	2.01	0.42
1:A:11[C]:SER:OG	1:A:72:ILE:HG23	2.19	0.42
7:B:228:HOH:O	1:C:7[B]:LYS:HA	2.20	0.41
1:K:31:VAL:HG21	6:L:104:PEG:H42	2.01	0.41
1:A:22[B]:LYS:HE3	7:A:219:HOH:O	2.21	0.41
1:A:30:GLY:HA3	4:A:104:GAI:HN22	1.86	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	79/83 (95%)	78 (99%)	1 (1%)	0	100	100
1	B	74/83 (89%)	73 (99%)	1 (1%)	0	100	100
1	C	77/83 (93%)	76 (99%)	1 (1%)	0	100	100
1	D	71/83 (86%)	69 (97%)	2 (3%)	0	100	100
1	E	72/83 (87%)	71 (99%)	1 (1%)	0	100	100
1	F	77/83 (93%)	76 (99%)	1 (1%)	0	100	100
1	G	78/83 (94%)	77 (99%)	1 (1%)	0	100	100
1	H	75/83 (90%)	72 (96%)	3 (4%)	0	100	100
1	I	78/83 (94%)	77 (99%)	1 (1%)	0	100	100
1	J	78/83 (94%)	77 (99%)	1 (1%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	K	75/83 (90%)	73 (97%)	2 (3%)	0	100	100
1	L	76/83 (92%)	75 (99%)	1 (1%)	0	100	100
All	All	910/996 (91%)	894 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	75/73 (103%)	75 (100%)	0	100	100
1	B	71/73 (97%)	71 (100%)	0	100	100
1	C	74/73 (101%)	74 (100%)	0	100	100
1	D	68/73 (93%)	68 (100%)	0	100	100
1	E	69/73 (94%)	69 (100%)	0	100	100
1	F	73/73 (100%)	73 (100%)	0	100	100
1	G	76/73 (104%)	76 (100%)	0	100	100
1	H	73/73 (100%)	71 (97%)	2 (3%)	44	15
1	I	76/73 (104%)	76 (100%)	0	100	100
1	J	76/73 (104%)	76 (100%)	0	100	100
1	K	73/73 (100%)	71 (97%)	2 (3%)	44	15
1	L	74/73 (101%)	72 (97%)	2 (3%)	44	15
All	All	878/876 (100%)	872 (99%)	6 (1%)	91	69

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	7[A]	LYS
1	H	7[B]	LYS
1	K	7[A]	LYS
1	K	7[B]	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	L	7[A]	LYS
1	L	7[B]	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 44 ligands modelled in this entry, 7 are monoatomic - leaving 37 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	PEG	K	104	-	6,6,6	0.47	0	5,5,5	0.32	0
2	MRD	E	101	-	7,7,7	0.24	0	9,10,10	0.24	0
3	MPD	H	102	-	7,7,7	0.27	0	9,10,10	0.25	0
3	MPD	J	102	-	7,7,7	0.25	0	9,10,10	0.23	0
2	MRD	A	102	-	7,7,7	0.30	0	9,10,10	0.17	0
3	MPD	L	102	-	7,7,7	0.27	0	9,10,10	0.26	0
3	MPD	I	102	-	7,7,7	0.26	0	9,10,10	0.30	0
3	MPD	G	102	-	7,7,7	0.27	0	9,10,10	0.23	0
2	MRD	H	101	-	7,7,7	0.26	0	9,10,10	0.26	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	MPD	B	102	-	7,7,7	0.29	0	9,10,10	0.40	0
2	MRD	G	101	-	7,7,7	0.30	0	9,10,10	0.34	0
3	MPD	K	102	-	7,7,7	0.29	0	9,10,10	0.28	0
3	MPD	E	102	-	7,7,7	0.31	0	9,10,10	0.28	0
4	GAI	B	103	-	3,3,3	6.16	3 (100%)	3,3,3	1.01	0
2	MRD	I	101	-	7,7,7	0.29	0	9,10,10	0.41	0
4	GAI	C	103	-	3,3,3	6.05	3 (100%)	3,3,3	0.97	0
3	MPD	C	102	-	7,7,7	0.27	0	9,10,10	0.25	0
4	GAI	G	104	-	3,3,3	4.46	3 (100%)	3,3,3	1.18	0
2	MRD	B	101	-	7,7,7	0.24	0	9,10,10	0.27	0
2	MRD	D	101	-	7,7,7	0.28	0	9,10,10	0.33	0
4	GAI	F	103	-	3,3,3	6.10	3 (100%)	3,3,3	1.10	0
3	MPD	A	103	-	7,7,7	0.26	0	9,10,10	0.27	0
6	PEG	J	104	-	6,6,6	0.51	0	5,5,5	0.34	0
4	GAI	E	103	-	3,3,3	4.54	3 (100%)	3,3,3	1.10	0
2	MRD	K	101	-	7,7,7	0.24	0	9,10,10	0.34	0
2	MRD	J	101	-	7,7,7	0.28	0	9,10,10	0.28	0
2	MRD	A	101	-	7,7,7	0.25	0	9,10,10	0.34	0
2	MRD	C	101	-	7,7,7	0.27	0	9,10,10	0.23	0
3	MPD	D	102	-	7,7,7	0.30	0	9,10,10	0.28	0
3	MPD	H	103	-	7,7,7	0.29	0	9,10,10	0.27	0
6	PEG	G	105	-	6,6,6	0.49	0	5,5,5	0.19	0
6	PEG	L	104	-	6,6,6	0.48	0	5,5,5	0.23	0
4	GAI	A	105	-	3,3,3	6.14	3 (100%)	3,3,3	1.05	0
4	GAI	A	104	-	3,3,3	6.08	3 (100%)	3,3,3	1.02	0
2	MRD	F	101	-	7,7,7	0.33	0	9,10,10	0.21	0
2	MRD	L	101	-	7,7,7	0.32	0	9,10,10	0.30	0
4	GAI	H	105	-	3,3,3	6.10	3 (100%)	3,3,3	1.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	K	104	-	-	1/4/4/4	-
2	MRD	E	101	-	-	0/5/5/5	-
3	MPD	H	102	-	-	0/5/5/5	-
3	MPD	J	102	-	-	0/5/5/5	-
2	MRD	A	102	-	-	0/5/5/5	-
3	MPD	L	102	-	-	0/5/5/5	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	I	102	-	-	0/5/5/5	-
3	MPD	G	102	-	-	0/5/5/5	-
2	MRD	H	101	-	-	0/5/5/5	-
3	MPD	B	102	-	-	0/5/5/5	-
2	MRD	G	101	-	-	0/5/5/5	-
3	MPD	K	102	-	-	0/5/5/5	-
3	MPD	E	102	-	-	0/5/5/5	-
2	MRD	I	101	-	-	0/5/5/5	-
3	MPD	C	102	-	-	1/5/5/5	-
2	MRD	B	101	-	-	0/5/5/5	-
2	MRD	D	101	-	-	0/5/5/5	-
3	MPD	A	103	-	-	1/5/5/5	-
6	PEG	J	104	-	-	1/4/4/4	-
2	MRD	K	101	-	-	0/5/5/5	-
2	MRD	J	101	-	-	0/5/5/5	-
2	MRD	A	101	-	-	0/5/5/5	-
2	MRD	C	101	-	-	0/5/5/5	-
3	MPD	D	102	-	-	1/5/5/5	-
3	MPD	H	103	-	-	0/5/5/5	-
6	PEG	G	105	-	-	4/4/4/4	-
6	PEG	L	104	-	-	2/4/4/4	-
2	MRD	F	101	-	-	0/5/5/5	-
2	MRD	L	101	-	-	0/5/5/5	-

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	103	GAI	C-N1	7.17	1.45	1.30
4	C	103	GAI	C-N1	7.16	1.45	1.30
4	A	104	GAI	C-N1	7.12	1.45	1.30
4	H	105	GAI	C-N1	7.11	1.45	1.30
4	A	105	GAI	C-N1	7.10	1.45	1.30
4	F	103	GAI	C-N1	7.02	1.45	1.30
4	B	103	GAI	C-N3	-5.89	1.25	1.36
4	A	105	GAI	C-N2	-5.85	1.25	1.36
4	F	103	GAI	C-N3	-5.82	1.25	1.36
4	H	105	GAI	C-N3	-5.80	1.25	1.36
4	A	104	GAI	C-N2	-5.76	1.25	1.36
4	C	103	GAI	C-N2	-5.73	1.25	1.36
4	E	103	GAI	C-N2	5.41	1.45	1.36
4	F	103	GAI	C-N2	5.34	1.45	1.36
4	A	105	GAI	C-N3	5.34	1.45	1.36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	103	GAI	C-N2	5.26	1.45	1.36
4	G	104	GAI	C-N3	5.26	1.45	1.36
4	H	105	GAI	C-N2	5.23	1.45	1.36
4	E	103	GAI	C-N3	5.22	1.45	1.36
4	A	104	GAI	C-N3	5.21	1.45	1.36
4	G	104	GAI	C-N2	5.11	1.45	1.36
4	C	103	GAI	C-N3	5.06	1.45	1.36
4	G	104	GAI	C-N1	-2.43	1.25	1.30
4	E	103	GAI	C-N1	-2.33	1.25	1.30

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	G	105	PEG	O2-C3-C4-O4
6	K	104	PEG	O1-C1-C2-O2
6	L	104	PEG	O2-C3-C4-O4
6	G	105	PEG	O1-C1-C2-O2
3	A	103	MPD	O2-C2-C3-C4
6	G	105	PEG	C1-C2-O2-C3
6	G	105	PEG	C4-C3-O2-C2
6	L	104	PEG	C1-C2-O2-C3
3	D	102	MPD	CM-C2-C3-C4
6	J	104	PEG	C4-C3-O2-C2
3	C	102	MPD	O2-C2-C3-C4

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	103	GAI	1	0
6	G	105	PEG	1	0
6	L	104	PEG	1	0
4	A	104	GAI	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	72/83 (86%)	-0.19	3 (4%) 36 40	9, 14, 34, 50	0
1	B	70/83 (84%)	-0.21	3 (4%) 35 39	9, 14, 37, 50	0
1	C	71/83 (85%)	-0.17	4 (5%) 24 26	10, 14, 41, 87	0
1	D	69/83 (83%)	-0.26	2 (2%) 51 56	9, 15, 40, 50	0
1	E	69/83 (83%)	-0.29	2 (2%) 51 56	9, 14, 37, 47	0
1	F	72/83 (86%)	-0.08	5 (6%) 16 17	9, 14, 37, 49	0
1	G	70/83 (84%)	-0.08	2 (2%) 51 56	9, 15, 38, 72	0
1	H	71/83 (85%)	-0.15	4 (5%) 24 26	9, 13, 42, 58	0
1	I	71/83 (85%)	-0.17	4 (5%) 24 26	9, 15, 41, 56	0
1	J	71/83 (85%)	-0.15	5 (7%) 16 17	9, 15, 51, 75	0
1	K	71/83 (85%)	-0.11	3 (4%) 36 40	9, 15, 42, 68	0
1	L	71/83 (85%)	-0.21	4 (5%) 24 26	10, 15, 41, 56	0
All	All	848/996 (85%)	-0.17	41 (4%) 30 33	9, 14, 43, 87	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	2[A]	SER	7.6
1	J	2[A]	SER	7.4
1	K	2[A]	SER	7.1
1	H	2[A]	SER	6.3
1	L	2[A]	SER	5.7
1	F	1	GLY	5.1
1	D	51	LYS	5.0
1	C	51	LYS	5.0
1	K	51	LYS	4.7
1	B	51	LYS	4.5
1	F	51	LYS	4.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	I	51	LYS	4.2
1	A	51	LYS	3.8
1	D	72	ILE	3.8
1	H	51	LYS	3.7
1	I	71	GLU	3.6
1	C	50	GLY	3.5
1	L	51	LYS	3.5
1	E	51	LYS	3.4
1	J	51	LYS	3.4
1	K	72	ILE	3.3
1	B	72	ILE	3.2
1	I	2[A]	SER	3.0
1	J	71	GLU	3.0
1	A	72	ILE	2.8
1	J	72	ILE	2.8
1	B	71	GLU	2.8
1	F	72	ILE	2.7
1	I	72	ILE	2.7
1	C	4[A]	MET	2.6
1	F	71	GLU	2.6
1	A	71	GLU	2.6
1	L	71	GLU	2.4
1	J	49	ASP	2.4
1	H	71	GLU	2.4
1	C	72	ILE	2.2
1	G	51	LYS	2.2
1	H	72	ILE	2.1
1	F	50	GLY	2.1
1	L	72	ILE	2.1
1	E	4	MET	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GAI	A	105	4/4	0.77	0.17	34,41,51,69	0
4	GAI	F	103	4/4	0.81	0.16	29,36,38,38	0
6	PEG	G	105	7/7	0.82	0.15	36,40,44,44	0
4	GAI	B	103	4/4	0.84	0.20	35,39,40,41	0
4	GAI	E	103	4/4	0.86	0.14	29,35,36,38	0
4	GAI	A	104	4/4	0.86	0.16	34,38,40,40	0
4	GAI	H	105	4/4	0.86	0.13	37,45,54,54	0
4	GAI	G	104	4/4	0.87	0.17	39,42,43,44	0
6	PEG	J	104	7/7	0.88	0.15	40,42,43,44	7
6	PEG	L	104	7/7	0.90	0.13	29,30,32,32	7
3	MPD	D	102	8/8	0.91	0.13	20,28,32,32	0
6	PEG	K	104	7/7	0.91	0.16	44,46,48,48	0
3	MPD	C	102	8/8	0.91	0.16	27,34,38,41	0
4	GAI	C	103	4/4	0.92	0.15	32,37,39,40	0
3	MPD	H	103	8/8	0.93	0.10	24,31,35,36	0
3	MPD	A	103	8/8	0.93	0.14	18,30,32,33	0
3	MPD	J	102	8/8	0.93	0.11	22,29,33,34	0
3	MPD	H	102	8/8	0.94	0.09	20,25,26,29	0
3	MPD	G	102	8/8	0.94	0.10	19,26,30,30	0
3	MPD	B	102	8/8	0.94	0.11	17,25,28,31	0
2	MRD	A	101	8/8	0.95	0.10	15,15,16,16	0
3	MPD	K	102	8/8	0.95	0.09	19,28,30,33	0
2	MRD	E	101	8/8	0.95	0.08	16,17,17,18	0
3	MPD	I	102	8/8	0.95	0.09	20,28,32,34	0
2	MRD	J	101	8/8	0.96	0.09	13,14,15,16	0
3	MPD	L	102	8/8	0.96	0.10	22,29,34,34	0
2	MRD	D	101	8/8	0.96	0.08	16,16,17,18	0
3	MPD	E	102	8/8	0.97	0.08	17,28,33,34	0
2	MRD	F	101	8/8	0.97	0.12	14,15,16,17	0
2	MRD	B	101	8/8	0.97	0.07	15,16,17,17	0
2	MRD	K	101	8/8	0.98	0.11	12,14,15,16	0
2	MRD	A	102	8/8	0.98	0.06	15,18,21,22	0
2	MRD	G	101	8/8	0.98	0.10	12,13,14,15	0
2	MRD	C	101	8/8	0.98	0.06	15,16,16,17	0
2	MRD	I	101	8/8	0.98	0.11	12,13,14,15	0
2	MRD	L	101	8/8	0.98	0.11	12,13,14,14	0
2	MRD	H	101	8/8	0.98	0.08	11,13,15,15	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	CL	F	102	1/1	0.99	0.07	35,35,35,35	0
5	CL	K	103	1/1	1.00	0.03	16,16,16,16	0
5	CL	L	103	1/1	1.00	0.03	18,18,18,18	0
5	CL	H	104	1/1	1.00	0.03	16,16,16,16	0
5	CL	G	103	1/1	1.00	0.03	17,17,17,17	0
5	CL	I	103	1/1	1.00	0.03	15,15,15,15	0
5	CL	J	103	1/1	1.00	0.03	17,17,17,17	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.