



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 6, 2022 – 04:05 PM EST

PDB ID : 7T0S  
Title : Crystal structure of the BCL6 BTB domain in complex with OICR-11864  
Authors : Kuntz, D.A.; Prive, G.G.  
Deposited on : 2021-11-30  
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.31.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	127	<div> <div>5%</div> <div>78%</div> <div>18%</div> <div>.</div> </div>
1	B	127	<div> <div>2%</div> <div>87%</div> <div>7%</div> <div>.</div> </div>
1	C	127	<div> <div>%</div> <div>86%</div> <div>10%</div> <div>.</div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6437 atoms, of which 3128 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Isoform 2 of B-cell lymphoma 6 protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	122	Total	C	H	N	O	S	0	5	0
			1978	626	997	169	178	8			
1	B	122	Total	C	H	N	O	S	0	2	0
			1982	623	1001	171	179	8			
1	C	122	Total	C	H	N	O	S	0	5	0
			2013	632	1022	173	178	8			

There are 15 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	3	GLY	-	expression tag	UNP P41182
A	4	SER	-	expression tag	UNP P41182
A	8	GLN	CYS	engineered mutation	UNP P41182
A	67	ARG	CYS	engineered mutation	UNP P41182
A	84	ASN	CYS	engineered mutation	UNP P41182
B	3	GLY	-	expression tag	UNP P41182
B	4	SER	-	expression tag	UNP P41182
B	8	GLN	CYS	engineered mutation	UNP P41182
B	67	ARG	CYS	engineered mutation	UNP P41182
B	84	ASN	CYS	engineered mutation	UNP P41182
C	3	GLY	-	expression tag	UNP P41182
C	4	SER	-	expression tag	UNP P41182
C	8	GLN	CYS	engineered mutation	UNP P41182
C	67	ARG	CYS	engineered mutation	UNP P41182
C	84	ASN	CYS	engineered mutation	UNP P41182

- Molecule 2 is 3-chloro-5-{1-[2-({5-chloro-2-[(3S)-3-methylmorpholin-4-yl]pyridin-4-yl}amino)-2-oxoethyl]-4-fluoro-1H-pyrrolo[2,3-b]pyridin-3-yl}-2-hydroxybenzamide (three-letter code: E2E) (formula: C<sub>26</sub>H<sub>23</sub>Cl<sub>2</sub>FN<sub>6</sub>O<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
			Total	C	Cl	F	H	N	O		
2	A	1	Total 62	26	2	1	23	6	4	0	0
2	B	1	Total 124	52	4	2	46	12	8	0	1
2	C	1	Total 62	26	2	1	23	6	4	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	Total 14	3	8	3	0	1

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	58	Total	O	0	3
			61	61		
5	B	63	Total	O	0	2
			65	65		
5	C	57	Total	O	0	4
			61	61		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Isoform 2 of B-cell lymphoma 6 protein



- Molecule 1: Isoform 2 of B-cell lymphoma 6 protein



- Molecule 1: Isoform 2 of B-cell lymphoma 6 protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.74Å 72.29Å 55.33Å 90.00° 108.32° 90.00°	Depositor
Resolution (Å)	29.78 – 1.86 55.87 – 1.86	Depositor EDS
% Data completeness (in resolution range)	99.5 (29.78-1.86) 99.8 (55.87-1.86)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.21 (at 1.86Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.212 , 0.249 0.211 , 0.249	Depositor DCC
$R_{free}$ test set	1379 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.3	Xtriage
Anisotropy	0.590	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.45 , 53.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6437	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 70.16 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.2755e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, E2E, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/1019	0.62	0/1377
1	B	0.35	0/1002	0.59	0/1352
1	C	0.36	0/1027	0.61	0/1385
All	All	0.39	0/3048	0.61	0/4114

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	981	997	994	15	0
1	B	981	1001	999	7	0
1	C	991	1022	1010	10	0
2	A	39	23	0	0	0
2	B	78	46	0	0	0
2	C	39	23	0	1	0
3	A	6	8	8	0	0
3	B	6	8	8	0	0
4	A	1	0	0	0	0
5	A	61	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	65	0	0	2	0
5	C	61	0	0	0	0
All	All	3309	3128	3019	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:39:SER:OG	1:B:40:ARG:N	2.34	0.61
1:A:35[A]:VAL:CG1	1:A:71:VAL:HG22	2.33	0.59
1:A:97:LEU:HD13	1:A:102:ILE:HD13	1.87	0.56
1:A:7:SER:OG	1:A:8:GLN:N	2.40	0.55
1:C:35[A]:VAL:CG1	1:C:71:VAL:HG22	2.37	0.54
1:B:98:ARG:NH2	1:C:7:SER:OG	2.41	0.54
1:A:97:LEU:HD21	1:A:105:VAL:HG11	1.91	0.52
1:C:58:TYR:CE1	1:C:62:THR:HG21	2.45	0.52
1:B:7:SER:CA	5:B:305:HOH:O	2.57	0.52
1:C:35[B]:VAL:HG22	1:C:44:ARG:HG2	1.91	0.51
1:C:58:TYR:CZ	1:C:62:THR:HG21	2.46	0.50
1:A:35[A]:VAL:HG12	1:A:71:VAL:HG22	1.95	0.49
1:B:38:VAL:HG22	1:B:74:LEU:HD12	1.96	0.48
1:C:58:TYR:CD1	2:C:201:E2E:C22	2.97	0.48
1:A:66:LYS:HB3	1:A:69:LEU:HD12	1.96	0.47
1:C:35[A]:VAL:HG12	1:C:71:VAL:HG22	1.99	0.45
1:A:58:TYR:CZ	1:A:62:THR:HG21	2.52	0.45
1:C:51:MET:CE	1:C:62:THR:HG23	2.47	0.45
1:A:102:ILE:HD13	1:A:124:PHE:CD2	2.53	0.44
1:A:38:VAL:HG22	1:A:74:LEU:HD12	2.00	0.43
1:B:86:LEU:HD21	1:B:117:VAL:HG11	2.01	0.43
1:A:86:LEU:CD2	1:A:117:VAL:HG11	2.49	0.42
1:A:39:SER:OG	1:A:40:ARG:N	2.52	0.42
1:B:7:SER:N	5:B:305:HOH:O	2.52	0.42
1:C:41:GLU:HG3	1:C:80:PRO:CB	2.50	0.42
1:A:35[B]:VAL:HG22	1:A:44:ARG:HG2	2.01	0.42
1:B:106:MET:HG3	1:B:121:CYS:HB2	2.01	0.42
1:A:35[B]:VAL:HG23	1:A:44:ARG:NH1	2.35	0.42
1:C:69:LEU:HD23	1:C:69:LEU:HA	1.90	0.41
1:A:86:LEU:HD21	1:A:117:VAL:HG11	2.02	0.41
1:A:7:SER:OG	1:A:9:ILE:HG13	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	125/127 (98%)	120 (96%)	5 (4%)	0	100	100
1	B	122/127 (96%)	119 (98%)	3 (2%)	0	100	100
1	C	125/127 (98%)	122 (98%)	3 (2%)	0	100	100
All	All	372/381 (98%)	361 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/115 (98%)	110 (97%)	3 (3%)	44	29
1	B	112/115 (97%)	109 (97%)	3 (3%)	44	29
1	C	114/115 (99%)	111 (97%)	3 (3%)	46	30
All	All	339/345 (98%)	330 (97%)	9 (3%)	43	29

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115	GLU
1	A	119	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	122	ARG
1	B	8	GLN
1	B	40	ARG
1	B	98	ARG
1	C	8	GLN
1	C	13	ARG
1	C	119	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	42	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	B	202	-	5,5,5	0.84	0	5,5,5	1.21	1 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	E2E	A	201	-	40,43,43	2.16	10 (25%)	48,63,63	1.68	7 (14%)
2	E2E	B	201[A]	-	40,43,43	2.13	10 (25%)	48,63,63	1.82	10 (20%)
3	GOL	A	202[A]	-	5,5,5	1.05	0	5,5,5	1.30	1 (20%)
2	E2E	C	201	-	40,43,43	2.17	11 (27%)	48,63,63	1.81	11 (22%)
2	E2E	B	201[B]	-	40,43,43	2.14	10 (25%)	48,63,63	1.75	8 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	202	-	-	2/4/4/4	-
2	E2E	A	201	-	-	0/20/31/31	0/5/5/5
2	E2E	B	201[A]	-	-	4/20/31/31	0/5/5/5
3	GOL	A	202[A]	-	-	0/4/4/4	-
2	E2E	C	201	-	-	1/20/31/31	0/5/5/5
2	E2E	B	201[B]	-	-	2/20/31/31	0/5/5/5

All (41) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	201[B]	E2E	C13-N15	6.85	1.46	1.33
2	C	201	E2E	C13-N15	6.81	1.46	1.33
2	B	201[A]	E2E	C13-N15	6.77	1.45	1.33
2	A	201	E2E	C13-N15	6.68	1.45	1.33
2	C	201	E2E	C05-C06	-5.66	1.39	1.49
2	A	201	E2E	C05-C06	-5.50	1.39	1.49
2	B	201[A]	E2E	C05-C06	-5.45	1.40	1.49
2	B	201[B]	E2E	C05-C06	-5.33	1.40	1.49
2	A	201	E2E	C17-N18	-4.98	1.32	1.38
2	B	201[B]	E2E	C20-N21	4.90	1.46	1.35
2	C	201	E2E	C17-N18	-4.89	1.32	1.38
2	B	201[A]	E2E	C20-N21	4.79	1.46	1.35
2	C	201	E2E	C20-N21	4.76	1.46	1.35
2	A	201	E2E	C20-N21	4.62	1.45	1.35
2	B	201[A]	E2E	C17-N18	-4.54	1.32	1.38
2	B	201[B]	E2E	C17-N18	-4.50	1.32	1.38
2	A	201	E2E	C01-C02	3.20	1.39	1.36
2	C	201	E2E	C01-C02	3.18	1.39	1.36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	201[B]	E2E	C01-C02	3.06	1.39	1.36
2	B	201[A]	E2E	C01-C02	3.04	1.39	1.36
2	C	201	E2E	O14-C13	-2.98	1.18	1.24
2	A	201	E2E	C12-C13	2.88	1.53	1.50
2	B	201[A]	E2E	O14-C13	-2.84	1.18	1.24
2	B	201[B]	E2E	O14-C13	-2.81	1.18	1.24
2	B	201[B]	E2E	C12-C13	2.78	1.53	1.50
2	A	201	E2E	O14-C13	-2.78	1.18	1.24
2	B	201[A]	E2E	C12-C13	2.74	1.53	1.50
2	B	201[B]	E2E	C22-N21	2.64	1.46	1.41
2	C	201	E2E	C22-N21	2.63	1.46	1.41
2	A	201	E2E	C22-N21	2.56	1.46	1.41
2	B	201[A]	E2E	C22-N21	2.46	1.46	1.41
2	C	201	E2E	C12-C13	2.33	1.53	1.50
2	A	201	E2E	C08-CL09	2.22	1.78	1.73
2	A	201	E2E	O36-C20	-2.22	1.18	1.23
2	B	201[B]	E2E	C08-CL09	2.22	1.78	1.73
2	B	201[A]	E2E	O36-C20	-2.21	1.18	1.23
2	C	201	E2E	O36-C20	-2.19	1.18	1.23
2	B	201[B]	E2E	O36-C20	-2.18	1.18	1.23
2	B	201[A]	E2E	C08-CL09	2.06	1.78	1.73
2	C	201	E2E	C08-CL09	2.04	1.78	1.73
2	C	201	E2E	C34-CL35	2.04	1.78	1.73

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	201	E2E	C01-C02-C04	-6.41	118.99	124.18
2	B	201[B]	E2E	C01-C02-C04	-6.40	118.99	124.18
2	B	201[A]	E2E	C01-C02-C04	-6.27	119.10	124.18
2	A	201	E2E	C01-C02-C04	-6.23	119.13	124.18
2	B	201[B]	E2E	C31-N25-C26	-4.29	106.93	114.66
2	C	201	E2E	C39-N38-C37	3.96	121.69	116.77
2	B	201[B]	E2E	C39-N38-C37	3.91	121.63	116.77
2	B	201[A]	E2E	C39-N38-C37	3.91	121.62	116.77
2	A	201	E2E	C39-N38-C37	3.85	121.55	116.77
2	B	201[A]	E2E	C31-N25-C26	-3.83	107.77	114.66
2	B	201[B]	E2E	F03-C02-C04	3.22	122.06	118.56
2	C	201	E2E	C31-N25-C26	-3.16	108.97	114.66
2	B	201[A]	E2E	F03-C02-C04	3.11	121.93	118.56
2	C	201	E2E	C30-O29-C28	3.10	114.27	109.97
2	C	201	E2E	F03-C02-C04	3.09	121.92	118.56

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	E2E	C31-N25-C26	-3.07	109.13	114.66
2	B	201[A]	E2E	C30-O29-C28	3.03	114.17	109.97
2	B	201[A]	E2E	C23-C24-N25	-2.94	118.84	122.31
2	A	201	E2E	F03-C02-C04	2.90	121.70	118.56
2	C	201	E2E	C05-C04-C37	-2.88	105.24	107.54
2	A	201	E2E	C05-C04-C37	-2.75	105.34	107.54
2	B	201[A]	E2E	C05-C04-C37	-2.61	105.46	107.54
2	B	201[A]	E2E	C01-C39-N38	-2.60	120.55	124.58
2	A	201	E2E	C01-C39-N38	-2.53	120.66	124.58
2	B	201[B]	E2E	C01-C39-N38	-2.53	120.66	124.58
3	A	202[A]	GOL	C3-C2-C1	-2.51	101.96	111.70
2	C	201	E2E	C01-C39-N38	-2.48	120.75	124.58
2	C	201	E2E	C12-C13-N15	2.45	122.06	118.29
2	A	201	E2E	O14-C13-N15	-2.42	119.14	122.58
2	B	201[B]	E2E	C05-C04-C37	-2.41	105.61	107.54
2	B	201[A]	E2E	C12-C13-N15	2.28	121.80	118.29
2	B	201[B]	E2E	C23-C24-N25	-2.26	119.64	122.31
2	C	201	E2E	C33-C34-C22	-2.25	118.75	120.06
3	B	202	GOL	C3-C2-C1	-2.22	103.06	111.70
2	B	201[A]	E2E	O14-C13-N15	-2.22	119.43	122.58
2	C	201	E2E	C33-N32-C24	2.21	121.33	117.30
2	C	201	E2E	O14-C13-C12	-2.11	117.76	120.24
2	B	201[B]	E2E	C34-C33-N32	-2.01	121.11	122.84

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	201[A]	E2E	C23-C24-N25-C26
2	B	201[A]	E2E	C23-C24-N25-C31
2	B	201[A]	E2E	N32-C24-N25-C31
2	B	201[B]	E2E	C23-C24-N25-C26
2	C	201	E2E	C23-C24-N25-C26
3	B	202	GOL	O1-C1-C2-O2
3	B	202	GOL	O1-C1-C2-C3
2	B	201[A]	E2E	N32-C24-N25-C26
2	B	201[B]	E2E	N32-C24-N25-C26

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	201	E2E	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




## Ligand E2E B 201 (A)



## Ligand E2E C 201







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	122/127 (96%)	0.03	6 (4%) 29 28	18, 30, 50, 75	0
1	B	122/127 (96%)	-0.16	2 (1%) 72 72	19, 29, 42, 54	0
1	C	122/127 (96%)	-0.13	1 (0%) 86 86	20, 30, 44, 55	0
All	All	366/381 (96%)	-0.08	9 (2%) 57 56	18, 30, 47, 75	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	8	GLN	3.1
1	B	65	LEU	2.9
1	A	7	SER	2.8
1	A	124	PHE	2.5
1	A	13	ARG	2.4
1	B	7	SER	2.2
1	C	13	ARG	2.2
1	A	125	ILE	2.2
1	A	97	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	CL	A	203	1/1	0.86	0.18	61,61,61,61	0
2	E2E	B	201[B]	39/39	0.90	0.15	21,28,39,43	62
2	E2E	B	201[A]	39/39	0.90	0.15	23,29,42,46	62
2	E2E	C	201	39/39	0.91	0.15	21,30,51,52	62
3	GOL	B	202	6/6	0.92	0.09	29,41,54,65	0
2	E2E	A	201	39/39	0.93	0.11	16,28,36,39	0
3	GOL	A	202[A]	6/6	0.94	0.11	21,25,28,28	14

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




### Electron density around E2E B 201 (B):

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around E2E B 201 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around E2E C 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around E2E A 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.