



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 12:23 am BST

PDB ID : 3T2G
Title : Fructose-1,6-bisphosphate aldolase/phosphatase from *Thermoproteus neutrophilus*, Y229F variant with DHAP
Authors : Du, J.; Say, R.; Lue, W.; Fuchs, G.; Einsle, O.
Deposited on : 2011-07-22
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	407	 66% 23% 5% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	13P	A	410	-	-	X	-
3	13P	A	411	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3032 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fructose-1,6-bisphosphate aldolase/phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	383	Total	C	N	O	S	0	0	0
			3005	1936	515	542	12			


There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	229	PHE	TYR	ENGINEERED MUTATION	UNP B1YAL1
A	400	LEU	-	EXPRESSION TAG	UNP B1YAL1
A	401	GLU	-	EXPRESSION TAG	UNP B1YAL1
A	402	HIS	-	EXPRESSION TAG	UNP B1YAL1
A	403	HIS	-	EXPRESSION TAG	UNP B1YAL1
A	404	HIS	-	EXPRESSION TAG	UNP B1YAL1
A	405	HIS	-	EXPRESSION TAG	UNP B1YAL1
A	406	HIS	-	EXPRESSION TAG	UNP B1YAL1
A	407	HIS	-	EXPRESSION TAG	UNP B1YAL1

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		

- Molecule 3 is 1,3-DIHYDROXYACETONEPHOSPHATE (three-letter code: 13P) (formula: C₃H₇O₆P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			10	3	6	1		
3	A	1	Total	C	O	P	0	0
			10	3	6	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	5	Total	O	0	0
			5	5		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fructose-1,6-bisphosphate aldolase/phosphatase



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	112.43Å 112.43Å 151.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.72 – 3.00 47.72 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.72-3.00) 99.9 (47.72-3.00)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	0.18	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.79 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.175 , 0.234 0.174 , 0.237	Depositor DCC
R_{free} test set	505 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	52.1	Xtriage
Anisotropy	0.214	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 37.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.012 for -1/2*h+1/2*k-1/2*l, 1/2*h-1/2*k-1/2*l, -h-k 0.032 for -1/2*h-1/2*k+1/2*l, -1/2*h-1/2*k-1/2*l, h-k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3032	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 13P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/3083	0.75	1/4173 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	208	ARG	NE-CZ-NH2	-5.47	117.57	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3005	0	3003	78	0
2	A	2	0	0	0	0
3	A	20	0	10	5	0
4	A	5	0	0	0	0
All	All	3032	0	3013	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:LEU:O	1:A:76:THR:HG22	1.37	1.18
1:A:141:LEU:HD11	1:A:145:LYS:HE3	1.54	0.89
3:A:410:13P:H32	3:A:411:13P:H31	1.55	0.88
1:A:293:MET:CE	1:A:301:LYS:HD2	2.05	0.87
1:A:72:LEU:O	1:A:76:THR:CG2	2.27	0.78
1:A:379:ASP:HB3	1:A:383:TYR:CE2	2.19	0.77
1:A:105:ASN:OD1	1:A:107:ARG:HG3	1.86	0.75
1:A:107:ARG:HH12	1:A:221:ARG:NH2	1.86	0.73
1:A:133:LYS:HG2	1:A:297:ASP:O	1.89	0.72
1:A:366:ALA:O	1:A:367:LYS:HB2	1.90	0.72
1:A:107:ARG:HH12	1:A:221:ARG:HH22	1.35	0.70
1:A:268:SER:HA	1:A:351:LEU:HD12	1.73	0.69
3:A:410:13P:C3	3:A:411:13P:H31	2.22	0.69
1:A:293:MET:HE1	1:A:301:LYS:HD2	1.75	0.69
1:A:293:MET:HE2	1:A:301:LYS:HD2	1.75	0.68
3:A:410:13P:H32	3:A:411:13P:C3	2.25	0.66
1:A:365:LEU:HA	1:A:367:LYS:O	1.96	0.66
1:A:356:MET:O	1:A:357:GLU:HB2	1.97	0.65
1:A:265:MET:SD	1:A:266:ARG:HG3	2.36	0.65
1:A:265:MET:HB2	1:A:296:PHE:O	1.99	0.63
1:A:293:MET:HE1	1:A:301:LYS:HB2	1.80	0.63
1:A:379:ASP:HB3	1:A:383:TYR:CD2	2.37	0.60
3:A:410:13P:C3	3:A:411:13P:C3	2.79	0.60
1:A:354:GLU:N	1:A:356:MET:O	2.36	0.58
1:A:293:MET:CE	1:A:301:LYS:CD	2.81	0.58
1:A:136:PRO:HA	1:A:235:PRO:HG3	1.85	0.57
1:A:126:ILE:HG22	1:A:240:ARG:HG2	1.87	0.57
1:A:60:HIS:CD2	1:A:69:ILE:HG13	2.40	0.56
1:A:47:VAL:O	1:A:287:ILE:HA	2.07	0.55
1:A:293:MET:HE1	1:A:301:LYS:CD	2.37	0.54
1:A:27:GLU:HG2	1:A:285:ARG:HH22	1.73	0.54
1:A:272:PRO:HG3	1:A:336:ILE:HB	1.91	0.53
1:A:49:ASN:ND2	1:A:290:GLY:H	2.07	0.53
1:A:366:ALA:O	1:A:367:LYS:CB	2.58	0.52
1:A:171:LEU:HD23	1:A:219:VAL:HG23	1.92	0.51
1:A:108:GLY:C	1:A:230:VAL:HG21	2.31	0.51
1:A:293:MET:HE2	1:A:301:LYS:CD	2.41	0.50
1:A:368:VAL:HG12	1:A:369:LYS:N	2.26	0.50
1:A:129:PHE:HB2	1:A:237:LEU:HB3	1.93	0.50
1:A:371:TYR:CD2	1:A:371:TYR:N	2.80	0.49
1:A:298:GLY:HA3	1:A:299:PRO:C	2.33	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:365:LEU:CA	1:A:367:LYS:O	2.61	0.49
1:A:4:THR:OG1	1:A:117:GLU:HG3	2.13	0.48
1:A:5:VAL:HG12	1:A:238:LEU:HD11	1.94	0.48
1:A:175:GLU:HB3	1:A:177:LYS:HE3	1.96	0.47
1:A:22:HIS:HB3	1:A:25:MET:HG3	1.95	0.47
1:A:194:ILE:HD13	1:A:202:ILE:HD11	1.96	0.47
1:A:181:LEU:HD13	1:A:190:LEU:HA	1.95	0.47
1:A:331:GLN:O	1:A:335:GLN:HG3	2.15	0.47
1:A:63:GLY:HA2	1:A:117:GLU:HG3	1.95	0.47
1:A:136:PRO:HG2	1:A:232:LYS:HB2	1.96	0.47
1:A:92:GLY:O	1:A:95:GLN:HB2	2.15	0.46
1:A:221:ARG:NE	1:A:221:ARG:H	2.14	0.45
1:A:138:ALA:HB1	1:A:261:VAL:HG11	1.99	0.45
1:A:364:ILE:HG13	1:A:364:ILE:H	1.48	0.45
1:A:230:VAL:HG12	1:A:231:GLY:N	2.31	0.45
1:A:298:GLY:CA	1:A:299:PRO:O	2.65	0.44
1:A:176:HIS:N	1:A:176:HIS:ND1	2.66	0.44
1:A:201:ILE:HG22	1:A:219:VAL:HG11	1.99	0.44
1:A:365:LEU:C	1:A:367:LYS:N	2.71	0.44
1:A:356:MET:O	1:A:357:GLU:CB	2.64	0.44
1:A:382:LYS:HA	1:A:385:GLU:HB2	1.98	0.44
1:A:27:GLU:CG	1:A:285:ARG:HH22	2.30	0.44
1:A:298:GLY:CA	1:A:299:PRO:C	2.87	0.43
3:A:410:13P:O3	3:A:411:13P:H32	2.18	0.43
1:A:379:ASP:HB3	1:A:383:TYR:HE2	1.79	0.43
1:A:32:LYS:HG2	1:A:76:THR:HB	2.01	0.43
1:A:185:GLU:OE2	1:A:208:ARG:NH2	2.37	0.43
1:A:141:LEU:HD12	1:A:141:LEU:O	2.19	0.42
1:A:203:ARG:HH12	1:A:221:ARG:HH12	1.68	0.42
1:A:27:GLU:HG2	1:A:285:ARG:NH2	2.35	0.42
1:A:149:ASP:HA	1:A:150:PRO:HD3	1.74	0.41
1:A:136:PRO:CG	1:A:232:LYS:HB2	2.50	0.41
1:A:316:GLY:HA3	1:A:317:GLY:HA2	1.81	0.41
1:A:319:ASP:C	1:A:319:ASP:OD1	2.58	0.41
1:A:384:ILE:O	1:A:388:VAL:HG23	2.19	0.41
1:A:379:ASP:O	1:A:380:ARG:C	2.59	0.41
1:A:182:LYS:HD2	1:A:208:ARG:HH21	1.86	0.41
1:A:265:MET:HB3	1:A:296:PHE:CE2	2.56	0.40
1:A:18:HIS:O	1:A:133:LYS:NZ	2.47	0.40
1:A:199:ARG:HD3	1:A:200:TYR:CE2	2.57	0.40
1:A:5:VAL:HG11	1:A:238:LEU:HD21	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:ALA:O	1:A:102:PHE:CB	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	379/407 (93%)	332 (88%)	34 (9%)	13 (3%)	3	20

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	101	ALA
1	A	102	PHE
1	A	103	SER
1	A	354	GLU
1	A	365	LEU
1	A	367	LYS
1	A	232	LYS
1	A	357	GLU
1	A	363	LYS
1	A	386	ALA
1	A	359	THR
1	A	104	GLY
1	A	234	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/331 (94%)	282 (91%)	29 (9%)	9	33

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	ASN
1	A	76	THR
1	A	78	LYS
1	A	95	GLN
1	A	99	LYS
1	A	100	ASP
1	A	113	VAL
1	A	115	GLU
1	A	117	GLU
1	A	127	ILE
1	A	161	SER
1	A	176	HIS
1	A	182	LYS
1	A	219	VAL
1	A	221	ARG
1	A	232	LYS
1	A	233	ASP
1	A	234	ASP
1	A	238	LEU
1	A	243	SER
1	A	284	GLU
1	A	286	ARG
1	A	331	GLN
1	A	341	ARG
1	A	355	GLU
1	A	367	LYS
1	A	369	LYS
1	A	382	LYS
1	A	389	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	49	ASN
1	A	95	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	13P	A	411	-	9,9,9	0.53	0	10,12,12	1.30	2 (20%)
3	13P	A	410	2	9,9,9	0.60	0	10,12,12	1.02	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	13P	A	411	-	-	2/7/8/8	-
3	13P	A	410	2	-	1/7/8/8	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	A	411	13P	O3P-P-O2P	2.16	115.88	107.64
3	A	411	13P	O2P-P-O1	-2.08	101.19	106.73

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	411	13P	O1-C1-C2-O2
3	A	411	13P	O2-C2-C3-O3
3	A	410	13P	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	411	13P	5	0
3	A	410	13P	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	383/407 (94%)	-0.45	6 (1%) 72 44	25, 37, 69, 78	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	230	VAL	2.5
1	A	101	ALA	2.4
1	A	100	ASP	2.3
1	A	99	LYS	2.2
1	A	358	TYR	2.1
1	A	357	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	13P	A	410	10/10	0.93	0.17	64,70,74,74	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MG	A	409	1/1	0.93	0.08	50,50,50,50	0
3	13P	A	411	10/10	0.94	0.21	37,38,40,41	10
2	MG	A	408	1/1	0.97	0.04	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.