



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 22, 2020 – 11:17 AM BST

PDB ID : 5T20
Title : Crystal Structure of Tarin Lectin bound to Trimannose
Authors : Pereira, P.R.; Meagher, J.L.; Stuckey, J.A.
Deposited on : 2016-08-22
Resolution : 1.91 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)
RSRZ outliers	127900	7793 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	110	<div> <div>0%</div> <div> <div></div> <div>88%</div> <div>10%</div> <div></div> </div> <div></div> </div>
1	C	110	<div> <div>3%</div> <div> <div></div> <div>96%</div> <div></div> </div> <div></div> </div>
1	E	110	<div> <div>4%</div> <div> <div></div> <div>95%</div> <div></div> </div> <div></div> </div>
1	G	110	<div> <div>3%</div> <div> <div></div> <div>86%</div> <div>12%</div> <div></div> </div> <div></div> </div>
1	I	110	<div> <div>2%</div> <div> <div></div> <div>93%</div> <div>5%</div> <div></div> </div> <div></div> </div>
1	K	110	<div> <div>3%</div> <div> <div></div> <div>95%</div> <div></div> </div> <div></div> </div>


Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	M	110	<div> <div>4%</div> <div>90%</div> <div>8%</div> <div>.</div> </div>
1	O	110	<div> <div>%</div> <div>92%</div> <div>6%</div> <div>.</div> </div>
2	B	111	<div> <div>%</div> <div>90%</div> <div>9%</div> <div>.</div> </div>
2	D	111	<div> <div>95%</div> <div>.</div> <div>.</div> </div>
2	F	111	<div> <div>2%</div> <div>92%</div> <div>7%</div> <div>.</div> </div>
2	H	111	<div> <div>2%</div> <div>93%</div> <div>6%</div> <div>.</div> </div>
2	J	111	<div> <div>%</div> <div>95%</div> <div>5%</div> <div>.</div> </div>
2	L	111	<div> <div>%</div> <div>88%</div> <div>11%</div> <div>.</div> </div>
2	N	111	<div> <div>%</div> <div>95%</div> <div>5%</div> <div>.</div> </div>
2	P	111	<div> <div>%</div> <div>89%</div> <div>10%</div> <div>.</div> </div>
3	Q	3	<div> <div>67%</div> <div>33%</div> </div>
3	R	3	<div> <div>33%</div> <div>67%</div> </div>
3	S	3	<div> <div>33%</div> <div>67%</div> </div>
3	T	3	<div> <div>100%</div> </div>
3	U	3	<div> <div>33%</div> <div>67%</div> </div>
3	W	3	<div> <div>33%</div> <div>67%</div> </div>
3	X	3	<div> <div>33%</div> <div>67%</div> </div>
3	Y	3	<div> <div>100%</div> </div>
3	Z	3	<div> <div>67%</div> <div>33%</div> </div>
3	a	3	<div> <div>33%</div> <div>67%</div> </div>
3	b	3	<div> <div>100%</div> </div>
3	c	3	<div> <div>100%</div> </div>
3	d	3	<div> <div>100%</div> </div>
3	e	3	<div> <div>33%</div> <div>67%</div> </div>
3	f	3	<div> <div>33%</div> <div>67%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	V	2	 100%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 14952 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lectin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	108	Total	C	N	O	S	0	0	0
			830	522	145	160	3			
1	C	109	Total	C	N	O	S	0	0	0
			830	523	143	161	3			
1	E	109	Total	C	N	O	S	0	0	0
			829	522	143	161	3			
1	G	108	Total	C	N	O	S	0	0	0
			830	522	145	160	3			
1	I	108	Total	C	N	O	S	0	0	0
			824	519	142	160	3			
1	K	108	Total	C	N	O	S	0	0	0
			830	522	145	160	3			
1	M	108	Total	C	N	O	S	0	0	0
			830	522	145	160	3			
1	O	108	Total	C	N	O	S	0	0	0
			825	520	142	160	3			

- Molecule 2 is a protein called Lectin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	109	Total	C	N	O	S	0	2	0
			872	555	149	165	3			
2	D	109	Total	C	N	O	S	0	1	0
			868	551	149	165	3			
2	F	109	Total	C	N	O	S	0	1	0
			868	551	149	165	3			
2	H	109	Total	C	N	O	S	0	1	0
			868	551	149	165	3			
2	J	109	Total	C	N	O	S	0	1	0
			866	550	149	164	3			
2	L	109	Total	C	N	O	S	0	1	0
			868	551	149	165	3			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	N	109	Total	C	N	O	S	0	1	0
			868	551	149	165	3			
2	P	109	Total	C	N	O	S	0	1	0
			868	551	149	165	3			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	Q	3	Total	C	O	0	0	0
			34	18	16			
3	R	3	Total	C	O	0	0	0
			34	18	16			
3	S	3	Total	C	O	0	0	0
			34	18	16			
3	T	3	Total	C	O	0	0	0
			34	18	16			
3	U	3	Total	C	O	0	0	0
			34	18	16			
3	W	3	Total	C	O	0	0	0
			34	18	16			
3	X	3	Total	C	O	0	0	0
			34	18	16			
3	Y	3	Total	C	O	0	0	0
			34	18	16			
3	Z	3	Total	C	O	0	0	0
			34	18	16			
3	a	3	Total	C	O	0	0	0
			34	18	16			
3	b	3	Total	C	O	0	0	0
			34	18	16			
3	c	3	Total	C	O	0	0	0
			34	18	16			
3	d	3	Total	C	O	0	0	0
			34	18	16			
3	e	3	Total	C	O	0	0	0
			34	18	16			
3	f	3	Total	C	O	0	0	0
			34	18	16			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	V	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 5 is alpha-D-mannopyranose (three-letter code: MAN) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	F	1	Total	C	O	0	0
			3	2	1		

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	L	1	Total	C	O	0	0
			4	2	2		
6	O	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	60	Total	O	0	0
			60	60		
7	B	59	Total	O	0	0
			59	59		
7	C	50	Total	O	0	0
			50	50		
7	D	52	Total	O	0	0
			52	52		
7	E	62	Total	O	0	0
			62	62		
7	F	53	Total	O	0	0
			53	53		
7	G	56	Total	O	0	0
			56	56		
7	H	49	Total	O	0	0
			49	49		
7	I	64	Total	O	0	0
			64	64		
7	J	42	Total	O	0	0
			42	42		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	K	50	Total 50	O 50	0	0
7	L	49	Total 49	O 49	0	0
7	M	56	Total 56	O 56	0	0
7	N	42	Total 42	O 42	0	0
7	O	51	Total 51	O 51	0	0
7	P	39	Total 39	O 39	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lectin




- Molecule 1: Lectin



- Molecule 1: Lectin



- Molecule 1: Lectin




- Molecule 1: Lectin




- Molecule 1: Lectin





• Molecule 1: Lectin



• Molecule 1: Lectin




• Molecule 2: Lectin




• Molecule 2: Lectin




• Molecule 2: Lectin




• Molecule 2: Lectin




• Molecule 2: Lectin




• Molecule 2: Lectin



• Molecule 2: Lectin



• Molecule 2: Lectin



• Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose



• Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose



• Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose



- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose

Chain T:  100%

MAN1
MAN2
MAN3

- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose

Chain U:  33% 67%

MAN1
MAN2
MAN3

- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose

Chain W:  33% 67%

MAN1
MAN2
MAN3

- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose

Chain X:  33% 67%

MAN1
MAN2
MAN3

- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose

Chain Y:  100%


MAN1
MAN2
MAN3

- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose

Chain Z:  67% 33%

MAN1
MAN2
MAN3

- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose

Chain a:  33% 67%



- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose

Chain b:  100%




- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose

Chain c:  100%



- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose

Chain d:  100%




- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose

Chain e:  33% 67%




- Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose

Chain f:  33% 67%



- Molecule 4: alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose

Chain V:  100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	78.28Å 78.83Å 92.94Å 76.19° 70.44° 59.80°	Depositor
Resolution (Å)	45.05 – 1.91 43.62 – 1.91	Depositor EDS
% Data completeness (in resolution range)	97.1 (45.05-1.91) 97.6 (43.62-1.91)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 1.91Å)	Xtriage
Refinement program	BUSTER 2.11.1	Depositor
R, R_{free}	0.209 , 0.244 0.202 , 0.238	Depositor DCC
R_{free} test set	6809 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	15.3	Xtriage
Anisotropy	0.192	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	14952	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/848	0.67	0/1155
1	C	0.49	0/848	0.67	0/1155
1	E	0.48	0/847	0.64	0/1154
1	G	0.46	0/848	0.66	0/1155
1	I	0.48	0/842	0.66	0/1148
1	K	0.49	0/848	0.67	0/1155
1	M	0.48	0/848	0.67	0/1155
1	O	0.47	0/843	0.66	0/1149
2	B	0.57	0/895	0.65	0/1210
2	D	0.56	0/888	0.65	0/1200
2	F	0.56	0/888	0.65	0/1200
2	H	0.57	0/888	0.67	0/1200
2	J	0.53	0/885	0.67	0/1194
2	L	0.54	0/888	0.66	0/1200
2	N	0.52	0/888	0.67	0/1200
2	P	0.53	0/888	0.67	0/1200
All	All	0.52	0/13880	0.66	0/18830

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	830	0	801	6	0
1	C	830	0	797	2	0
1	E	829	0	795	3	0
1	G	830	0	801	8	0
1	I	824	0	790	4	0
1	K	830	0	801	3	0
1	M	830	0	801	6	0
1	O	825	0	792	5	0
2	B	872	0	832	5	0
2	D	868	0	823	2	0
2	F	868	0	823	4	0
2	H	868	0	823	3	0
2	J	866	0	822	2	0
2	L	868	0	823	8	0
2	N	868	0	823	3	0
2	P	868	0	823	6	0
3	Q	34	0	30	0	0
3	R	34	0	30	0	0
3	S	34	0	30	0	0
3	T	34	0	30	0	0
3	U	34	0	30	0	0
3	W	34	0	30	0	0
3	X	34	0	30	0	0
3	Y	34	0	30	0	0
3	Z	34	0	30	0	0
3	a	34	0	30	0	0
3	b	34	0	30	0	0
3	c	34	0	30	0	0
3	d	34	0	30	0	0
3	e	34	0	30	0	0
3	f	34	0	30	0	0
4	V	23	0	20	0	0
5	F	3	0	3	0	0
6	L	4	0	6	2	0
6	O	4	0	6	0	0
7	A	60	0	0	0	0
7	B	59	0	0	0	0
7	C	50	0	0	1	0
7	D	52	0	0	0	0
7	E	62	0	0	0	0
7	F	53	0	0	0	0
7	G	56	0	0	1	0
7	H	49	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	I	64	0	0	0	0
7	J	42	0	0	0	0
7	K	50	0	0	0	0
7	L	49	0	0	1	0
7	M	56	0	0	0	0
7	N	42	0	0	0	0
7	O	51	0	0	0	0
7	P	39	0	0	1	0
All	All	14952	0	13455	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 59 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:11:THR:HG22	1:G:54:THR:HG22	1.71	0.70
1:A:83:TYR:CD1	1:A:97:GLY:HA3	2.35	0.62
1:O:103:ILE:HB	2:P:98:ALA:HB3	1.82	0.60
2:P:83:SER:OG	7:P:601:HOH:O	2.13	0.60
2:L:19:GLY:HA2	6:L:204:EDO:H22	1.85	0.58

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	106/110 (96%)	103 (97%)	3 (3%)	0	100	100
1	C	107/110 (97%)	104 (97%)	3 (3%)	0	100	100
1	E	107/110 (97%)	105 (98%)	2 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	106/110 (96%)	104 (98%)	2 (2%)	0	100	100
1	I	106/110 (96%)	102 (96%)	4 (4%)	0	100	100
1	K	106/110 (96%)	102 (96%)	4 (4%)	0	100	100
1	M	106/110 (96%)	103 (97%)	3 (3%)	0	100	100
1	O	106/110 (96%)	103 (97%)	3 (3%)	0	100	100
2	B	107/111 (96%)	105 (98%)	2 (2%)	0	100	100
2	D	106/111 (96%)	103 (97%)	3 (3%)	0	100	100
2	F	106/111 (96%)	102 (96%)	4 (4%)	0	100	100
2	H	106/111 (96%)	104 (98%)	2 (2%)	0	100	100
2	J	105/111 (95%)	103 (98%)	2 (2%)	0	100	100
2	L	106/111 (96%)	104 (98%)	2 (2%)	0	100	100
2	N	106/111 (96%)	104 (98%)	2 (2%)	0	100	100
2	P	106/111 (96%)	105 (99%)	1 (1%)	0	100	100
All	All	1698/1768 (96%)	1656 (98%)	42 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	91/93 (98%)	89 (98%)	2 (2%)	52	45
1	C	90/93 (97%)	90 (100%)	0	100	100
1	E	90/93 (97%)	90 (100%)	0	100	100
1	G	91/93 (98%)	90 (99%)	1 (1%)	73	72
1	I	90/93 (97%)	90 (100%)	0	100	100
1	K	91/93 (98%)	91 (100%)	0	100	100
1	M	91/93 (98%)	91 (100%)	0	100	100
1	O	90/93 (97%)	90 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	94/94 (100%)	93 (99%)	1 (1%)	73	72
2	D	93/94 (99%)	93 (100%)	0	100	100
2	F	93/94 (99%)	92 (99%)	1 (1%)	73	72
2	H	93/94 (99%)	92 (99%)	1 (1%)	73	72
2	J	93/94 (99%)	92 (99%)	1 (1%)	73	72
2	L	93/94 (99%)	93 (100%)	0	100	100
2	N	93/94 (99%)	93 (100%)	0	100	100
2	P	93/94 (99%)	92 (99%)	1 (1%)	73	72
All	All	1469/1496 (98%)	1461 (100%)	8 (0%)	88	89

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	93[A]	ARG
2	P	93[A]	ARG
2	H	93[A]	ARG
2	B	93[A]	ARG
1	G	31	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	28	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

47 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MAN	Q	1	3	12,12,12	0.67	0	17,17,17	0.76	0
3	MAN	Q	2	3	11,11,12	1.02	0	15,15,17	0.82	0
3	MAN	Q	3	3	11,11,12	0.80	0	15,15,17	1.72	1 (6%)
3	MAN	R	1	3	12,12,12	0.87	0	17,17,17	0.84	1 (5%)
3	MAN	R	2	3	11,11,12	0.64	0	15,15,17	0.82	0
3	MAN	R	3	3	11,11,12	0.93	0	15,15,17	1.22	2 (13%)
3	MAN	S	1	3	12,12,12	0.67	0	17,17,17	0.67	0
3	MAN	S	2	3	11,11,12	1.27	1 (9%)	15,15,17	0.84	1 (6%)
3	MAN	S	3	3	11,11,12	0.86	0	15,15,17	1.02	1 (6%)
3	MAN	T	1	3	12,12,12	0.80	0	17,17,17	0.94	1 (5%)
3	MAN	T	2	3	11,11,12	0.73	0	15,15,17	0.79	1 (6%)
3	MAN	T	3	3	11,11,12	1.12	1 (9%)	15,15,17	0.69	0
3	MAN	U	1	3	12,12,12	0.85	0	17,17,17	0.80	1 (5%)
3	MAN	U	2	3	11,11,12	0.93	1 (9%)	15,15,17	0.89	1 (6%)
3	MAN	U	3	3	11,11,12	0.83	0	15,15,17	0.91	0
4	MAN	V	1	4	12,12,12	0.73	0	17,17,17	0.92	1 (5%)
4	MAN	V	2	4	11,11,12	0.79	0	15,15,17	0.88	1 (6%)
3	MAN	W	1	3	12,12,12	0.91	0	17,17,17	0.77	0
3	MAN	W	2	3	11,11,12	0.91	0	15,15,17	0.77	1 (6%)
3	MAN	W	3	3	11,11,12	0.69	0	15,15,17	1.91	2 (13%)
3	MAN	X	1	3	12,12,12	0.94	0	17,17,17	0.91	1 (5%)
3	MAN	X	2	3	11,11,12	0.81	0	15,15,17	0.85	0
3	MAN	X	3	3	11,11,12	0.96	1 (9%)	15,15,17	1.16	1 (6%)
3	MAN	Y	1	3	12,12,12	0.75	0	17,17,17	0.87	1 (5%)
3	MAN	Y	2	3	11,11,12	1.11	0	15,15,17	0.72	1 (6%)
3	MAN	Y	3	3	11,11,12	0.92	1 (9%)	15,15,17	1.46	2 (13%)
3	MAN	Z	1	3	12,12,12	0.86	0	17,17,17	0.81	1 (5%)
3	MAN	Z	2	3	11,11,12	0.67	0	15,15,17	0.67	0
3	MAN	Z	3	3	11,11,12	0.74	0	15,15,17	0.83	0
3	MAN	a	1	3	12,12,12	0.73	0	17,17,17	0.67	1 (5%)
3	MAN	a	2	3	11,11,12	0.93	0	15,15,17	0.67	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MAN	a	3	3	11,11,12	0.86	0	15,15,17	1.31	1 (6%)
3	MAN	b	1	3	12,12,12	0.72	0	17,17,17	0.68	0
3	MAN	b	2	3	11,11,12	0.81	0	15,15,17	0.69	0
3	MAN	b	3	3	11,11,12	0.89	0	15,15,17	0.70	0
3	MAN	c	1	3	12,12,12	0.72	0	17,17,17	0.67	1 (5%)
3	MAN	c	2	3	11,11,12	1.19	1 (9%)	15,15,17	0.82	0
3	MAN	c	3	3	11,11,12	0.87	0	15,15,17	1.42	1 (6%)
3	MAN	d	1	3	12,12,12	0.86	0	17,17,17	1.01	1 (5%)
3	MAN	d	2	3	11,11,12	0.71	0	15,15,17	1.01	1 (6%)
3	MAN	d	3	3	11,11,12	0.89	1 (9%)	15,15,17	0.66	0
3	MAN	e	1	3	12,12,12	0.52	0	17,17,17	0.93	1 (5%)
3	MAN	e	2	3	11,11,12	0.89	0	15,15,17	0.64	0
3	MAN	e	3	3	11,11,12	0.96	0	15,15,17	1.33	2 (13%)
3	MAN	f	1	3	12,12,12	0.86	0	17,17,17	0.78	1 (5%)
3	MAN	f	2	3	11,11,12	0.78	0	15,15,17	1.64	1 (6%)
3	MAN	f	3	3	11,11,12	0.60	0	15,15,17	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	Q	1	3	-	0/2/22/22	0/1/1/1
3	MAN	Q	2	3	-	0/2/19/22	0/1/1/1
3	MAN	Q	3	3	-	2/2/19/22	0/1/1/1
3	MAN	R	1	3	-	0/2/22/22	0/1/1/1
3	MAN	R	2	3	-	0/2/19/22	0/1/1/1
3	MAN	R	3	3	-	0/2/19/22	0/1/1/1
3	MAN	S	1	3	-	0/2/22/22	0/1/1/1
3	MAN	S	2	3	-	0/2/19/22	0/1/1/1
3	MAN	S	3	3	-	1/2/19/22	0/1/1/1
3	MAN	T	1	3	-	0/2/22/22	0/1/1/1
3	MAN	T	2	3	-	1/2/19/22	0/1/1/1
3	MAN	T	3	3	-	0/2/19/22	0/1/1/1
3	MAN	U	1	3	-	0/2/22/22	0/1/1/1
3	MAN	U	2	3	-	0/2/19/22	0/1/1/1
3	MAN	U	3	3	-	2/2/19/22	0/1/1/1
4	MAN	V	1	4	-	0/2/22/22	0/1/1/1
4	MAN	V	2	4	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	W	1	3	-	0/2/22/22	0/1/1/1
3	MAN	W	2	3	-	0/2/19/22	0/1/1/1
3	MAN	W	3	3	-	2/2/19/22	0/1/1/1
3	MAN	X	1	3	-	0/2/22/22	0/1/1/1
3	MAN	X	2	3	-	2/2/19/22	0/1/1/1
3	MAN	X	3	3	-	0/2/19/22	0/1/1/1
3	MAN	Y	1	3	-	0/2/22/22	0/1/1/1
3	MAN	Y	2	3	-	0/2/19/22	0/1/1/1
3	MAN	Y	3	3	-	1/2/19/22	0/1/1/1
3	MAN	Z	1	3	-	1/2/22/22	0/1/1/1
3	MAN	Z	2	3	-	0/2/19/22	0/1/1/1
3	MAN	Z	3	3	-	0/2/19/22	0/1/1/1
3	MAN	a	1	3	-	0/2/22/22	0/1/1/1
3	MAN	a	2	3	-	0/2/19/22	0/1/1/1
3	MAN	a	3	3	-	2/2/19/22	0/1/1/1
3	MAN	b	1	3	-	0/2/22/22	0/1/1/1
3	MAN	b	2	3	-	0/2/19/22	0/1/1/1
3	MAN	b	3	3	-	2/2/19/22	0/1/1/1
3	MAN	c	1	3	-	0/2/22/22	0/1/1/1
3	MAN	c	2	3	-	0/2/19/22	0/1/1/1
3	MAN	c	3	3	-	2/2/19/22	0/1/1/1
3	MAN	d	1	3	-	0/2/22/22	0/1/1/1
3	MAN	d	2	3	-	0/2/19/22	0/1/1/1
3	MAN	d	3	3	-	0/2/19/22	0/1/1/1
3	MAN	e	1	3	-	0/2/22/22	0/1/1/1
3	MAN	e	2	3	-	0/2/19/22	0/1/1/1
3	MAN	e	3	3	-	1/2/19/22	0/1/1/1
3	MAN	f	1	3	-	0/2/22/22	0/1/1/1
3	MAN	f	2	3	-	2/2/19/22	0/1/1/1
3	MAN	f	3	3	-	0/2/19/22	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	c	2	MAN	C2-C3	3.03	1.57	1.52
3	S	2	MAN	O5-C1	2.79	1.48	1.43
3	Y	3	MAN	C2-C3	2.48	1.56	1.52
3	T	3	MAN	O5-C1	2.35	1.47	1.43
3	U	2	MAN	O5-C1	2.29	1.47	1.43

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	W	3	MAN	O5-C5-C6	6.70	117.70	107.20
3	Q	3	MAN	O5-C5-C6	6.22	116.95	107.20
3	f	2	MAN	O5-C5-C6	5.74	116.21	107.20
3	c	3	MAN	O5-C5-C6	4.93	114.93	107.20
3	a	3	MAN	O5-C5-C6	4.37	114.05	107.20

There are no chirality outliers.


5 of 21 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	a	3	MAN	O5-C5-C6-O6
3	f	2	MAN	O5-C5-C6-O6
3	a	3	MAN	C4-C5-C6-O6
3	W	3	MAN	O5-C5-C6-O6
3	U	3	MAN	C4-C5-C6-O6



There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











Oligosaccharide Chain W



Oligosaccharide Chain X







5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	EDO	L	204	-	3,3,3	0.42	0	2,2,2	0.61	0
5	MAN	F	500	-	2,2,12	0.48	0	1,1,17	0.41	0
6	EDO	O	204	-	3,3,3	0.53	0	2,2,2	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	L	204	-	-	1/1/1/1	-
6	EDO	O	204	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	L	204	EDO	O1-C1-C2-O2
6	O	204	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	L	204	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	108/110 (98%)	0.04	1 (0%) 84 85	9, 15, 30, 42	1 (0%)
1	C	109/110 (99%)	0.26	3 (2%) 53 56	9, 15, 30, 49	1 (0%)
1	E	109/110 (99%)	0.32	4 (3%) 41 44	10, 17, 29, 70	1 (0%)
1	G	108/110 (98%)	0.13	3 (2%) 53 56	9, 15, 31, 46	1 (0%)
1	I	108/110 (98%)	0.29	2 (1%) 66 69	11, 17, 33, 44	1 (0%)
1	K	108/110 (98%)	0.15	3 (2%) 53 56	10, 16, 30, 44	1 (0%)
1	M	108/110 (98%)	0.14	4 (3%) 41 44	10, 16, 38, 46	1 (0%)
1	O	108/110 (98%)	0.28	1 (0%) 84 85	11, 17, 33, 44	1 (0%)
2	B	110/111 (99%)	0.16	1 (0%) 84 85	9, 15, 28, 38	2 (1%)
2	D	110/111 (99%)	0.16	0 100 100	8, 14, 28, 40	2 (1%)
2	F	110/111 (99%)	0.12	2 (1%) 68 71	8, 14, 27, 50	2 (1%)
2	H	110/111 (99%)	0.24	2 (1%) 68 71	11, 16, 27, 43	2 (1%)
2	J	110/111 (99%)	0.16	1 (0%) 84 85	10, 16, 31, 52	2 (1%)
2	L	110/111 (99%)	0.26	1 (0%) 84 85	11, 17, 31, 47	2 (1%)
2	N	110/111 (99%)	0.32	1 (0%) 84 85	12, 19, 33, 48	2 (1%)
2	P	110/111 (99%)	0.23	1 (0%) 84 85	10, 17, 30, 48	2 (1%)
All	All	1746/1768 (98%)	0.21	30 (1%) 70 72	8, 16, 31, 70	24 (1%)

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	43	TYR	5.7
1	E	109	GLY	4.4
1	C	109	GLY	4.3
1	E	108	PRO	3.2
1	K	67	ASP	3.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MAN	a	3	11/12	0.61	0.34	45,51,52,53	0
3	MAN	c	3	11/12	0.62	0.28	40,44,47,48	0
3	MAN	U	3	11/12	0.70	0.23	30,36,40,42	0
3	MAN	S	3	11/12	0.79	0.25	33,38,40,40	0
3	MAN	Q	3	11/12	0.79	0.19	37,42,45,45	0
3	MAN	Z	2	11/12	0.80	0.29	36,41,47,48	0
3	MAN	T	2	11/12	0.81	0.27	31,36,37,38	0
4	MAN	V	1	12/12	0.82	0.20	21,30,36,37	0
3	MAN	T	3	11/12	0.84	0.20	26,29,31,31	0
4	MAN	V	2	11/12	0.85	0.27	29,33,36,37	0
3	MAN	d	1	12/12	0.85	0.15	28,33,36,36	0
3	MAN	W	3	11/12	0.85	0.17	31,36,38,40	0
3	MAN	d	3	11/12	0.86	0.23	30,32,34,35	0
3	MAN	f	2	11/12	0.86	0.21	35,39,43,45	0
3	MAN	Z	1	12/12	0.87	0.20	27,30,32,37	0
3	MAN	Z	3	11/12	0.87	0.19	25,28,31,32	0
3	MAN	f	1	12/12	0.88	0.19	26,27,30,31	0
3	MAN	X	1	12/12	0.88	0.14	19,26,27,30	0
3	MAN	b	3	11/12	0.89	0.16	24,29,31,32	0
3	MAN	Y	3	11/12	0.89	0.12	19,26,29,30	0
3	MAN	d	2	11/12	0.89	0.22	33,34,40,42	0
3	MAN	T	1	12/12	0.90	0.18	20,27,31,33	0
3	MAN	R	1	12/12	0.90	0.19	18,25,27,30	0
3	MAN	b	2	11/12	0.90	0.16	32,35,38,39	0
3	MAN	R	3	11/12	0.91	0.16	20,24,29,31	0
3	MAN	b	1	12/12	0.91	0.13	24,29,32,33	0
3	MAN	R	2	11/12	0.92	0.18	29,31,37,40	0
3	MAN	X	3	11/12	0.92	0.16	20,24,26,27	0
3	MAN	e	3	11/12	0.92	0.09	23,26,28,28	0
3	MAN	X	2	11/12	0.92	0.14	26,29,33,35	0
3	MAN	f	3	11/12	0.93	0.14	20,23,28,29	0
3	MAN	Y	1	12/12	0.93	0.12	12,14,19,20	0

Continued on next page...

Continued from previous page...



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MAN	e	2	11/12	0.93	0.13	13,16,20,21	0
3	MAN	Y	2	11/12	0.93	0.12	11,13,20,20	0
3	MAN	c	2	11/12	0.94	0.11	11,13,17,21	0
3	MAN	Q	1	12/12	0.94	0.14	12,16,24,30	0
3	MAN	S	2	11/12	0.94	0.12	13,15,18,21	0
3	MAN	U	2	11/12	0.94	0.11	14,15,18,18	0
3	MAN	Q	2	11/12	0.95	0.11	13,14,19,20	0
3	MAN	W	1	12/12	0.95	0.10	13,15,23,27	0
3	MAN	U	1	12/12	0.95	0.10	12,16,19,20	0
3	MAN	W	2	11/12	0.95	0.11	13,14,18,23	0
3	MAN	S	1	12/12	0.96	0.11	13,15,21,27	0
3	MAN	a	1	12/12	0.96	0.10	13,20,30,38	0
3	MAN	a	2	11/12	0.96	0.10	12,15,23,28	0
3	MAN	e	1	12/12	0.97	0.12	10,13,19,20	0
3	MAN	c	1	12/12	0.97	0.10	12,17,26,31	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain R:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain S:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain T:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain U:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around Chain W:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain X:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain Y:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain Z:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	EDO	O	204	4/4	0.65	0.24	33,35,37,42	0
5	MAN	F	500	3/12	0.84	0.12	33,33,35,36	0
6	EDO	L	204	4/4	0.87	0.17	18,23,25,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.