



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 13, 2020 – 09:50 pm BST

PDB ID : 5T2K  
Title : Geobacillus stearothermophilus HemQ with Manganese-Coproporphyrin III  
Authors : Gauss, G.H.; Celis, A.I.; Dubois, J.L.; Peters, J.W.  
Deposited on : 2016-08-23  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	<div> <div>90%</div> <div>6%</div> <div>.</div> </div>
1	B	248	<div> <div>96%</div> <div>.</div> <div>.</div> </div>
1	C	248	<div> <div>%</div> <div>92%</div> <div>5%</div> <div>.</div> </div>
1	D	248	<div> <div>3%</div> <div>89%</div> <div>8%</div> <div>.</div> </div>
1	E	248	<div> <div>%</div> <div>90%</div> <div>6%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	76R	A	301	X	-	-	-
2	76R	B	301	X	-	-	-
2	76R	C	301	X	-	-	-
2	76R	D	301	X	-	-	-
2	76R	E	301	X	-	-	-

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 21683 atoms, of which 9919 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative heme-dependent peroxidase GT50\_08830.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	238	Total	C	H	N	O	S	0	9	0
			3958	1295	1939	338	376	10			
1	B	243	Total	C	H	N	O	S	0	8	0
			4041	1318	1984	344	385	10			
1	C	241	Total	C	H	N	O	S	0	8	0
			4008	1308	1966	339	383	12			
1	D	241	Total	C	H	N	O	S	0	7	0
			3989	1305	1953	339	382	10			
1	E	238	Total	C	H	N	O	S	0	6	0
			3909	1280	1917	330	371	11			

- Molecule 2 is [3,3',3'',3'''-(3,8,13,17-tetramethylporphyrin-2,7,12,18-tetrayl-kappa 4 N 2 1 ,N 22 ,N 23 ,N 24 )tetra(propanoato)(2-)]manganese (three-letter code: 76R) (formula: C<sub>36</sub>H<sub>36</sub>MnN<sub>4</sub>O<sub>8</sub>).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	Mn	N	O	0	0
			81	36	32	1	4	8		
2	B	1	Total	C	H	Mn	N	O	0	0
			81	36	32	1	4	8		
2	C	1	Total	C	H	Mn	N	O	0	0
			81	36	32	1	4	8		
2	D	1	Total	C	H	Mn	N	O	0	0
			81	36	32	1	4	8		
2	E	1	Total	C	H	Mn	N	O	0	0
			81	36	32	1	4	8		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	282	Total	O	0	0
			282	282		
3	B	304	Total	O	0	0
			304	304		
3	C	279	Total	O	0	0
			279	279		
3	D	241	Total	O	0	0
			241	241		
3	E	267	Total	O	0	0
			267	267		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative heme-dependent peroxidase GT50\_08830

Chain A: 



- Molecule 1: Putative heme-dependent peroxidase GT50\_08830

Chain B: 





- Molecule 1: Putative heme-dependent peroxidase GT50\_08830

Chain C: 





- Molecule 1: Putative heme-dependent peroxidase GT50\_08830

Chain D: 



- Molecule 1: Putative heme-dependent peroxidase GT50\_08830

Chain E: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	131.02Å 93.46Å 132.54Å 90.00° 105.31° 90.00°	Depositor
Resolution (Å)	38.40 – 1.80 38.40 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (38.40-1.80) 99.3 (38.40-1.80)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.37 (at 1.79Å)	Xtriage
Refinement program	PHENIX (1.10.1 _2155: ???)	Depositor
R, $R_{free}$	0.154 , 0.176 0.155 , 0.177	Depositor DCC
$R_{free}$ test set	7157 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.0	Xtriage
Anisotropy	0.582	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 53.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	21683	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 76R

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/2078	0.62	0/2813
1	B	0.47	0/2115	0.62	0/2864
1	C	0.47	0/2092	0.62	0/2831
1	D	0.44	0/2087	0.60	0/2827
1	E	0.45	0/2045	0.60	0/2768
All	All	0.46	0/10417	0.61	0/14103

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2019	1939	1952	14	0
1	B	2057	1984	1978	5	0
1	C	2042	1966	1973	11	0
1	D	2036	1953	1963	14	0
1	E	1992	1917	1930	13	0
2	A	49	32	0	0	0
2	B	49	32	0	0	0
2	C	49	32	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	49	32	0	0	0
2	E	49	32	0	0	0
3	A	282	0	0	9	1
3	B	304	0	0	4	0
3	C	279	0	0	5	3
3	D	241	0	0	4	1
3	E	267	0	0	6	0
All	All	11764	9919	9796	54	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:220:GLU:OE1	3:C:401:HOH:O	1.93	0.86
1:C:220:GLU:OE2	1:C:224:ARG:NH1	2.11	0.83
1:E:137:LYS:NZ	3:E:402:HOH:O	2.12	0.82
1:B:45[A]:ASP:OD1	3:B:401:HOH:O	2.05	0.73
1:A:81:GLU:OE2	3:A:401:HOH:O	2.07	0.71
1:B:173:MET:SD	1:B:176:ARG:NH2	2.62	0.71
1:C:240:GLU:OE1	3:C:402:HOH:O	2.08	0.69
1:B:45[B]:ASP:OD2	3:B:402:HOH:O	2.11	0.69
1:E:173:MET:SD	1:E:176:ARG:NH2	2.64	0.69
1:D:224:ARG:HG3	1:E:158[A]:MET:SD	2.32	0.69
1:C:173[A]:MET:SD	1:C:176:ARG:NH2	2.67	0.67
1:B:65:GLY:O	3:B:403:HOH:O	2.13	0.66
1:D:173:MET:SD	1:D:176:ARG:NH2	2.69	0.66
1:A:137:LYS:NZ	3:A:405:HOH:O	2.21	0.64
1:A:220:GLU:OE1	3:A:403:HOH:O	2.16	0.63
1:A:51:GLU:OE2	3:A:404:HOH:O	2.16	0.60
1:D:166:GLU:OE1	1:D:169:ARG:NH1	2.34	0.59
1:A:137:LYS:NZ	3:A:412:HOH:O	2.36	0.58
1:E:166:GLU:OE1	1:E:169:ARG:NH1	2.37	0.57
1:E:151:GLN:OE1	3:E:404:HOH:O	2.18	0.57
1:C:183[B]:THR:HG23	1:C:202:PHE:HB2	1.86	0.56
1:A:19[B]:THR:HG22	3:A:420:HOH:O	2.05	0.55
1:C:151:GLN:NE2	3:C:408:HOH:O	2.38	0.55
1:A:19[B]:THR:HG23	1:A:101:ALA:HB2	1.89	0.54
1:C:220:GLU:CG	1:C:224:ARG:HD3	2.38	0.54
1:E:137:LYS:NZ	3:E:415:HOH:O	2.42	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:LEU:HD21	1:E:233[B]:VAL:HG12	1.91	0.53
1:B:46:GLN:NE2	3:B:408:HOH:O	2.41	0.52
1:A:74:ILE:HD13	1:A:85:ILE:HG21	1.90	0.52
1:E:181:LYS:NZ	3:E:403:HOH:O	2.16	0.52
1:E:28:LEU:O	1:E:33:ARG:NH1	2.44	0.51
1:C:65:GLY:O	3:C:403:HOH:O	2.20	0.50
1:D:19[B]:THR:HG23	1:D:101:ALA:HB2	1.93	0.50
1:D:74:ILE:HD13	1:D:85:ILE:HG21	1.93	0.50
1:A:111:SER:CB	3:A:402:HOH:O	2.54	0.48
1:D:12:TYR:CD2	1:D:107[B]:VAL:HG22	2.49	0.47
1:A:183[B]:THR:HG23	1:A:202:PHE:HB2	1.97	0.47
1:D:19[B]:THR:HG22	3:D:404:HOH:O	2.15	0.47
1:C:159:LEU:O	1:C:164[B]:ARG:NH1	2.48	0.46
1:D:162:GLU:HG3	3:D:437:HOH:O	2.16	0.46
1:A:111:SER:HB3	3:A:402:HOH:O	2.16	0.45
1:A:28:LEU:O	1:A:33[A]:ARG:NH1	2.50	0.45
1:E:65:GLY:O	3:E:406:HOH:O	2.21	0.45
1:D:224:ARG:CG	1:E:158[A]:MET:SD	3.05	0.44
1:C:169:ARG:NH1	3:C:406:HOH:O	2.37	0.44
1:D:65:GLY:O	3:D:401:HOH:O	2.21	0.44
1:A:166:GLU:HG3	3:A:583:HOH:O	2.17	0.43
1:C:220:GLU:HG3	1:C:224:ARG:HD3	2.01	0.43
1:E:2:SER:N	3:E:423:HOH:O	2.54	0.41
1:D:58:HIS:HA	1:D:73:MET:O	2.21	0.41
1:D:183[A]:THR:CG2	1:D:202:PHE:HB2	2.51	0.41
1:D:177:LYS:HE2	3:D:475:HOH:O	2.21	0.40
1:E:58:HIS:HA	1:E:73:MET:O	2.22	0.40
1:D:28:LEU:O	1:D:33:ARG:NH1	2.55	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:601:HOH:O	3:C:601:HOH:O[2_556]	1.40	0.80
3:D:624:HOH:O	3:D:624:HOH:O[2_556]	1.49	0.71
3:C:438:HOH:O	3:C:438:HOH:O[2_556]	1.88	0.32
3:A:662:HOH:O	3:C:554:HOH:O[3_545]	2.19	0.01

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	243/248 (98%)	238 (98%)	5 (2%)	0	100	100
1	B	247/248 (100%)	242 (98%)	5 (2%)	0	100	100
1	C	245/248 (99%)	241 (98%)	4 (2%)	0	100	100
1	D	244/248 (98%)	239 (98%)	5 (2%)	0	100	100
1	E	240/248 (97%)	235 (98%)	5 (2%)	0	100	100
All	All	1219/1240 (98%)	1195 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/218 (100%)	218 (100%)	1 (0%)	88	87
1	B	223/218 (102%)	223 (100%)	0	100	100
1	C	221/218 (101%)	221 (100%)	0	100	100
1	D	220/218 (101%)	220 (100%)	0	100	100
1	E	215/218 (99%)	215 (100%)	0	100	100
All	All	1098/1090 (101%)	1097 (100%)	1 (0%)	93	92

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	112	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	83	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	76R	E	301	1	28,56,56	2.01	9 (32%)	20,90,90	2.31	6 (30%)
2	76R	D	301	1	28,56,56	2.03	9 (32%)	20,90,90	2.98	7 (35%)
2	76R	C	301	1	28,56,56	2.11	9 (32%)	20,90,90	2.74	9 (45%)
2	76R	B	301	1	28,56,56	1.97	8 (28%)	20,90,90	2.88	6 (30%)
2	76R	A	301	1	28,56,56	1.94	8 (28%)	20,90,90	2.55	6 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	76R	E	301	1	1/1/9/9	1/12/60/60	-
2	76R	D	301	1	1/1/9/9	1/12/60/60	-
2	76R	C	301	1	1/1/9/9	2/12/60/60	-
2	76R	B	301	1	1/1/9/9	0/12/60/60	-
2	76R	A	301	1	1/1/9/9	1/12/60/60	-

All (43) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	301	76R	C1B-NB	5.05	1.45	1.36
2	D	301	76R	C1B-NB	4.91	1.45	1.36
2	E	301	76R	C1B-NB	4.73	1.45	1.36
2	B	301	76R	C1B-NB	4.27	1.44	1.36
2	A	301	76R	C1B-NB	4.07	1.44	1.36
2	E	301	76R	C3B-C2B	3.64	1.48	1.37
2	A	301	76R	C3B-C2B	3.61	1.48	1.37
2	C	301	76R	C2A-C3A	3.60	1.48	1.37
2	D	301	76R	C2A-C3A	3.60	1.48	1.37
2	C	301	76R	C3B-C2B	3.54	1.48	1.37
2	D	301	76R	C3B-C2B	3.52	1.48	1.37
2	C	301	76R	C3C-C2C	3.48	1.48	1.37
2	C	301	76R	C3D-C2D	3.46	1.47	1.37
2	D	301	76R	C3D-C2D	3.41	1.47	1.37
2	B	301	76R	C3B-C2B	3.36	1.47	1.37
2	A	301	76R	C3D-C2D	3.35	1.47	1.37
2	E	301	76R	C2A-C3A	3.33	1.47	1.37
2	B	301	76R	C2A-C3A	3.29	1.47	1.37
2	A	301	76R	C2A-C3A	3.26	1.47	1.37
2	E	301	76R	C3D-C2D	3.21	1.47	1.37
2	B	301	76R	C1D-C2D	3.14	1.49	1.42
2	B	301	76R	C3C-C2C	3.11	1.46	1.37
2	B	301	76R	C3D-C2D	3.07	1.46	1.37
2	E	301	76R	C3C-C2C	3.03	1.46	1.37
2	C	301	76R	C1D-C2D	3.01	1.49	1.42
2	D	301	76R	C1A-C2A	3.00	1.49	1.42
2	E	301	76R	C4A-C3A	2.94	1.49	1.42
2	A	301	76R	C3C-C2C	2.93	1.46	1.37
2	A	301	76R	C1D-C2D	2.91	1.49	1.42
2	A	301	76R	C4A-C3A	2.91	1.49	1.42

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	301	76R	C1D-C2D	2.89	1.49	1.42
2	D	301	76R	C1D-C2D	2.89	1.49	1.42
2	C	301	76R	C1A-C2A	2.85	1.49	1.42
2	C	301	76R	C4A-C3A	2.81	1.48	1.42
2	D	301	76R	C3C-C2C	2.79	1.45	1.37
2	D	301	76R	C4A-C3A	2.77	1.48	1.42
2	E	301	76R	C1A-C2A	2.71	1.48	1.42
2	B	301	76R	C4A-C3A	2.70	1.48	1.42
2	B	301	76R	C1A-C2A	2.69	1.48	1.42
2	A	301	76R	C1A-C2A	2.57	1.48	1.42
2	C	301	76R	C1B-CHB	2.01	1.46	1.41
2	E	301	76R	C4B-C3B	2.00	1.47	1.42
2	D	301	76R	C1B-CHB	2.00	1.46	1.41

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	301	76R	CBD-CAD-C3D	-6.88	99.79	112.49
2	C	301	76R	CBD-CAD-C3D	-6.70	100.14	112.49
2	B	301	76R	C1C-C2C-C3C	-6.32	102.60	107.00
2	D	301	76R	CBC-CAC-C3C	-6.23	101.00	112.49
2	A	301	76R	C1C-C2C-C3C	-6.18	102.70	107.00
2	B	301	76R	CBB-CAB-C3B	-6.14	101.17	112.48
2	D	301	76R	C1C-C2C-C3C	-6.13	102.73	107.00
2	C	301	76R	C1C-C2C-C3C	-5.92	102.88	107.00
2	E	301	76R	C1C-C2C-C3C	-5.88	102.90	107.00
2	B	301	76R	CBD-CAD-C3D	-5.72	101.93	112.49
2	B	301	76R	CBC-CAC-C3C	-5.44	102.44	112.49
2	A	301	76R	CBC-CAC-C3C	-5.34	102.64	112.49
2	E	301	76R	CBC-CAC-C3C	-4.71	103.81	112.49
2	A	301	76R	CBB-CAB-C3B	-4.51	104.17	112.48
2	A	301	76R	C1B-C2B-C3B	-4.21	104.07	107.00
2	C	301	76R	CBB-CAB-C3B	-3.80	105.48	112.48
2	D	301	76R	CBB-CAB-C3B	-3.71	105.64	112.48
2	C	301	76R	CAD-CBD-CGD	3.54	118.60	112.67
2	D	301	76R	CAD-CBD-CGD	3.50	118.54	112.67
2	A	301	76R	CBA-CAA-C2A	-3.47	106.09	112.48
2	C	301	76R	CBC-CAC-C3C	-3.44	106.14	112.49
2	C	301	76R	C1B-C2B-C3B	-3.41	104.62	107.00
2	E	301	76R	CBB-CAB-C3B	-3.29	106.42	112.48
2	E	301	76R	CBA-CAA-C2A	-3.22	106.54	112.48
2	B	301	76R	C1B-C2B-C3B	-3.04	104.88	107.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	301	76R	CBD-CAD-C3D	-3.03	106.90	112.49
2	E	301	76R	C1B-C2B-C3B	-3.01	104.90	107.00
2	D	301	76R	C1B-C2B-C3B	-2.97	104.93	107.00
2	C	301	76R	CBA-CAA-C2A	-2.47	107.93	112.48
2	B	301	76R	CMC-C2C-C3C	2.44	129.54	124.94
2	C	301	76R	CMC-C2C-C3C	2.22	129.12	124.94
2	C	301	76R	CMA-C3A-C2A	2.18	129.06	124.94
2	D	301	76R	CMA-C3A-C2A	2.10	128.90	124.94
2	A	301	76R	CAD-CBD-CGD	-2.04	109.25	112.67

All (5) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	E	301	76R	NA
2	D	301	76R	NA
2	C	301	76R	NA
2	B	301	76R	NA
2	A	301	76R	NA

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	301	76R	C3B-CAB-CBB-CGB
2	D	301	76R	C3B-CAB-CBB-CGB
2	A	301	76R	C3B-CAB-CBB-CGB
2	C	301	76R	C2C-C3C-CAC-CBC
2	C	301	76R	C3B-CAB-CBB-CGB

There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.







## Ligand 76R D 301




Bond lengths



Bond angles




Torsions




Rings


## Ligand 76R C 301




Bond lengths



Bond angles




Torsions




Rings


## Ligand 76R B 301




Bond lengths




Bond angles



Torsions



Rings



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	238/248 (95%)	-0.48	1 (0%)	92 90	13, 21, 37, 57	1 (0%)
1	B	243/248 (97%)	-0.48	1 (0%)	92 90	12, 21, 35, 54	0
1	C	241/248 (97%)	-0.49	2 (0%)	86 84	15, 23, 38, 53	0
1	D	241/248 (97%)	-0.27	8 (3%)	46 40	16, 25, 45, 71	0
1	E	238/248 (95%)	-0.42	3 (1%)	77 74	14, 23, 36, 54	0
All	All	1201/1240 (96%)	-0.43	15 (1%)	79 76	12, 23, 38, 71	1 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	125	PRO	4.1
1	D	121	PRO	3.9
1	D	120	ASP	3.9
1	D	122	TYR	3.7
1	D	123	GLN	3.4
1	D	124	ILE	3.3
1	D	125	PRO	3.1
1	B	112	ASN	3.0
1	C	3	GLU	2.8
1	A	125	PRO	2.4
1	E	126	GLU	2.2
1	D	112	ASN	2.1
1	D	127	VAL	2.1
1	C	6	GLN	2.1
1	E	124	ILE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	76R	C	301	49/49	0.94	0.11	26,35,50,56	0
2	76R	D	301	49/49	0.95	0.11	19,31,59,65	0
2	76R	B	301	49/49	0.96	0.11	17,24,39,63	0
2	76R	E	301	49/49	0.97	0.11	15,27,58,79	0
2	76R	A	301	49/49	0.97	0.09	16,23,52,73	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around 76R C 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around 76R D 301:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around 76R B 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 76R E 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around 76R A 301:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers ⓘ

There are no such residues in this entry.