



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 05:56 pm BST

PDB ID : 3T3O  
Title : Molecular basis for the recognition and cleavage of RNA (CUGG) by the bi-functional 5'-3' exo/endoribonuclease RNase J  
Authors : Dorleans, A.; Li de la Sierra-Gallay, I.; Piton, J.; Zig, L.; Gilet, L.; Putzer, H.; Condon, C.  
Deposited on : 2011-07-25  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)
RNA backbone	3102	1008 (2.84-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	562	<div> <div>10%</div> <div>81%</div> <div>16%</div> <div>..</div> </div>
2	B	5	<div> <div>20%</div> <div>20%</div> <div>40%</div> <div>40%</div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4545 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Metal dependent hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	553	Total	C	N	O	S	0	0	0
			4330	2778	761	777	14			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	EXPRESSION TAG	UNP Q72JJ7
A	-6	SER	-	EXPRESSION TAG	UNP Q72JJ7
A	-5	HIS	-	EXPRESSION TAG	UNP Q72JJ7
A	-4	HIS	-	EXPRESSION TAG	UNP Q72JJ7
A	-3	HIS	-	EXPRESSION TAG	UNP Q72JJ7
A	-2	HIS	-	EXPRESSION TAG	UNP Q72JJ7
A	-1	HIS	-	EXPRESSION TAG	UNP Q72JJ7
A	0	HIS	-	EXPRESSION TAG	UNP Q72JJ7
A	1	SER	-	EXPRESSION TAG	UNP Q72JJ7
A	77	ALA	HIS	ENGINEERED MUTATION	UNP Q72JJ7

- Molecule 2 is a RNA chain called O2'methyl-RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	5	Total	C	N	O	P	0	0	0
			111	51	17	38	5			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	96	Total	O	0	0
			96	96		
5	B	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Metal dependent hydrolase



#### • Molecule 2: O2'methyl-RNA



## 4 Data and refinement statistics

Property	Value	Source
Space group	F 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	99.35Å 117.94Å 229.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.76 – 2.50 26.95 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (27.76-2.50) 98.3 (26.95-2.50)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.23 (at 2.50Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, $R_{free}$	0.227 , 0.285 0.237 , 0.299	Depositor DCC
$R_{free}$ test set	1182 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.5	Xtriage
Anisotropy	0.690	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 65.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4545	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OMC, GOL, ZN, OMG, OMU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.40	0/4427	0.65	0/5993
2	B	1.12	0/22	1.63	1/34 (2.9%)
All	All	0.41	0/4449	0.66	1/6027 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	905	U	O4'-C1'-N1	5.88	112.90	108.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4330	0	4428	39	0
2	B	111	0	62	2	0
3	A	1	0	0	0	0
4	A	6	0	8	0	0
5	A	96	0	0	0	0
5	B	1	0	0	0	0
All	All	4545	0	4498	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:248:GLN:HE22	1:A:282:LYS:H	1.22	0.84
1:A:268:MET:HE3	1:A:304:LEU:HB3	1.75	0.68
1:A:54:LEU:HD23	1:A:447:VAL:HG21	1.80	0.63
1:A:487:VAL:HG11	1:A:509:ALA:HB3	1.88	0.55
1:A:241:ALA:HA	1:A:268:MET:HG3	1.88	0.55
1:A:150:HIS:CG	1:A:151:SER:H	2.25	0.55
1:A:319:LEU:HD21	1:A:327:MET:HB3	1.90	0.53
1:A:345:ASN:O	1:A:349:VAL:HG23	2.08	0.53
1:A:175:LEU:HB2	1:A:381:GLU:HG2	1.93	0.51
1:A:465:ASP:HB2	1:A:547:MET:CE	2.42	0.50
1:A:536:LYS:HE3	1:A:540:LYS:HE3	1.95	0.49
1:A:239:THR:HG22	1:A:338:SER:HB3	1.93	0.49
1:A:465:ASP:HB2	1:A:547:MET:HE2	1.95	0.49
1:A:290:LEU:HD11	1:A:327:MET:HG3	1.95	0.48
1:A:408:LYS:HE2	1:A:421:THR:HB	1.95	0.48
1:A:9:HIS:HB3	1:A:433:THR:HG22	1.95	0.48
1:A:13:ILE:HG12	1:A:429:VAL:HG22	1.97	0.46
1:A:254:ALA:HB2	1:A:302:LEU:HD13	1.97	0.46
1:A:335:VAL:HG21	1:A:356:LEU:HD22	1.98	0.46
1:A:293:VAL:HG22	1:A:301:VAL:HG22	1.97	0.45
1:A:336:ILE:HG12	1:A:364:LEU:HD12	1.99	0.45
1:A:377:ALA:HB1	1:A:381:GLU:HB2	1.98	0.44
1:A:240:PHE:CZ	2:B:901:OMC:H5"	2.52	0.44
1:A:200:LEU:HD11	1:A:395:LEU:HB2	2.00	0.44
1:A:23:LYS:HB2	1:A:56:PRO:HD3	2.00	0.44
1:A:228:ILE:HG12	1:A:235:VAL:HG11	1.99	0.43
2:B:903:OMG:H8	2:B:903:OMG:H5"	1.83	0.43
1:A:293:VAL:HA	1:A:296:LEU:HD12	2.00	0.43
1:A:480:LEU:HD12	1:A:486:VAL:HB	2.01	0.43
1:A:4:GLY:HA3	1:A:434:ARG:HH22	1.84	0.42
1:A:72:VAL:HG21	1:A:143:LEU:HD22	2.00	0.42
1:A:25:ILE:HG21	1:A:56:PRO:HG3	2.02	0.42
1:A:397:TRP:CG	1:A:398:HIS:N	2.88	0.42
1:A:244:ILE:HD11	1:A:271:PHE:HB3	2.02	0.41
1:A:311:GLN:HG2	1:A:311:GLN:H	1.55	0.41
1:A:319:LEU:HD11	1:A:327:MET:HE2	2.02	0.41
1:A:238:THR:HA	1:A:305:ALA:O	2.21	0.41

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:PRO:HB2	1:A:45:GLU:H	1.68	0.41
1:A:194:GLY:HA2	1:A:198:VAL:HG23	2.04	0.40
1:A:33:GLU:HG2	1:A:69:LYS:HE3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	551/562 (98%)	517 (94%)	26 (5%)	8 (2%)	10 18

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	44	PRO
1	A	47	GLY
1	A	94	LYS
1	A	322	GLU
1	A	65	ARG
1	A	25	ILE
1	A	232	PRO
1	A	531	TYR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	461/470 (98%)	433 (94%)	28 (6%)	18	36

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	ASP
1	A	35	PHE
1	A	59	ASP
1	A	65	ARG
1	A	133	ASP
1	A	144	ASP
1	A	151	SER
1	A	172	ASP
1	A	219	GLU
1	A	230	ARG
1	A	266	ARG
1	A	273	ARG
1	A	293	VAL
1	A	311	GLN
1	A	313	MET
1	A	337	LEU
1	A	370	LYS
1	A	384	LEU
1	A	395	LEU
1	A	407	PHE
1	A	438	GLU
1	A	463	LEU
1	A	500	ARG
1	A	513	LEU
1	A	514	LYS
1	A	523	LEU
1	A	549	LEU
1	A	553	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	HIS
1	A	136	GLN
1	A	155	ASN
1	A	248	GLN
1	A	379	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	406	ASN

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	4/5 (80%)	3 (75%)	0

All (3) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	902	OMU
2	B	903	OMG
2	B	905	U

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	OMC	B	901	2	19,23,23	1.53	3 (15%)	24,34,34	1.97	5 (20%)
2	OMU	B	902	2	14,22,23	1.62	3 (21%)	14,31,34	1.23	1 (7%)
2	OMG	B	903	2	18,26,27	1.57	3 (16%)	20,38,41	2.33	3 (15%)
2	OMG	B	904	2	18,26,27	1.60	3 (16%)	20,38,41	2.32	4 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OMC	B	901	2	-	4/10/28/28	0/2/2/2
2	OMU	B	902	2	-	0/7/27/28	0/2/2/2
2	OMG	B	903	2	-	2/5/27/28	0/3/3/3
2	OMG	B	904	2	-	0/5/27/28	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	904	OMG	C6-N1	4.70	1.41	1.33
2	B	903	OMG	C6-N1	4.43	1.40	1.33
2	B	902	OMU	O2'-CM2	4.36	1.57	1.42
2	B	901	OMC	P-OP3	-3.87	1.39	1.54
2	B	904	OMG	C2-N1	3.21	1.41	1.35
2	B	901	OMC	C6-N1	3.19	1.39	1.35
2	B	901	OMC	O2'-CM2	3.18	1.53	1.42
2	B	903	OMG	C2-N1	3.02	1.40	1.35
2	B	902	OMU	C4-N3	2.97	1.38	1.33
2	B	903	OMG	C8-N7	-2.63	1.30	1.34
2	B	904	OMG	C8-N7	-2.53	1.30	1.34
2	B	902	OMU	C6-N1	2.14	1.38	1.35

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	904	OMG	C5-C6-N1	-7.49	113.18	123.43
2	B	903	OMG	C5-C6-N1	-7.45	113.23	123.43
2	B	901	OMC	CM2-O2'-C2'	-6.71	96.93	114.52
2	B	904	OMG	C6-N1-C2	5.00	123.87	115.93
2	B	903	OMG	C6-N1-C2	4.94	123.77	115.93
2	B	902	OMU	C5-C4-N3	-4.05	114.39	123.31
2	B	901	OMC	C2-N3-C4	4.03	120.42	116.34
2	B	904	OMG	N3-C2-N1	-2.92	123.32	127.22
2	B	901	OMC	OP3-P-O5'	2.91	114.48	106.73
2	B	903	OMG	N3-C2-N1	-2.80	123.48	127.22
2	B	904	OMG	CM2-O2'-C2'	-2.68	107.50	114.52
2	B	901	OMC	C6-N1-C2	-2.63	117.03	121.20
2	B	901	OMC	OP3-P-OP1	-2.23	101.95	110.68

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	901	OMC	O4'-C1'-N1-C6
2	B	901	OMC	C5'-O5'-P-OP1
2	B	901	OMC	C5'-O5'-P-OP2
2	B	901	OMC	C5'-O5'-P-OP3
2	B	903	OMG	C3'-C4'-C5'-O5'
2	B	903	OMG	O4'-C4'-C5'-O5'

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	901	OMC	1	0
2	B	903	OMG	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	A	701	-	5,5,5	0.49	0	5,5,5	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	701	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	701	GOL	O1-C1-C2-C3
4	A	701	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	553/562 (98%)	0.51	54 (9%) <b>7</b> <b>7</b>	37, 65, 100, 116	0
2	B	1/5 (20%)	4.45	1 (100%) <b>0</b> <b>0</b>	114, 114, 114, 114	0
All	All	554/567 (97%)	0.52	55 (9%) <b>7</b> <b>7</b>	37, 65, 101, 116	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	GLY	5.8
1	A	47	GLY	5.2
1	A	325	ALA	5.0
1	A	327	MET	4.7
1	A	48	MET	4.6
2	B	905	U	4.4
1	A	3	GLN	4.4
1	A	239	THR	4.2
1	A	237	VAL	4.0
1	A	515	ASN	3.6
1	A	295	ASP	3.4
1	A	238	THR	3.4
1	A	322	GLU	3.4
1	A	524	GLU	3.3
1	A	336	ILE	3.0
1	A	532	TYR	3.0
1	A	523	LEU	3.0
1	A	326	LYS	2.9
1	A	305	ALA	2.9
1	A	456	GLY	2.8
1	A	525	ARG	2.8
1	A	302	LEU	2.8
1	A	2	SER	2.8
1	A	313	MET	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	474	LEU	2.7
1	A	303	ILE	2.7
1	A	202	ILE	2.6
1	A	37	LEU	2.6
1	A	236	PHE	2.6
1	A	118	GLY	2.5
1	A	511	GLU	2.5
1	A	519	GLU	2.4
1	A	121	PRO	2.4
1	A	531	TYR	2.4
1	A	169	HIS	2.4
1	A	536	LYS	2.4
1	A	344	GLY	2.3
1	A	168	VAL	2.3
1	A	171	GLY	2.3
1	A	50	GLY	2.2
1	A	201	LEU	2.2
1	A	268	MET	2.2
1	A	304	LEU	2.2
1	A	482	GLY	2.2
1	A	291	GLU	2.2
1	A	312	PRO	2.2
1	A	323	GLY	2.1
1	A	334	THR	2.1
1	A	520	LYS	2.1
1	A	6	PRO	2.1
1	A	497	ALA	2.1
1	A	500	ARG	2.0
1	A	39	GLY	2.0
1	A	28	PHE	2.0
1	A	367	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	OMC	B	901	22/22	0.88	0.21	102,104,105,105	0
2	OMU	B	902	21/22	0.88	0.18	97,102,105,105	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	OMG	B	904	24/25	0.88	0.20	104,109,113,113	0
2	OMG	B	903	24/25	0.89	0.17	102,104,106,106	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	A	701	6/6	0.89	0.25	77,78,78,79	0
3	ZN	A	600	1/1	0.95	0.06	94,94,94,94	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.