



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 03:42 am BST

PDB ID : 3T4B  
Title : Crystal Structure of the HCV IRES pseudoknot domain  
Authors : Berry, K.E.; Waghay, S.; Mortimer, S.A.; Bai, Y.; Doudna, J.A.  
Deposited on : 2011-07-25  
Resolution : 3.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1020 (3.62-3.50)
Clashscore	141614	1100 (3.62-3.50)
RSRZ outliers	127900	1009 (3.64-3.48)
RNA backbone	3102	1008 (4.10-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	84	<div> <div></div> <div>48%</div> <div>37%</div> <div>13%</div> <div>.</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1841 atoms, of which 28 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called HCV IRES pseudoknot domain plus crystallization module.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	84	Total	C	H	N	O	P	1	0	0
			1824	798	28	327	587	84			


- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	17	Total	Ni	0	0
			17	17		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HCV IRES pseudoknot domain plus crystallization module



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	113.39Å 113.39Å 88.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.09 – 3.55 40.09 – 3.55	Depositor EDS
% Data completeness (in resolution range)	99.7 (40.09-3.55) 99.7 (40.09-3.55)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.25 (at 3.57Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, $R_{free}$	0.221 , 0.254 0.215 , 0.243	Depositor DCC
$R_{free}$ test set	368 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	142.3	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.17 , 58.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	1841	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	148.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, A23

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.33	0/1980	0.83	3/3089 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	306	U	C6-N1-C1'	5.34	128.68	121.20
1	A	290	U	C5-C4-O4	-5.26	122.74	125.90
1	A	298	A	N1-C6-N6	-5.25	115.45	118.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1796	28	905	43	0
2	A	17	0	0	0	0
All	All	1813	28	905	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:289:C:H4'	1:A:290:U:OP1	1.70	0.91
1:A:307:G:C6	1:A:330:A:C2	2.84	0.66
1:A:304:C:N3	1:A:305:U:C4	2.69	0.60
1:A:306:U:O2'	1:A:307:G:O5'	2.19	0.59
1:A:329:U:O2'	1:A:330:A:P	2.61	0.59
1:A:329:U:HO2'	1:A:330:A:P	2.27	0.57
1:A:304:C:N4	1:A:305:U:O4	2.39	0.55
1:A:325:C:H2'	1:A:325:C:O2	2.06	0.54
1:A:291:G:H1'	1:A:314:C:C2	2.44	0.53
1:A:306:U:O2'	1:A:307:G:H8	1.93	0.52
1:A:293:C:H2'	1:A:294:U:C6	2.45	0.51
1:A:289:C:O2'	1:A:290:U:H5'	2.11	0.51
1:A:294:U:H2'	1:A:295:G:H2'	1.93	0.51
1:A:328:G:C6	1:A:329:U:C4	2.98	0.50
1:A:125:C:H2'	1:A:126:C:C6	2.47	0.50
1:A:126:C:H2'	1:A:127:U:C6	2.47	0.49
1:A:9:G:C2'	1:A:10:G:H5'	2.43	0.49
1:A:328:G:C2'	1:A:329:U:H5'	2.42	0.48
1:A:290:U:O2'	1:A:315:C:OP2	2.30	0.48
1:A:329:U:C2'	1:A:330:A:O5'	2.62	0.47
1:A:329:U:O2'	1:A:330:A:O5'	2.33	0.47
1:A:306:U:O2'	1:A:307:G:C8	2.69	0.46
1:A:129:C:O2'	1:A:130:C:H5'	2.16	0.46
1:A:292:C:O2'	1:A:293:C:H5'	2.17	0.45
1:A:298:A:N3	1:A:298:A:H2'	2.32	0.44
1:A:328:G:C5	1:A:329:U:C4	3.06	0.44
1:A:304:C:N3	1:A:305:U:C5	2.86	0.44
1:A:311:G:N2	1:A:326:U:C2	2.86	0.44
1:A:331:G:H5'	1:A:332:A23:H5'	1.99	0.44
1:A:288:A:C6	1:A:289:C:C4	3.05	0.43
1:A:288:A:C5	1:A:289:C:C5	3.06	0.43
1:A:306:U:O2'	1:A:307:G:P	2.77	0.43
1:A:303:G:C6	1:A:304:C:C4	3.07	0.43
1:A:6:G:H2'	1:A:7:G:O4'	2.18	0.43
1:A:137:G:C4	1:A:288:A:N1	2.87	0.42
1:A:329:U:O2'	1:A:330:A:OP1	2.31	0.42
1:A:321:A:C2	1:A:322:G:C4	3.07	0.42
1:A:307:G:H5''	1:A:308:C:OP2	2.20	0.42
1:A:134:A:C2'	1:A:135:G:H5'	2.49	0.42
1:A:304:C:C4	1:A:305:U:O4	2.74	0.40
1:A:137:G:N3	1:A:288:A:C2	2.89	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:291:G:C4	1:A:292:C:C5	3.09	0.40
1:A:9:G:H2'	1:A:10:G:H5'	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	83/84 (98%)	16 (19%)	3 (3%)

All (16) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	135	G
1	A	1	G
1	A	10	G
1	A	14	C
1	A	19	U
1	A	289	C
1	A	290	U
1	A	291	G
1	A	297	U
1	A	306	U
1	A	307	G
1	A	324	U
1	A	325	C
1	A	330	A
1	A	331	G
1	A	332	A23



All (3) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	289	C
1	A	290	U
1	A	329	U

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	A23	A	332	1	19,28,29	0.77	0	19,43,46	2.04	4 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	A23	A	332	1	-	2/3/35/36	0/4/4/4

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	332	A23	O3'-PC-O1C	-5.81	100.42	115.76
1	A	332	A23	O2'-C2'-C3'	5.10	114.44	105.08
1	A	332	A23	O3'-C3'-C2'	2.54	109.73	105.08
1	A	332	A23	C5-C6-N6	2.22	123.73	120.35

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	332	A23	O4'-C4'-C5'-O5'
1	A	332	A23	C3'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	332	A23	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 17 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	83/84 (98%)	-0.78	1 (1%) 79 65	109, 147, 197, 259	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	324	U	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	A23	A	332	25/26	0.92	0.14	143,146,180,182	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NI	A	1017	1/1	0.44	0.15	157,157,157,157	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NI	A	1003	1/1	0.64	0.18	130,130,130,130	0
2	NI	A	1015	1/1	0.72	0.24	205,205,205,205	0
2	NI	A	1010	1/1	0.73	0.10	166,166,166,166	0
2	NI	A	1013	1/1	0.83	0.20	179,179,179,179	1
2	NI	A	1011	1/1	0.84	0.22	249,249,249,249	0
2	NI	A	1006	1/1	0.86	0.12	130,130,130,130	0
2	NI	A	1016	1/1	0.86	0.09	164,164,164,164	0
2	NI	A	1008	1/1	0.86	0.06	191,191,191,191	0
2	NI	A	1005	1/1	0.87	0.15	118,118,118,118	0
2	NI	A	1009	1/1	0.88	0.08	191,191,191,191	0
2	NI	A	1012	1/1	0.89	0.26	192,192,192,192	0
2	NI	A	1007	1/1	0.89	0.20	143,143,143,143	0
2	NI	A	1002	1/1	0.89	0.21	126,126,126,126	0
2	NI	A	1001	1/1	0.93	0.20	123,123,123,123	0
2	NI	A	1014	1/1	0.94	0.11	211,211,211,211	0
2	NI	A	1004	1/1	0.96	0.20	134,134,134,134	0

## 6.5 Other polymers

There are no such residues in this entry.