



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 05:17 AM BST

PDB ID : 5T4H
Title : Human DPP4 in complex with ligand 34n
Authors : Scapin, G.
Deposited on : 2016-08-29
Resolution : 2.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3797 (2.64-2.60)
Clashscore	141614	4168 (2.64-2.60)
Ramachandran outliers	138981	4093 (2.64-2.60)
Sidechain outliers	138945	4093 (2.64-2.60)
RSRZ outliers	127900	3731 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	728	
1	B	728	
2	C	2	
2	D	2	
2	E	2	
2	F	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	2	 100%
2	H	2	 100%
2	I	2	 100%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 13686 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl peptidase 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	728	Total	C	N	O	S	0	6	0
			5994	3845	988	1135	26			
1	B	728	Total	C	N	O	S	0	2	0
			5972	3833	982	1130	27			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	THR	-	expression tag	UNP P27487
B	39	THR	-	expression tag	UNP P27487

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	F	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	G	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	H	2	Total	C	N	O	0	0	0
			28	16	2	10			

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	I	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is 2-({2-[(3R)-3-aminopiperidin-1-yl]-3-(but-2-yn-1-yl)-4-oxo-3,4-dihydro-5H-imidazo[2,1-b]purin-5-yl}methyl)benzonitrile (three-letter code: 75J) (formula: $C_{24}H_{24}N_8O$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			33	24	8	1		
4	B	1	Total	C	N	O	0	0
			33	24	8	1		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	715	Total	O	0	0
			715	715		
6	B	672	Total	O	0	0
			672	672		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dipeptidyl peptidase 4




- Molecule 1: Dipeptidyl peptidase 4




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  50% 50%


MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%


MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%


MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  100%

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	118.12Å 126.01Å 137.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	92.91 – 2.61 89.61 – 2.61	Depositor EDS
% Data completeness (in resolution range)	99.8 (92.91-2.61) 99.8 (89.61-2.61)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.99 (at 2.62Å)	Xtriage
Refinement program	BUSTER 2.9.4	Depositor
R, R_{free}	0.148 , 0.226 0.151 , 0.227	Depositor DCC
R_{free} test set	3168 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	27.1	Xtriage
Anisotropy	0.479	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 61.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13686	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, NAG, 75J

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/6191	0.72	0/8419
1	B	0.51	0/6152	0.72	0/8366
All	All	0.51	0/12343	0.72	0/16785

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5994	0	5700	23	0
1	B	5972	0	5689	16	0
2	C	28	0	25	1	0
2	D	28	0	25	0	0
2	E	28	0	25	1	0
2	F	28	0	25	1	0
2	G	28	0	25	0	0
2	H	28	0	25	0	0
2	I	28	0	25	0	0
3	A	42	0	39	0	0
3	B	28	0	26	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	33	0	0	0	0
4	B	33	0	0	0	0
5	A	1	0	0	0	0
6	A	715	0	0	1	0
6	B	672	0	0	0	0
All	All	13686	0	11629	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:LYS:HZ1	1:A:712:HIS:HD2	1.41	0.69
1:A:258:LYS:NZ	1:A:712:HIS:HD2	1.93	0.65
1:A:78:VAL:HG22	1:A:89:PHE:HB2	1.79	0.65
1:A:143:ILE:HD13	1:A:178:PRO:HB2	1.82	0.62
1:B:640:LEU:HD11	1:B:650:GLY:HA3	1.80	0.61
1:B:302:ASP:HB3	1:B:314:GLN:HB2	1.83	0.60
1:B:726:VAL:HG23	1:B:728:VAL:HG23	1.84	0.59
1:B:224:ALA:HB1	1:B:268:PHE:CZ	2.41	0.56
1:B:114:ILE:HG23	1:B:135:TYR:HB3	1.88	0.56
6:A:1399:HOH:O	2:E:2:NAG:H62	2.08	0.54
2:F:1:NAG:O3	2:F:2:NAG:H2	2.09	0.52
1:A:640:LEU:HD11	1:A:650:GLY:HA3	1.91	0.52
1:B:501:ASP:O	1:B:505:GLN:HG2	2.09	0.51
1:A:153:GLN:HE22	1:A:170:ASN:ND2	2.08	0.51
1:B:134:ILE:HD13	1:B:178:PRO:HB3	1.92	0.51
1:B:102:ILE:HD13	1:B:116:LEU:HD22	1.94	0.48
1:A:356:ARG:HD3	1:A:551:CYS:SG	2.54	0.48
1:B:49:LEU:HB3	1:B:749:GLN:HG2	1.96	0.48
1:A:69:LEU:HD13	1:A:107:ILE:HD13	1.95	0.47
1:A:301:CYS:O	1:A:358:ARG:NH1	2.48	0.47
1:A:230:ASP:OD1	1:A:264:PRO:HB3	2.15	0.47
1:B:105:TYR:HB2	1:B:114:ILE:HD11	1.96	0.47
1:B:272:ASN:HD21	1:B:274:ASP:HB2	1.80	0.46
1:B:626:ILE:O	1:B:650:GLY:HA2	2.16	0.45
1:B:109:PRO:HG2	1:B:158:SER:O	2.17	0.45
1:A:60:LEU:HB2	1:A:68:TYR:CD1	2.52	0.44
2:C:1:NAG:H4	2:C:2:NAG:H2	1.63	0.44
1:B:510:PRO:HD3	1:B:569:SER:HB2	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:90:LEU:HD21	1:B:95:PHE:HE2	1.84	0.42
1:A:162:HIS:HD2	1:A:178:PRO:HD3	1.84	0.42
1:A:90:LEU:HD21	1:A:95:PHE:HE2	1.83	0.42
1:A:224:ALA:HB1	1:A:268:PHE:CZ	2.55	0.42
1:A:105:TYR:HB2	1:A:114:ILE:HD11	2.02	0.42
1:A:550:PRO:HD2	1:A:666:TYR:OH	2.20	0.41
1:A:659:TRP:HB3	1:A:667:THR:CG2	2.51	0.41
1:A:107:ILE:HG13	1:A:114:ILE:HD12	2.03	0.41
1:A:726:VAL:HG23	1:A:728:VAL:HG23	2.02	0.41
1:A:547:TYR:C	1:A:547:TYR:CD1	2.94	0.41
1:A:649:CYS:HA	1:A:699:GLU:O	2.21	0.41
1:B:147:ARG:HE	3:B:803:NAG:H83	1.85	0.41
1:A:127:SER:HB3	1:A:211:TYR:CD1	2.56	0.41
1:A:482:LEU:HB2	1:A:494:LEU:HD11	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	732/728 (100%)	703 (96%)	28 (4%)	1 (0%)	51	74
1	B	728/728 (100%)	696 (96%)	31 (4%)	1 (0%)	51	74
All	All	1460/1456 (100%)	1399 (96%)	59 (4%)	2 (0%)	51	74

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	320	GLN
1	A	320	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	659/653 (101%)	638 (97%)	21 (3%)	39	63
1	B	655/653 (100%)	637 (97%)	18 (3%)	44	69
All	All	1314/1306 (101%)	1275 (97%)	39 (3%)	42	66

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	LYS
1	A	71	LYS
1	A	78	VAL
1	A	97	GLU
1	A	133	ASP
1	A	147	ARG
1	A	202	VAL
1	A	263	ASN
1	A	366	LEU
1	A	385	CYS
1	A	471[A]	ARG
1	A	471[B]	ARG
1	A	472	CYS
1	A	520	ASN
1	A	547	TYR
1	A	566	TYR
1	A	627	TRP
1	A	696	LYS
1	A	704	HIS
1	A	745	SER
1	A	764	SER
1	B	51	ASN
1	B	90	LEU
1	B	91	GLU
1	B	97	GLU
1	B	107	ILE
1	B	129	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	138	ASN
1	B	214	LEU
1	B	230	ASP
1	B	244	GLU
1	B	272	ASN
1	B	323	SER
1	B	385	CYS
1	B	450	ASN
1	B	487	ASN
1	B	547	TYR
1	B	627	TRP
1	B	685	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	123	GLN
1	A	169	ASN
1	A	170	ASN
1	A	263	ASN
1	A	572	ASN
1	A	712	HIS
1	A	731	GLN
1	B	169	ASN
1	B	272	ASN
1	B	345	HIS
1	B	487	ASN
1	B	533	HIS
1	B	572	ASN
1	B	712	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	1,2	14,14,15	1.22	1 (7%)	17,19,21	2.09	5 (29%)
2	NAG	C	2	2	14,14,15	1.83	4 (28%)	17,19,21	2.70	7 (41%)
2	NAG	D	1	1,2	14,14,15	1.42	4 (28%)	17,19,21	2.01	4 (23%)
2	NAG	D	2	2	14,14,15	1.67	2 (14%)	17,19,21	2.23	7 (41%)
2	NAG	E	1	1,2	14,14,15	1.55	2 (14%)	17,19,21	2.00	4 (23%)
2	NAG	E	2	2	14,14,15	1.66	2 (14%)	17,19,21	2.39	6 (35%)
2	NAG	F	1	1,2	14,14,15	1.10	1 (7%)	17,19,21	1.90	4 (23%)
2	NAG	F	2	2	14,14,15	2.47	5 (35%)	17,19,21	2.90	7 (41%)
2	NAG	G	1	1,2	14,14,15	1.65	2 (14%)	17,19,21	1.63	4 (23%)
2	NAG	G	2	2	14,14,15	1.82	5 (35%)	17,19,21	1.63	4 (23%)
2	NAG	H	1	1,2	14,14,15	1.38	2 (14%)	17,19,21	2.04	6 (35%)
2	NAG	H	2	2	14,14,15	1.53	2 (14%)	17,19,21	1.70	5 (29%)
2	NAG	I	1	1,2	14,14,15	1.94	4 (28%)	17,19,21	2.16	8 (47%)
2	NAG	I	2	2	14,14,15	2.09	7 (50%)	17,19,21	1.79	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	NAG	D	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
2	NAG	E	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	4/6/23/26	0/1/1/1
2	NAG	F	1	1,2	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	F	2	2	-	2/6/23/26	0/1/1/1
2	NAG	G	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	G	2	2	-	2/6/23/26	0/1/1/1
2	NAG	H	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	H	2	2	-	0/6/23/26	0/1/1/1
2	NAG	I	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	I	2	2	-	0/6/23/26	0/1/1/1

All (43) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	2	NAG	C1-C2	5.30	1.60	1.52
2	G	1	NAG	C1-C2	4.92	1.59	1.52
2	C	2	NAG	C1-C2	4.69	1.59	1.52
2	F	2	NAG	C2-N2	4.48	1.53	1.46
2	I	1	NAG	C1-C2	3.93	1.58	1.52
2	E	2	NAG	C1-C2	3.83	1.58	1.52
2	F	2	NAG	C3-C2	3.82	1.60	1.52
2	I	2	NAG	C4-C3	3.67	1.61	1.52
2	H	2	NAG	C1-C2	3.56	1.57	1.52
2	E	1	NAG	C1-C2	3.55	1.57	1.52
2	H	1	NAG	C1-C2	3.55	1.57	1.52
2	G	2	NAG	C4-C3	3.43	1.61	1.52
2	E	2	NAG	C4-C5	3.31	1.60	1.53
2	D	2	NAG	C3-C2	3.24	1.59	1.52
2	I	1	NAG	O3-C3	3.20	1.50	1.43
2	G	2	NAG	C4-C5	3.18	1.59	1.53
2	D	2	NAG	C1-C2	2.86	1.56	1.52
2	I	2	NAG	C3-C2	2.84	1.58	1.52
2	D	1	NAG	O7-C7	2.84	1.29	1.23
2	I	2	NAG	C4-C5	2.83	1.59	1.53
2	C	1	NAG	C1-C2	2.80	1.56	1.52
2	I	2	NAG	C1-C2	2.69	1.56	1.52
2	F	2	NAG	C4-C3	2.62	1.59	1.52
2	H	2	NAG	C3-C2	2.61	1.58	1.52
2	C	2	NAG	C2-N2	2.51	1.50	1.46
2	I	2	NAG	O5-C5	2.43	1.48	1.43
2	G	2	NAG	C3-C2	2.40	1.57	1.52
2	C	2	NAG	C4-C5	2.33	1.57	1.53
2	I	1	NAG	C6-C5	2.32	1.59	1.51
2	G	2	NAG	O4-C4	2.32	1.48	1.43
2	I	1	NAG	C4-C3	2.28	1.58	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	2	NAG	C7-N2	2.23	1.42	1.34
2	D	1	NAG	C4-C3	2.23	1.58	1.52
2	G	2	NAG	C1-C2	2.22	1.55	1.52
2	D	1	NAG	C7-N2	2.21	1.42	1.34
2	I	2	NAG	C6-C5	2.20	1.59	1.51
2	I	2	NAG	C2-N2	2.14	1.50	1.46
2	F	1	NAG	C1-C2	2.13	1.55	1.52
2	E	1	NAG	C4-C3	2.11	1.57	1.52
2	D	1	NAG	C1-C2	2.08	1.55	1.52
2	C	2	NAG	C4-C3	2.06	1.57	1.52
2	G	1	NAG	C4-C5	2.05	1.57	1.53
2	H	1	NAG	O5-C5	2.02	1.47	1.43

All (76) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	2	NAG	C2-N2-C7	7.68	133.84	122.90
2	D	2	NAG	C1-O5-C5	5.95	120.25	112.19
2	C	2	NAG	C2-N2-C7	5.59	130.86	122.90
2	E	2	NAG	O5-C1-C2	5.57	120.09	111.29
2	C	2	NAG	C4-C3-C2	5.26	118.73	111.02
2	D	1	NAG	O3-C3-C2	-5.11	98.90	109.47
2	F	2	NAG	O5-C1-C2	4.82	118.89	111.29
2	F	2	NAG	C4-C3-C2	4.65	117.83	111.02
2	C	1	NAG	C1-O5-C5	4.63	118.46	112.19
2	E	2	NAG	C2-N2-C7	4.48	129.29	122.90
2	H	1	NAG	C2-N2-C7	4.28	129.00	122.90
2	C	2	NAG	O5-C1-C2	4.22	117.96	111.29
2	E	1	NAG	C3-C4-C5	4.14	117.62	110.24
2	H	1	NAG	C1-C2-N2	4.05	117.41	110.49
2	E	1	NAG	C4-C3-C2	4.04	116.93	111.02
2	G	2	NAG	C4-C3-C2	4.00	116.87	111.02
2	F	1	NAG	C4-C3-C2	3.88	116.70	111.02
2	C	2	NAG	C1-C2-N2	3.87	117.09	110.49
2	I	2	NAG	C3-C4-C5	3.83	117.08	110.24
2	F	1	NAG	O3-C3-C2	-3.77	101.66	109.47
2	E	2	NAG	C1-O5-C5	3.75	117.28	112.19
2	D	1	NAG	C4-C3-C2	3.68	116.42	111.02
2	I	1	NAG	O3-C3-C4	3.65	118.78	110.35
2	F	1	NAG	C1-C2-N2	3.50	116.47	110.49
2	G	1	NAG	C8-C7-N2	-3.48	110.20	116.10
2	F	2	NAG	O7-C7-N2	3.41	128.22	121.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	1	NAG	C8-C7-N2	-3.38	110.38	116.10
2	E	1	NAG	C1-C2-N2	3.36	116.22	110.49
2	F	2	NAG	C1-O5-C5	3.34	116.72	112.19
2	F	1	NAG	C1-O5-C5	3.31	116.67	112.19
2	H	2	NAG	O3-C3-C4	-3.30	102.71	110.35
2	I	1	NAG	O6-C6-C5	3.27	122.52	111.29
2	H	1	NAG	C4-C3-C2	3.11	115.58	111.02
2	C	1	NAG	C8-C7-N2	-3.10	110.85	116.10
2	C	1	NAG	C4-C3-C2	3.10	115.55	111.02
2	I	2	NAG	C2-N2-C7	3.09	127.30	122.90
2	I	1	NAG	O3-C3-C2	3.06	115.79	109.47
2	G	1	NAG	C1-C2-N2	3.05	115.69	110.49
2	I	1	NAG	C4-C3-C2	-3.03	106.57	111.02
2	E	2	NAG	C1-C2-N2	-2.98	105.40	110.49
2	C	1	NAG	O4-C4-C3	-2.89	103.66	110.35
2	D	1	NAG	C8-C7-N2	-2.83	111.30	116.10
2	D	2	NAG	O5-C5-C6	-2.82	102.78	107.20
2	I	1	NAG	O5-C1-C2	2.79	115.69	111.29
2	I	1	NAG	C1-C2-N2	-2.73	105.83	110.49
2	E	2	NAG	O5-C5-C4	2.67	117.32	110.83
2	C	2	NAG	C3-C4-C5	2.63	114.94	110.24
2	D	2	NAG	O3-C3-C2	2.60	114.85	109.47
2	H	2	NAG	C4-C3-C2	2.60	114.82	111.02
2	I	1	NAG	C1-O5-C5	2.56	115.66	112.19
2	D	2	NAG	C2-N2-C7	2.54	126.52	122.90
2	D	2	NAG	O7-C7-N2	2.54	126.62	121.95
2	F	2	NAG	C8-C7-N2	-2.53	111.81	116.10
2	F	2	NAG	C1-C2-N2	2.53	114.81	110.49
2	C	2	NAG	O7-C7-N2	2.53	126.60	121.95
2	D	2	NAG	C8-C7-N2	-2.49	111.89	116.10
2	C	2	NAG	C1-O5-C5	2.47	115.53	112.19
2	E	2	NAG	O4-C4-C3	-2.46	104.67	110.35
2	H	2	NAG	C1-C2-N2	-2.44	106.31	110.49
2	I	2	NAG	O5-C1-C2	-2.40	107.49	111.29
2	C	1	NAG	O3-C3-C2	2.36	114.35	109.47
2	G	2	NAG	C2-N2-C7	-2.35	119.56	122.90
2	G	1	NAG	O7-C7-N2	2.34	126.26	121.95
2	I	2	NAG	O4-C4-C3	2.30	115.66	110.35
2	E	1	NAG	O5-C5-C6	2.29	110.80	107.20
2	H	2	NAG	C1-O5-C5	2.27	115.27	112.19
2	H	1	NAG	O5-C5-C6	2.24	110.71	107.20
2	G	2	NAG	C3-C4-C5	2.17	114.12	110.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	2	NAG	O5-C5-C6	2.15	110.58	107.20
2	D	1	NAG	C2-N2-C7	-2.13	119.87	122.90
2	G	1	NAG	O4-C4-C3	-2.11	105.47	110.35
2	G	2	NAG	O3-C3-C4	2.08	115.15	110.35
2	H	2	NAG	O5-C1-C2	2.06	114.54	111.29
2	D	2	NAG	C6-C5-C4	-2.05	108.21	113.00
2	H	1	NAG	O7-C7-N2	2.03	125.68	121.95
2	I	1	NAG	O4-C4-C5	-2.02	104.28	109.30

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	NAG	C1-C2-N2-C7
2	F	2	NAG	O5-C5-C6-O6
2	D	2	NAG	C4-C5-C6-O6
2	E	2	NAG	O5-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6
2	G	2	NAG	C4-C5-C6-O6
2	E	2	NAG	C8-C7-N2-C2
2	E	2	NAG	O7-C7-N2-C2
2	G	2	NAG	O5-C5-C6-O6
2	F	2	NAG	C4-C5-C6-O6
2	E	2	NAG	C4-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6


There are no ring outliers.


5 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	2	NAG	1	0
2	E	2	NAG	1	0
2	C	1	NAG	1	0
2	C	2	NAG	1	0
2	F	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

















5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAG	B	803	1	14,14,15	1.92	4 (28%)	17,19,21	1.88	3 (17%)
4	75J	B	811	-	30,37,37	1.87	7 (23%)	32,53,53	2.91	14 (43%)
3	NAG	B	810	1	14,14,15	1.87	4 (28%)	17,19,21	1.86	4 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	75J	A	810	-	30,37,37	1.85	6 (20%)	32,53,53	2.26	12 (37%)
3	NAG	A	807	1	14,14,15	2.23	5 (35%)	17,19,21	2.40	7 (41%)
3	NAG	A	806	1	14,14,15	1.02	0	17,19,21	1.78	3 (17%)
3	NAG	A	801	1	14,14,15	1.21	1 (7%)	17,19,21	2.22	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	803	1	-	2/6/23/26	0/1/1/1
4	75J	B	811	-	-	2/8/24/24	0/5/5/5
3	NAG	B	810	1	-	0/6/23/26	0/1/1/1
4	75J	A	810	-	-	1/8/24/24	0/5/5/5
3	NAG	A	807	1	-	2/6/23/26	0/1/1/1
3	NAG	A	806	1	-	0/6/23/26	0/1/1/1
3	NAG	A	801	1	-	0/6/23/26	0/1/1/1

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	810	75J	C13-N8	6.18	1.47	1.38
3	A	807	NAG	C1-C2	5.68	1.60	1.52
4	B	811	75J	C13-N8	4.59	1.45	1.38
4	B	811	75J	C17-C16	4.01	1.51	1.36
4	B	811	75J	C19-N18	-3.91	1.28	1.35
3	B	810	NAG	C4-C5	3.87	1.61	1.53
3	B	803	NAG	C1-C2	3.81	1.58	1.52
4	A	810	75J	C13-C12	3.73	1.47	1.41
4	A	810	75J	C17-C16	3.72	1.50	1.36
4	B	811	75J	C13-C12	3.60	1.47	1.41
3	B	803	NAG	C3-C2	3.17	1.59	1.52
3	B	810	NAG	O4-C4	3.10	1.50	1.43
3	A	801	NAG	C1-C2	2.97	1.56	1.52
4	B	811	75J	C9-N15	2.89	1.39	1.35
3	B	810	NAG	C3-C2	2.85	1.58	1.52
4	A	810	75J	C6-C5	2.80	1.44	1.40
3	A	807	NAG	O5-C1	2.59	1.47	1.43
3	A	807	NAG	C4-C5	2.52	1.58	1.53
3	B	810	NAG	C1-C2	2.38	1.55	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	810	75J	C3-C1	2.35	1.44	1.38
4	B	811	75J	C23-N22	2.34	1.51	1.46
3	B	803	NAG	C4-C5	2.23	1.57	1.53
4	B	811	75J	C11-N18	-2.22	1.31	1.33
3	A	807	NAG	C7-N2	2.09	1.41	1.34
3	B	803	NAG	O5-C5	2.08	1.47	1.43
3	A	807	NAG	O4-C4	2.06	1.47	1.43
4	A	810	75J	C16-N15	2.01	1.46	1.37

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	811	75J	C12-C13-N8	-10.15	109.87	120.30
4	B	811	75J	C13-C12-C11	7.24	124.61	119.96
4	A	810	75J	C12-C13-N8	-6.41	113.72	120.30
4	A	810	75J	C13-C12-C11	5.37	123.41	119.96
3	A	806	NAG	C1-C2-N2	4.89	118.84	110.49
3	B	803	NAG	C1-O5-C5	4.52	118.31	112.19
3	A	807	NAG	C1-O5-C5	4.19	117.87	112.19
3	A	801	NAG	C4-C3-C2	-4.12	104.97	111.02
3	B	810	NAG	C1-O5-C5	4.08	117.72	112.19
4	A	810	75J	C16-C17-N10	-3.95	103.58	106.83
4	B	811	75J	C29-C21-N20	3.90	117.60	112.10
3	A	807	NAG	O4-C4-C5	3.84	118.82	109.30
4	A	810	75J	C25-C26-C27	3.80	116.18	110.85
4	B	811	75J	C16-C17-N10	-3.75	103.75	106.83
3	A	801	NAG	O3-C3-C4	3.67	118.84	110.35
3	B	810	NAG	O4-C4-C5	3.67	118.41	109.30
3	A	807	NAG	O5-C1-C2	3.61	116.99	111.29
4	B	811	75J	C23-N22-C19	-3.58	114.37	122.03
3	A	801	NAG	O4-C4-C3	3.48	118.40	110.35
4	B	811	75J	C25-C26-C27	-3.43	106.05	110.85
3	A	807	NAG	O7-C7-N2	3.32	128.06	121.95
3	A	801	NAG	O5-C1-C2	3.24	116.40	111.29
3	A	807	NAG	O5-C5-C6	3.20	112.22	107.20
3	A	801	NAG	C8-C7-N2	-3.08	110.89	116.10
3	A	807	NAG	C1-C2-N2	3.06	115.72	110.49
3	A	807	NAG	C8-C7-N2	-3.06	110.92	116.10
3	A	806	NAG	O4-C4-C5	3.03	116.82	109.30
4	B	811	75J	C7-N8-C13	-2.96	114.22	117.79
3	B	810	NAG	C8-C7-N2	-2.91	111.17	116.10
4	B	811	75J	C27-N22-C19	-2.88	115.87	122.03

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	810	75J	C29-C21-N20	2.84	116.11	112.10
3	B	803	NAG	C4-C3-C2	2.83	115.17	111.02
4	A	810	75J	C23-N22-C19	-2.76	116.11	122.03
3	B	810	NAG	O5-C5-C6	-2.74	102.90	107.20
3	A	801	NAG	C1-O5-C5	2.59	115.70	112.19
3	B	803	NAG	C2-N2-C7	2.57	126.56	122.90
4	B	811	75J	C21-C29-C30	-2.47	170.88	175.98
4	A	810	75J	C5-C6-C32	-2.42	117.83	120.14
4	A	810	75J	C1-C2-C6	-2.33	117.36	120.36
4	A	810	75J	C23-C24-N28	-2.21	107.02	111.60
4	B	811	75J	C25-C24-N28	-2.21	104.86	111.03
4	A	810	75J	C24-C23-N22	2.20	112.41	109.64
4	B	811	75J	C23-C24-N28	-2.19	107.07	111.60
4	B	811	75J	C3-C1-C2	2.17	123.49	120.19
4	B	811	75J	C12-C11-N18	2.13	115.26	110.87
4	A	810	75J	C12-C11-N18	2.13	115.25	110.87
4	B	811	75J	C25-C24-C23	2.11	113.36	110.65
3	A	806	NAG	O4-C4-C3	-2.10	105.49	110.35
4	A	810	75J	C27-N22-C19	-2.05	117.63	122.03

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	803	NAG	O5-C5-C6-O6
3	A	807	NAG	C4-C5-C6-O6
3	A	807	NAG	O5-C5-C6-O6
3	B	803	NAG	C4-C5-C6-O6
4	B	811	75J	N20-C21-C29-C30
4	B	811	75J	C4-C5-C7-N8
4	A	810	75J	C4-C5-C7-N8


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	803	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	728/728 (100%)	-0.72	1 (0%) 95 95	13, 24, 49, 73	0
1	B	728/728 (100%)	-0.71	1 (0%) 95 95	13, 24, 50, 80	0
All	All	1456/1456 (100%)	-0.72	2 (0%) 95 95	13, 24, 50, 80	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	88	VAL	2.1
1	A	39	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	F	2	14/15	0.57	0.25	82,86,88,88	0
2	NAG	E	2	14/15	0.57	0.36	85,89,92,92	0
2	NAG	C	2	14/15	0.78	0.20	74,78,81,81	0
2	NAG	I	2	14/15	0.85	0.20	47,51,53,54	0
2	NAG	G	2	14/15	0.88	0.14	45,49,52,52	0
2	NAG	D	2	14/15	0.88	0.15	55,59,61,62	0
2	NAG	E	1	14/15	0.89	0.24	60,64,66,66	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	F	1	14/15	0.89	0.11	60,63,66,66	0
2	NAG	C	1	14/15	0.90	0.18	59,63,66,66	0
2	NAG	D	1	14/15	0.92	0.16	39,43,45,46	0
2	NAG	I	1	14/15	0.94	0.16	23,27,30,31	0
2	NAG	H	2	14/15	0.94	0.19	50,54,56,56	0
2	NAG	G	1	14/15	0.95	0.12	38,41,44,44	0
2	NAG	H	1	14/15	0.98	0.10	28,32,34,35	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain E:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain H:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	B	803	14/15	0.76	0.21	74,77,80,80	0
3	NAG	A	801	14/15	0.85	0.16	50,54,57,58	0
3	NAG	A	806	14/15	0.88	0.16	35,39,42,42	0
3	NAG	B	810	14/15	0.89	0.20	44,48,50,50	0
3	NAG	A	807	14/15	0.90	0.15	45,49,51,51	0
5	NA	A	811	1/1	0.93	0.14	34,34,34,34	0
4	75J	A	810	33/33	0.98	0.12	6,13,18,19	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	75J	B	811	33/33	0.98	0.14	16,23,28,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 75J A 810:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 75J B 811:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.