



wwPDB X-ray Structure Validation Summary Report ⓘ

May 15, 2020 – 09:56 pm BST

PDB ID : 5T5N
Title : Calcium-activated chloride channel bestrophin-1 (BEST1), triple mutant: I76A, F80A, F84A; in complex with an Fab antibody fragment, chloride, and calcium
Authors : Long, S.B.; Vaisey, G.
Deposited on : 2016-08-31
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

i

X-RAY DIFFRACTION

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

[illegible]

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	212	
2	J	212	
2	L	212	
2	N	212	
3	G	217	
3	I	217	
3	K	217	
3	M	217	
3	O	217	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CL	A	502	-	-	-	X
4	CL	A	506	-	-	-	X
4	CL	B	503	-	-	-	X
4	CL	C	506	-	-	-	X
4	CL	D	503	-	-	-	X
4	CL	E	506	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 30705 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called bestrophin-1 (BEST1).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	366	Total	C	N	O	S	0	0	0
			2979	1947	484	535	13			
1	B	366	Total	C	N	O	S	0	0	0
			2979	1947	484	535	13			
1	C	366	Total	C	N	O	S	0	0	0
			2979	1947	484	535	13			
1	D	366	Total	C	N	O	S	0	0	0
			2979	1947	484	535	13			
1	E	366	Total	C	N	O	S	0	0	0
			2979	1947	484	535	13			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	76	ALA	ILE	engineered mutation	UNP E1C3A0
A	80	ALA	PHE	engineered mutation	UNP E1C3A0
A	84	ALA	PHE	engineered mutation	UNP E1C3A0
A	406	GLU	-	expression tag	UNP E1C3A0
A	407	GLY	-	expression tag	UNP E1C3A0
A	408	GLU	-	expression tag	UNP E1C3A0
A	409	GLU	-	expression tag	UNP E1C3A0
A	410	PHE	-	expression tag	UNP E1C3A0
B	76	ALA	ILE	engineered mutation	UNP E1C3A0
B	80	ALA	PHE	engineered mutation	UNP E1C3A0
B	84	ALA	PHE	engineered mutation	UNP E1C3A0
B	406	GLU	-	expression tag	UNP E1C3A0
B	407	GLY	-	expression tag	UNP E1C3A0
B	408	GLU	-	expression tag	UNP E1C3A0
B	409	GLU	-	expression tag	UNP E1C3A0
B	410	PHE	-	expression tag	UNP E1C3A0
C	76	ALA	ILE	engineered mutation	UNP E1C3A0
C	80	ALA	PHE	engineered mutation	UNP E1C3A0
C	84	ALA	PHE	engineered mutation	UNP E1C3A0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	406	GLU	-	expression tag	UNP E1C3A0
C	407	GLY	-	expression tag	UNP E1C3A0
C	408	GLU	-	expression tag	UNP E1C3A0
C	409	GLU	-	expression tag	UNP E1C3A0
C	410	PHE	-	expression tag	UNP E1C3A0
D	76	ALA	ILE	engineered mutation	UNP E1C3A0
D	80	ALA	PHE	engineered mutation	UNP E1C3A0
D	84	ALA	PHE	engineered mutation	UNP E1C3A0
D	406	GLU	-	expression tag	UNP E1C3A0
D	407	GLY	-	expression tag	UNP E1C3A0
D	408	GLU	-	expression tag	UNP E1C3A0
D	409	GLU	-	expression tag	UNP E1C3A0
D	410	PHE	-	expression tag	UNP E1C3A0
E	76	ALA	ILE	engineered mutation	UNP E1C3A0
E	80	ALA	PHE	engineered mutation	UNP E1C3A0
E	84	ALA	PHE	engineered mutation	UNP E1C3A0
E	406	GLU	-	expression tag	UNP E1C3A0
E	407	GLY	-	expression tag	UNP E1C3A0
E	408	GLU	-	expression tag	UNP E1C3A0
E	409	GLU	-	expression tag	UNP E1C3A0
E	410	PHE	-	expression tag	UNP E1C3A0

- Molecule 2 is a protein called Fab antibody fragment, light chain (10D10).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	212	Total	C	N	O	S	0	0	0
			1591	990	266	329	6			
2	H	212	Total	C	N	O	S	0	0	0
			1591	990	266	329	6			
2	J	212	Total	C	N	O	S	0	0	0
			1591	990	266	329	6			
2	L	212	Total	C	N	O	S	0	0	0
			1591	990	266	329	6			
2	N	212	Total	C	N	O	S	0	0	0
			1591	990	266	329	6			

- Molecule 3 is a protein called Fab antibody fragment, heavy chain (10D10).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	211	Total	C	N	O	S	0	0	0
			1558	991	254	305	8			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	I	211	Total	C	N	O	S	0	0	0
			1558	991	254	305	8			
3	K	211	Total	C	N	O	S	0	0	0
			1558	991	254	305	8			
3	M	211	Total	C	N	O	S	0	0	0
			1558	991	254	305	8			
3	O	211	Total	C	N	O	S	0	0	0
			1558	991	254	305	8			

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	3	Total	Cl	0	0
			3	3		
4	A	3	Total	Cl	0	0
			3	3		
4	D	3	Total	Cl	0	0
			3	3		
4	C	3	Total	Cl	0	0
			3	3		
4	E	3	Total	Cl	0	0
			3	3		

- Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	K	0	0
			2	2		
5	C	2	Total	K	0	0
			2	2		
5	E	1	Total	K	0	0
			1	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	C	1	Total	C	O	0	0
			6	3	3		
6	D	1	Total	C	O	0	0
			6	3	3		
6	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	1	Total	Ca	0	0
			1	1		
7	A	1	Total	Ca	0	0
			1	1		
7	D	1	Total	Ca	0	0
			1	1		
7	C	1	Total	Ca	0	0
			1	1		
7	E	1	Total	Ca	0	0
			1	1		

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	2	Total 2	O 2	0	0
8	B	2	Total 2	O 2	0	0
8	C	2	Total 2	O 2	0	0
8	D	2	Total 2	O 2	0	0
8	E	2	Total 2	O 2	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: bestrophin-1 (BEST1)

Chain A: 




- Molecule 1: bestrophin-1 (BEST1)


Chain B: 



- Molecule 1: bestrophin-1 (BEST1)


Chain C: 






- Molecule 1: bestrophin-1 (BEST1)

Chain D: 70% 18% 11%




- Molecule 1: bestrophin-1 (BEST1)

Chain E: 71% 18% 11%



- Molecule 2: Fab antibody fragment, light chain (10D10)


Chain F: 2% 82% 18%





- Molecule 2: Fab antibody fragment, light chain (10D10)

Chain H: 84% 15% 1%









- Molecule 2: Fab antibody fragment, light chain (10D10)




- Molecule 2: Fab antibody fragment, light chain (10D10)




- Molecule 2: Fab antibody fragment, light chain (10D10)



- Molecule 3: Fab antibody fragment, heavy chain (10D10)







- Molecule 3: Fab antibody fragment, heavy chain (10D10)

Chain I: 79% 18% .




- Molecule 3: Fab antibody fragment, heavy chain (10D10)

Chain K: 32% 80% 17% .




- Molecule 3: Fab antibody fragment, heavy chain (10D10)

Chain M: 4% 81% 16% .



- Molecule 3: Fab antibody fragment, heavy chain (10D10)

Chain O: 24% 81% 16% .





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	98.24Å 243.88Å 172.05Å 90.00° 93.84° 90.00°	Depositor
Resolution (Å)	37.95 – 3.10 37.95 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (37.95-3.10) 99.9 (37.95-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 3.12Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???), CNS 1.3	Depositor
R, R_{free}	0.218 , 0.242 0.217 , 0.242	Depositor DCC
R_{free} test set	7276 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	81.7	Xtriage
Anisotropy	0.489	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 59.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	30705	wwPDB-VP
Average B, all atoms (Å ²)	101.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, K, CA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/3064	0.50	0/4178
1	B	0.30	0/3064	0.50	0/4178
1	C	0.31	0/3064	0.51	0/4178
1	D	0.30	0/3064	0.50	0/4178
1	E	0.30	0/3064	0.50	1/4178 (0.0%)
2	F	0.28	0/1630	0.54	0/2225
2	H	0.30	0/1630	0.54	0/2225
2	J	0.32	0/1630	0.56	0/2225
2	L	0.34	1/1630 (0.1%)	0.57	0/2225
2	N	0.32	0/1630	0.57	0/2225
3	G	0.28	0/1602	0.53	0/2202
3	I	0.27	0/1602	0.54	1/2202 (0.0%)
3	K	0.26	0/1602	0.53	0/2202
3	M	0.28	0/1602	0.54	0/2202
3	O	0.28	0/1602	0.53	1/2202 (0.0%)
All	All	0.30	1/31480 (0.0%)	0.52	3/43025 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	L	111	ALA	C-N	5.40	1.46	1.34

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	11	LEU	CA-CB-CG	5.44	127.82	115.30
3	O	11	LEU	CA-CB-CG	5.39	127.69	115.30
1	E	202	ARG	NE-CZ-NH1	-5.15	117.73	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	203	ASP	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2979	0	2917	53	0
1	B	2979	0	2917	59	0
1	C	2979	0	2917	52	0
1	D	2979	0	2917	59	0
1	E	2979	0	2917	59	0
2	F	1591	0	1449	23	0
2	H	1591	0	1449	22	0
2	J	1591	0	1449	37	0
2	L	1591	0	1449	24	0
2	N	1591	0	1449	25	0
3	G	1558	0	1441	21	0
3	I	1558	0	1441	30	0
3	K	1558	0	1441	27	0
3	M	1558	0	1441	27	0
3	O	1558	0	1441	24	0
4	A	3	0	0	0	0
4	B	3	0	0	1	0
4	C	3	0	0	0	0
4	D	3	0	0	0	0
4	E	3	0	0	0	0
5	A	2	0	0	0	0
5	C	2	0	0	0	0
5	E	1	0	0	0	0
6	A	6	0	8	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	6	0	8	2	0
6	C	6	0	8	1	0
6	D	6	0	8	1	0
6	E	6	0	8	1	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
7	D	1	0	0	0	0
7	E	1	0	0	0	0
8	A	2	0	0	0	0
8	B	2	0	0	0	0
8	C	2	0	0	0	0
8	D	2	0	0	0	0
8	E	2	0	0	0	0
All	All	30705	0	29075	452	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 452 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:O:38:LYS:HE2	3:O:40:ARG:HD2	1.64	0.79
3:I:38:LYS:HE2	3:I:40:ARG:HD2	1.66	0.77
3:M:38:LYS:HE2	3:M:40:ARG:HD2	1.67	0.77
3:G:38:LYS:HE2	3:G:40:ARG:HD2	1.66	0.76
3:K:38:LYS:HE2	3:K:40:ARG:HD2	1.65	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	364/409 (89%)	349 (96%)	14 (4%)	1 (0%)	41	73
1	B	364/409 (89%)	349 (96%)	14 (4%)	1 (0%)	41	73
1	C	364/409 (89%)	349 (96%)	14 (4%)	1 (0%)	41	73
1	D	364/409 (89%)	349 (96%)	14 (4%)	1 (0%)	41	73
1	E	364/409 (89%)	349 (96%)	14 (4%)	1 (0%)	41	73
2	F	210/212 (99%)	191 (91%)	18 (9%)	1 (0%)	29	64
2	H	210/212 (99%)	193 (92%)	16 (8%)	1 (0%)	29	64
2	J	210/212 (99%)	192 (91%)	17 (8%)	1 (0%)	29	64
2	L	210/212 (99%)	191 (91%)	18 (9%)	1 (0%)	29	64
2	N	210/212 (99%)	191 (91%)	18 (9%)	1 (0%)	29	64
3	G	207/217 (95%)	201 (97%)	5 (2%)	1 (0%)	29	64
3	I	207/217 (95%)	201 (97%)	5 (2%)	1 (0%)	29	64
3	K	207/217 (95%)	201 (97%)	5 (2%)	1 (0%)	29	64
3	M	207/217 (95%)	201 (97%)	5 (2%)	1 (0%)	29	64
3	O	207/217 (95%)	201 (97%)	5 (2%)	1 (0%)	29	64
All	All	3905/4190 (93%)	3708 (95%)	182 (5%)	15 (0%)	34	69

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	202	ARG
1	B	202	ARG
1	C	202	ARG
1	D	202	ARG
1	E	202	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	324/367 (88%)	316 (98%)	8 (2%)	47	75

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	324/367 (88%)	316 (98%)	8 (2%)	47	75
1	C	324/367 (88%)	315 (97%)	9 (3%)	43	73
1	D	324/367 (88%)	316 (98%)	8 (2%)	47	75
1	E	324/367 (88%)	316 (98%)	8 (2%)	47	75
2	F	173/187 (92%)	170 (98%)	3 (2%)	60	83
2	H	173/187 (92%)	169 (98%)	4 (2%)	50	77
2	J	173/187 (92%)	170 (98%)	3 (2%)	60	83
2	L	173/187 (92%)	170 (98%)	3 (2%)	60	83
2	N	173/187 (92%)	170 (98%)	3 (2%)	60	83
3	G	166/190 (87%)	166 (100%)	0	100	100
3	I	166/190 (87%)	166 (100%)	0	100	100
3	K	166/190 (87%)	166 (100%)	0	100	100
3	M	166/190 (87%)	166 (100%)	0	100	100
3	O	166/190 (87%)	166 (100%)	0	100	100
All	All	3315/3720 (89%)	3258 (98%)	57 (2%)	60	83

5 of 57 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	2	THR
1	D	236	TYR
2	L	81	GLU
1	D	37	LEU
1	D	104	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 40 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	89	GLN
2	J	3	GLN
2	N	89	GLN
2	H	198	HIS
2	J	50	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 30 ligands modelled in this entry, 25 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	C	501	-	5,5,5	0.39	0	5,5,5	0.40	0
6	GOL	E	501	-	5,5,5	0.39	0	5,5,5	0.33	0
6	GOL	B	501	-	5,5,5	0.39	0	5,5,5	0.31	0
6	GOL	D	501	-	5,5,5	0.41	0	5,5,5	0.31	0
6	GOL	A	504	-	5,5,5	0.38	0	5,5,5	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	C	501	-	-	1/4/4/4	-
6	GOL	E	501	-	-	1/4/4/4	-
6	GOL	B	501	-	-	1/4/4/4	-
6	GOL	D	501	-	-	1/4/4/4	-
6	GOL	A	504	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	504	GOL	O1-C1-C2-O2
6	B	501	GOL	O1-C1-C2-O2
6	E	501	GOL	O1-C1-C2-O2
6	D	501	GOL	O1-C1-C2-O2
6	C	501	GOL	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	501	GOL	1	0
6	E	501	GOL	1	0
6	B	501	GOL	2	0
6	D	501	GOL	1	0
6	A	504	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	366/409 (89%)	-0.33	1 (0%) 94 88	45, 70, 94, 130	0
1	B	366/409 (89%)	-0.35	0 100 100	50, 72, 95, 137	0
1	C	366/409 (89%)	-0.35	0 100 100	49, 70, 90, 130	0
1	D	366/409 (89%)	-0.37	1 (0%) 94 88	49, 67, 91, 140	0
1	E	366/409 (89%)	-0.36	0 100 100	45, 68, 92, 137	0
2	F	212/212 (100%)	0.06	5 (2%) 59 37	84, 120, 142, 171	0
2	H	212/212 (100%)	-0.03	2 (0%) 84 69	76, 111, 133, 143	0
2	J	212/212 (100%)	1.80	71 (33%) 0 0	88, 182, 231, 245	0
2	L	212/212 (100%)	-0.02	4 (1%) 66 46	67, 105, 124, 141	0
2	N	212/212 (100%)	1.23	57 (26%) 0 0	79, 153, 223, 239	0
3	G	211/217 (97%)	0.27	16 (7%) 13 5	75, 120, 159, 176	0
3	I	211/217 (97%)	-0.16	1 (0%) 91 81	76, 109, 140, 159	0
3	K	211/217 (97%)	1.46	70 (33%) 0 0	87, 148, 235, 247	0
3	M	211/217 (97%)	-0.06	9 (4%) 35 17	64, 103, 150, 171	0
3	O	211/217 (97%)	0.94	51 (24%) 0 0	66, 112, 228, 246	0
All	All	3945/4190 (94%)	0.13	288 (7%) 15 6	45, 90, 212, 247	0

The worst 5 of 288 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	N	120	PRO	13.0
2	J	150	ILE	11.9
3	O	182	SER	11.7
2	J	181	LEU	11.3
3	K	183	SER	11.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CL	B	505	1/1	0.66	0.34	104,104,104,104	0
4	CL	C	506	1/1	0.66	0.43	104,104,104,104	0
4	CL	B	503	1/1	0.69	0.53	129,129,129,129	0
4	CL	D	505	1/1	0.69	0.36	104,104,104,104	0
4	CL	E	503	1/1	0.70	0.37	136,136,136,136	0
4	CL	E	506	1/1	0.70	0.50	104,104,104,104	0
4	CL	D	503	1/1	0.70	0.57	124,124,124,124	0
4	CL	A	502	1/1	0.72	0.52	102,102,102,102	0
4	CL	A	506	1/1	0.75	0.48	106,106,106,106	0
5	K	E	504	1/1	0.78	0.35	133,133,133,133	0
6	GOL	A	504	6/6	0.85	0.38	67,93,109,110	0
6	GOL	D	501	6/6	0.86	0.29	55,74,95,99	0
4	CL	C	503	1/1	0.86	0.40	117,117,117,117	0
6	GOL	E	501	6/6	0.89	0.23	59,71,99,101	0
4	CL	C	505	1/1	0.89	0.19	107,107,107,107	0
6	GOL	B	501	6/6	0.90	0.27	60,82,99,104	0
6	GOL	C	501	6/6	0.92	0.32	61,77,106,108	0
4	CL	E	505	1/1	0.93	0.08	56,56,56,56	0
5	K	A	507	1/1	0.94	0.19	95,95,95,95	0
5	K	A	503	1/1	0.94	0.10	91,91,91,91	0
7	CA	D	502	1/1	0.95	0.13	53,53,53,53	0
5	K	C	507	1/1	0.95	0.16	92,92,92,92	0
4	CL	A	501	1/1	0.95	0.15	85,85,85,85	0
4	CL	B	504	1/1	0.96	0.09	68,68,68,68	0
5	K	C	504	1/1	0.96	0.12	92,92,92,92	0
4	CL	D	504	1/1	0.97	0.08	57,57,57,57	0
7	CA	C	502	1/1	0.98	0.17	67,67,67,67	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	CA	A	505	1/1	0.98	0.15	70,70,70,70	0
7	CA	E	502	1/1	0.98	0.11	49,49,49,49	0
7	CA	B	502	1/1	0.99	0.20	79,79,79,79	0

6.5 Other polymers [i](#)

There are no such residues in this entry.