



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 09:55 AM BST

PDB ID : 3T6C
Title : Crystal structure of an enolase from pantoea ananatis (efi target efi-501676) with bound d-gluconate and mg
Authors : Vetting, M.W.; Toro, R.; Bhosle, R.; Wasserman, S.R.; Morisco, L.L.; Hillerich, B.; Washington, E.; Scott Glenn, A.; Chowdhury, S.; Evans, B.; Hammonds, J.; Zencheck, W.D.; Imker, H.J.; Gerlt, J.A.; Almo, S.C.; Enzyme Function Initiative (EFI)
Deposited on : 2011-07-28
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	<div> <div>4%</div> <div> <div></div> <div>87%</div> <div>7%</div> <div>6%</div> </div> </div>
1	B	440	<div> <div>%</div> <div> <div></div> <div>88%</div> <div>5%</div> <div>6%</div> </div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7706 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative ManD family dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	415	Total	C	N	O	S	0	15	0
			3353	2136	570	627	20			
1	B	414	Total	C	N	O	S	0	14	0
			3332	2125	565	622	20			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	MET	-	expression tag	UNP D4GJ14
A	-21	HIS	-	expression tag	UNP D4GJ14
A	-20	HIS	-	expression tag	UNP D4GJ14
A	-19	HIS	-	expression tag	UNP D4GJ14
A	-18	HIS	-	expression tag	UNP D4GJ14
A	-17	HIS	-	expression tag	UNP D4GJ14
A	-16	HIS	-	expression tag	UNP D4GJ14
A	-15	SER	-	expression tag	UNP D4GJ14
A	-14	SER	-	expression tag	UNP D4GJ14
A	-13	GLY	-	expression tag	UNP D4GJ14
A	-12	VAL	-	expression tag	UNP D4GJ14
A	-11	ASP	-	expression tag	UNP D4GJ14
A	-10	LEU	-	expression tag	UNP D4GJ14
A	-9	GLY	-	expression tag	UNP D4GJ14
A	-8	THR	-	expression tag	UNP D4GJ14
A	-7	GLU	-	expression tag	UNP D4GJ14
A	-6	ASN	-	expression tag	UNP D4GJ14
A	-5	LEU	-	expression tag	UNP D4GJ14
A	-4	TYR	-	expression tag	UNP D4GJ14
A	-3	PHE	-	expression tag	UNP D4GJ14
A	-2	GLN	-	expression tag	UNP D4GJ14
A	-1	SER	-	expression tag	UNP D4GJ14
A	0	MET	-	expression tag	UNP D4GJ14
B	-22	MET	-	expression tag	UNP D4GJ14
B	-21	HIS	-	expression tag	UNP D4GJ14

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-20	HIS	-	expression tag	UNP D4GJ14
B	-19	HIS	-	expression tag	UNP D4GJ14
B	-18	HIS	-	expression tag	UNP D4GJ14
B	-17	HIS	-	expression tag	UNP D4GJ14
B	-16	HIS	-	expression tag	UNP D4GJ14
B	-15	SER	-	expression tag	UNP D4GJ14
B	-14	SER	-	expression tag	UNP D4GJ14
B	-13	GLY	-	expression tag	UNP D4GJ14
B	-12	VAL	-	expression tag	UNP D4GJ14
B	-11	ASP	-	expression tag	UNP D4GJ14
B	-10	LEU	-	expression tag	UNP D4GJ14
B	-9	GLY	-	expression tag	UNP D4GJ14
B	-8	THR	-	expression tag	UNP D4GJ14
B	-7	GLU	-	expression tag	UNP D4GJ14
B	-6	ASN	-	expression tag	UNP D4GJ14
B	-5	LEU	-	expression tag	UNP D4GJ14
B	-4	TYR	-	expression tag	UNP D4GJ14
B	-3	PHE	-	expression tag	UNP D4GJ14
B	-2	GLN	-	expression tag	UNP D4GJ14
B	-1	SER	-	expression tag	UNP D4GJ14
B	0	MET	-	expression tag	UNP D4GJ14

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0

- Molecule 3 is D-gluconic acid (three-letter code: GCO) (formula: C₆H₁₂O₇).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		
3	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

- Molecule 6 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total X 1 1	0	0
6	A	1	Total X 1 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	457	Total O 457 457	0	0
7	B	497	Total O 497 497	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative ManD family dehydratase



- Molecule 1: Putative ManD family dehydratase



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	130.79 Å 130.79 Å 104.68 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	21.22 – 1.60 39.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	92.5 (21.22-1.60) 91.4 (39.00-1.60)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.53 (at 1.60 Å)	Xtriage
Refinement program	PHENIX 1.7 _650	Depositor
R, R_{free}	0.139 , 0.168 0.143 , 0.171	Depositor DCC
R_{free} test set	1975 reflections (1.67%)	wwPDB-VP
Wilson B-factor (Å ²)	11.4	Xtriage
Anisotropy	0.064	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 55.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7706	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UNX, MG, EDO, GCO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	0/3435	0.86	1/4669 (0.0%)
1	B	0.76	0/3413	0.87	2/4640 (0.0%)
All	All	0.77	0/6848	0.86	3/9309 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	1
All	All	0	4

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	82	ARG	NE-CZ-NH2	-6.60	117.00	120.30
1	A	236	MET	CG-SD-CE	-5.89	90.78	100.20
1	B	44	ARG	NE-CZ-NH1	5.69	123.15	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	44	ARG	Mainchain
1	A	67	ARG	Mainchain
1	A	80	TYR	Mainchain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	80	TYR	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3353	0	3302	19	0
1	B	3332	0	3289	17	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	13	0	10	0	0
3	B	13	0	10	0	0
4	A	1	0	0	0	0
5	A	16	0	24	1	0
5	B	20	0	30	1	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	A	457	0	0	4	0
7	B	497	0	0	4	0
All	All	7706	0	6665	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205:ARG:HG3	7:A:723:HOH:O	1.53	1.06
1:B:143:LYS:HE3	7:B:580:HOH:O	1.80	0.82
1:B:362[B]:GLU:HG3	7:B:724:HOH:O	1.88	0.73
1:A:255:GLU:OE2	5:A:701:EDO:H22	1.94	0.66
1:A:362[A]:GLU:HG2	1:A:391:LEU:HD23	1.77	0.66
1:B:255:GLU:OE2	5:B:707:EDO:H22	2.02	0.59
1:A:180:GLN:HG2	1:A:180:GLN:O	2.07	0.55
1:A:128:THR:HG21	1:A:140:ALA:HA	1.91	0.53
1:A:3:ASN:HB3	7:A:775:HOH:O	2.10	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:122[B]:ILE:HD11	1:B:343:LEU:HG	1.92	0.51
1:B:391[B]:LEU:HD12	1:B:394:LYS:HD2	1.94	0.50
1:A:132:ASP:OD2	1:A:177:LYS:CE	2.60	0.49
1:A:176:ALA:CB	1:A:179:ILE:HD12	2.43	0.49
1:B:128:THR:CG2	1:B:143:LYS:HG3	2.42	0.49
1:B:152:ARG:HD3	7:B:697:HOH:O	2.14	0.47
1:A:179:ILE:HG23	1:A:198:ALA:HB1	1.96	0.47
1:B:129:ASP:HB2	7:B:822:HOH:O	2.14	0.46
1:A:164:ASP:HA	1:A:172:ARG:HG3	1.98	0.46
1:B:4:LEU:HD12	1:B:29:ASN:OD1	2.15	0.46
1:B:327:PRO:HG2	1:B:329:ASP:OD1	2.17	0.45
1:A:183[A]:ARG:NH1	7:A:697:HOH:O	2.49	0.45
1:A:353:TYR:CZ	1:A:370:VAL:HG21	2.52	0.44
1:A:132:ASP:OD2	1:A:177:LYS:HE2	2.17	0.44
1:B:194:PHE:O	1:B:410:PRO:HD3	2.19	0.43
1:A:54[A]:GLU:HG2	1:B:46:TYR:CD2	2.53	0.43
1:A:327:PRO:HG2	1:A:329:ASP:OD1	2.20	0.42
1:B:128:THR:HG21	1:B:140:ALA:HA	2.02	0.42
1:B:343:LEU:HD11	1:B:377[B]:VAL:HG22	2.02	0.41
1:B:203:ILE:HA	1:B:203:ILE:HD13	1.88	0.41
1:A:174:VAL:HG22	7:A:761:HOH:O	2.19	0.41
1:A:299[B]:ILE:HD13	1:A:341:LEU:CD1	2.51	0.41
1:A:179:ILE:O	1:A:181:PRO:HD3	2.21	0.41
1:B:11:THR:HG23	1:B:53:ASP:OD1	2.21	0.41
1:A:122[A]:ILE:HD12	1:A:342:ASP:HB3	2.03	0.40
1:B:327:PRO:HA	1:B:354:THR:OG1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	428/440 (97%)	419 (98%)	7 (2%)	2 (0%)	29	11
1	B	426/440 (97%)	417 (98%)	7 (2%)	2 (0%)	29	11
All	All	854/880 (97%)	836 (98%)	14 (2%)	4 (0%)	29	11

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	80	TYR
1	B	80	TYR
1	A	327	PRO
1	B	327	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	357/365 (98%)	354 (99%)	3 (1%)	81	70
1	B	355/365 (97%)	353 (99%)	2 (1%)	86	77
All	All	712/730 (98%)	707 (99%)	5 (1%)	84	73

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	PHE
1	A	169	ILE
1	A	247	PHE
1	B	41	PHE
1	B	152	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 2 are unknown and 3 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	B	704	-	3,3,3	0.48	0	2,2,2	0.17	0
3	GCO	A	601	2	9,12,12	1.24	1 (11%)	12,16,16	1.71	1 (8%)
3	GCO	B	601	2	9,12,12	1.69	3 (33%)	12,16,16	2.20	5 (41%)
5	EDO	B	708	-	3,3,3	1.16	0	2,2,2	0.56	0
5	EDO	A	706	-	3,3,3	0.38	0	2,2,2	0.09	0
5	EDO	A	703	-	3,3,3	0.61	0	2,2,2	0.39	0
5	EDO	B	709	-	3,3,3	0.56	0	2,2,2	0.16	0
5	EDO	B	702	-	3,3,3	0.93	0	2,2,2	0.19	0
5	EDO	A	705	-	3,3,3	1.12	0	2,2,2	0.74	0
5	EDO	B	707	-	3,3,3	0.55	0	2,2,2	0.71	0
5	EDO	A	701	-	3,3,3	0.70	0	2,2,2	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	704	-	-	0/1/1/1	-
3	GCO	A	601	2	-	4/14/18/18	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GCO	B	601	2	-	4/14/18/18	-
5	EDO	B	708	-	-	0/1/1/1	-
5	EDO	A	706	-	-	0/1/1/1	-
5	EDO	A	703	-	-	1/1/1/1	-
5	EDO	B	709	-	-	0/1/1/1	-
5	EDO	B	702	-	-	0/1/1/1	-
5	EDO	A	705	-	-	0/1/1/1	-
5	EDO	B	707	-	-	0/1/1/1	-
5	EDO	A	701	-	-	0/1/1/1	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	601	GCO	C3-C2	-3.56	1.49	1.53
3	A	601	GCO	O5-C5	-2.35	1.38	1.43
3	B	601	GCO	O5-C5	-2.27	1.38	1.43
3	B	601	GCO	O3-C3	-2.04	1.38	1.43

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	601	GCO	O2-C2-C3	4.77	118.18	107.39
3	A	601	GCO	O2-C2-C3	4.70	118.02	107.39
3	B	601	GCO	C4-C3-C2	-3.07	108.16	113.58
3	B	601	GCO	O3-C3-C2	-2.95	102.57	108.95
3	B	601	GCO	O6-C6-C5	2.08	115.60	111.07
3	B	601	GCO	C6-C5-C4	2.05	116.86	112.41

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	601	GCO	C2-C3-C4-O4
3	A	601	GCO	O3-C3-C4-C5
3	B	601	GCO	C2-C3-C4-C5
3	B	601	GCO	C2-C3-C4-O4
3	B	601	GCO	O3-C3-C4-C5
3	A	601	GCO	O3-C3-C4-O4
3	B	601	GCO	O3-C3-C4-O4
3	A	601	GCO	C2-C3-C4-C5
5	A	703	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	707	EDO	1	0
5	A	701	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	415/440 (94%)	-0.58	17 (4%) 37 34	8, 12, 31, 61	0
1	B	414/440 (94%)	-0.62	3 (0%) 87 87	8, 12, 24, 42	0
All	All	829/880 (94%)	-0.60	20 (2%) 59 56	8, 12, 25, 61	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	169	ILE	5.4
1	A	179	ILE	4.4
1	A	174	VAL	4.2
1	A	176	ALA	4.0
1	A	177	LYS	3.9
1	A	166	LEU	3.7
1	A	171	ASN	3.7
1	A	168	LEU	3.6
1	A	178	ASN	3.5
1	A	173[A]	MET	3.4
1	A	172	ARG	3.4
1	A	180	GLN	3.3
1	A	164	ASP	3.0
1	A	170	ALA	2.7
1	A	162	GLY	2.6
1	A	175	LYS	2.6
1	B	31	PRO	2.5
1	B	175	LYS	2.2
1	A	161	ALA	2.2
1	B	177	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	EDO	A	705	4/4	0.89	0.12	17,20,24,25	0
5	EDO	A	703	4/4	0.92	0.19	26,38,40,47	0
5	EDO	A	701	4/4	0.92	0.11	19,27,31,33	0
5	EDO	B	702	4/4	0.93	0.09	19,21,26,30	0
5	EDO	B	708	4/4	0.93	0.10	14,20,21,25	0
5	EDO	B	707	4/4	0.94	0.11	22,24,28,34	0
5	EDO	B	704	4/4	0.95	0.10	28,28,29,32	0
3	GCO	B	601	13/13	0.96	0.07	10,14,26,27	0
5	EDO	A	706	4/4	0.97	0.07	23,25,25,37	0
5	EDO	B	709	4/4	0.97	0.07	19,19,22,26	0
3	GCO	A	601	13/13	0.98	0.05	9,12,15,15	0
6	UNX	A	418	1/1	1.00	0.46	2,2,2,2	1
4	CL	A	901	1/1	1.00	0.03	10,10,10,10	0
6	UNX	B	418	1/1	1.00	0.43	2,2,2,2	1
2	MG	B	501	1/1	1.00	0.06	8,8,8,8	0
2	MG	A	501	1/1	1.00	0.04	8,8,8,8	0

6.5 Other polymers [i](#)

There are no such residues in this entry.