



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 11:30 am BST

PDB ID : 3T8V
Title : A bestatin-based chemical biology strategy reveals distinct roles for malaria M1- and M17-family aminopeptidases
Authors : McGowan, S.; Klemba, M.; Greebaum, D.C.
Deposited on : 2011-08-01
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	895	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> % 96% .. </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	BTJ	A	1092	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8284 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called M1 family aminopeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	889	Total	C	N	O	S	0	6	0
			7233	4658	1176	1371	28			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	213	GLN	ASN	ENGINEERED MUTATION	UNP O96935
A	223	GLN	ASN	ENGINEERED MUTATION	UNP O96935
A	378	PRO	HIS	ENGINEERED MUTATION	UNP O96935
A	501	GLN	ASN	ENGINEERED MUTATION	UNP O96935
A	572	ALA	GLU	ENGINEERED MUTATION	UNP O96935
A	745	GLN	ASN	ENGINEERED MUTATION	UNP O96935
A	795	GLN	ASN	ENGINEERED MUTATION	UNP O96935
A	1069	GLN	ASN	ENGINEERED MUTATION	UNP O96935
A	1085	HIS	-	EXPRESSION TAG	UNP O96935
A	1086	HIS	-	EXPRESSION TAG	UNP O96935
A	1087	HIS	-	EXPRESSION TAG	UNP O96935
A	1088	HIS	-	EXPRESSION TAG	UNP O96935
A	1089	HIS	-	EXPRESSION TAG	UNP O96935
A	1090	HIS	-	EXPRESSION TAG	UNP O96935


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Mg	0	0
			3	3		

- Molecule 4 is N-[(2-{2-[(N-{(2S,3R)-3-amino-4-[4-(benzyloxy)phenyl]-2-hydroxybutanoyl}-L-alanyl)amino]ethoxy}ethoxy)acetyl]-4-benzoyl-L-phenylalanyl-N 6 -hex-5-ynoyllysineamide (three-letter code: BTJ) (formula: C₅₄H₆₇N₇O₁₁).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			40	29	4	7		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1007	Total	O	0	0
			1007	1007		

i

- Molecule 1: M1 family aminopeptidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	75.54Å 108.82Å 118.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.04 – 1.80 37.77 – 1.70	Depositor EDS
% Data completeness (in resolution range)	(Not available) (40.04-1.80) 98.4 (37.77-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 1.70Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.160 , 0.198 0.166 , 0.203	Depositor DCC
R_{free} test set	5254 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	16.7	Xtriage
Anisotropy	0.262	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8284	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG, BTJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/7415	0.60	0/10041

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7233	0	7092	20	0
2	A	1	0	0	0	0
3	A	3	0	0	0	0
4	A	40	0	32	6	0
5	A	1007	0	0	8	0
All	All	8284	0	7124	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1041:GLU:CD	5:A:1234:HOH:O	2.05	0.95
1:A:1041:GLU:CG	5:A:1234:HOH:O	2.21	0.87
1:A:1041:GLU:HG2	5:A:1234:HOH:O	1.84	0.77
1:A:1041:GLU:OE2	5:A:1234:HOH:O	1.98	0.76
1:A:969:ARG:HG3	5:A:1216:HOH:O	1.92	0.70
1:A:358:ASP:OD2	5:A:1270:HOH:O	2.09	0.70
1:A:459:VAL:HB	4:A:1092:BTJ:H5A	1.89	0.55
1:A:568:TYR:HB3	1:A:574:PHE:CE1	2.41	0.55
4:A:1092:BTJ:C52	4:A:1092:BTJ:H8A	2.38	0.52
1:A:571:MET:O	1:A:574:PHE:HB2	2.11	0.51
1:A:512:TRP:HB3	1:A:571:MET:SD	2.50	0.51
1:A:656:PHE:O	1:A:745:GLN:HG3	2.12	0.50
1:A:480:LYS:HG3	5:A:1274:HOH:O	2.13	0.49
4:A:1092:BTJ:H7A	5:A:1218:HOH:O	2.13	0.49
1:A:572:ALA:O	4:A:1092:BTJ:H51	2.15	0.47
1:A:332:ARG:HG3	1:A:335:MET:HG2	1.97	0.47
1:A:550:VAL:HG12	1:A:577:THR:HG21	1.97	0.47
1:A:811:LEU:HB3	1:A:867:MET:SD	2.54	0.46
1:A:462:MET:SD	4:A:1092:BTJ:H42	2.57	0.44
1:A:271:TYR:CZ	1:A:273:ASN:HA	2.54	0.43
1:A:538:THR:HG21	1:A:819:GLY:HA3	2.02	0.41
1:A:458:ASN:HB3	4:A:1092:BTJ:H54	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	893/895 (100%)	877 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	797/826 (96%)	786 (99%)	11 (1%)	67 59

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	370	HIS
1	A	436	ASP
1	A	555	ASP
1	A	574	PHE
1	A	688	PHE
1	A	702	LYS
1	A	840	LEU
1	A	1041	GLU
1	A	1045	LEU
1	A	1052	LYS
1	A	1076	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	212	ASN
1	A	217	ASN
1	A	219	ASN
1	A	527	ASN
1	A	745	GLN
1	A	882	ASN
1	A	915	ASN
1	A	1005	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	BTJ	A	1092	2	41,41,75	3.77	18 (43%)	50,52,96	3.99	23 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BTJ	A	1092	2	-	21/40/40/77	0/2/2/4

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1092	BTJ	C40-C41	-10.86	1.41	1.54
4	A	1092	BTJ	C39-N5	-8.86	1.14	1.34
4	A	1092	BTJ	C47-C46	-8.29	1.22	1.38
4	A	1092	BTJ	O11-C40	-7.36	1.27	1.42
4	A	1092	BTJ	O10-C46	-6.19	1.23	1.37
4	A	1092	BTJ	C9-N2	6.11	1.47	1.34
4	A	1092	BTJ	O1-C3	-6.03	1.11	1.23
4	A	1092	BTJ	C55-C50	4.30	1.48	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1092	BTJ	C51-C50	4.07	1.47	1.38
4	A	1092	BTJ	C41-N6	-3.84	1.34	1.47
4	A	1092	BTJ	C44-C43	-3.83	1.30	1.38
4	A	1092	BTJ	C10-N2	3.15	1.52	1.47
4	A	1092	BTJ	C42-C41	-3.12	1.49	1.53
4	A	1092	BTJ	C48-C43	3.12	1.45	1.38
4	A	1092	BTJ	C40-C39	-3.11	1.46	1.52
4	A	1092	BTJ	C54-C55	-2.74	1.33	1.38
4	A	1092	BTJ	C45-C46	-2.49	1.33	1.38
4	A	1092	BTJ	C48-C47	-2.12	1.34	1.38

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1092	BTJ	C44-C45-C46	10.73	132.86	119.73
4	A	1092	BTJ	C25-C10-N2	10.24	130.41	109.79
4	A	1092	BTJ	C40-C39-N5	-8.96	103.91	116.25
4	A	1092	BTJ	O9-C39-C40	8.08	134.57	120.00
4	A	1092	BTJ	C47-C48-C43	7.31	131.08	121.03
4	A	1092	BTJ	C49-O10-C46	6.87	134.62	117.65
4	A	1092	BTJ	C8-O3-C7	6.45	129.93	113.13
4	A	1092	BTJ	O3-C8-C9	6.24	125.78	112.38
4	A	1092	BTJ	C48-C43-C44	-5.51	109.51	118.17
4	A	1092	BTJ	C10-N2-C9	4.47	129.14	123.00
4	A	1092	BTJ	C3-C2-N5	4.35	122.37	111.60
4	A	1092	BTJ	O11-C40-C39	4.29	119.89	110.63
4	A	1092	BTJ	C45-C44-C43	-4.22	115.22	121.03
4	A	1092	BTJ	C4-N1-C3	4.09	129.89	122.59
4	A	1092	BTJ	C11-C10-N2	4.05	117.95	109.79
4	A	1092	BTJ	C40-C41-N6	3.98	121.50	111.38
4	A	1092	BTJ	C42-C43-C48	3.71	128.27	120.91
4	A	1092	BTJ	C25-C10-C11	-3.51	103.77	111.58
4	A	1092	BTJ	C47-C46-C45	-3.33	115.05	120.18
4	A	1092	BTJ	C48-C47-C46	-2.96	116.11	119.73
4	A	1092	BTJ	C1-C2-N5	2.94	115.89	110.38
4	A	1092	BTJ	C5-C4-N1	-2.77	105.28	111.83
4	A	1092	BTJ	C2-N5-C39	-2.58	115.70	121.29

There are no chirality outliers.

All (21) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1092	BTJ	O4-C9-N2-C10
4	A	1092	BTJ	C8-C9-N2-C10
4	A	1092	BTJ	N5-C39-C40-O11
4	A	1092	BTJ	N5-C39-C40-C41
4	A	1092	BTJ	O9-C39-C40-O11
4	A	1092	BTJ	O9-C39-C40-C41
4	A	1092	BTJ	O11-C40-C41-N6
4	A	1092	BTJ	O11-C40-C41-C42
4	A	1092	BTJ	C39-C40-C41-N6
4	A	1092	BTJ	C39-C40-C41-C42
4	A	1092	BTJ	C41-C42-C43-C48
4	A	1092	BTJ	N1-C4-C5-O2
4	A	1092	BTJ	C41-C42-C43-C44
4	A	1092	BTJ	C9-C8-O3-C7
4	A	1092	BTJ	O3-C8-C9-N2
4	A	1092	BTJ	O3-C8-C9-O4
4	A	1092	BTJ	C40-C41-C42-C43
4	A	1092	BTJ	C6-C7-O3-C8
4	A	1092	BTJ	C4-C5-O2-C6
4	A	1092	BTJ	C7-C6-O2-C5
4	A	1092	BTJ	C1-C2-C3-O1

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1092	BTJ	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	889/895 (99%)	-0.39	10 (1%) 80 78	9, 17, 34, 54	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	196	PRO	5.4
1	A	200	TYR	3.2
1	A	571	MET	3.1
1	A	634	ASP	2.7
1	A	572	ALA	2.3
1	A	366	PRO	2.2
1	A	635	ASN	2.2
1	A	574	PHE	2.1
1	A	636	SER	2.1
1	A	543[A]	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	BTJ	A	1092	40/72	0.84	0.19	12,35,91,131	0
3	MG	A	1091	1/1	0.91	0.11	39,39,39,39	0
3	MG	A	3	1/1	0.97	0.20	34,34,34,34	0
3	MG	A	2	1/1	0.99	0.11	23,23,23,23	0
2	ZN	A	1	1/1	1.00	0.07	13,13,13,13	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.