



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 6, 2020 – 03:20 PM BST

PDB ID : 6T8J  
Title : Crystal structure of formate dehydrogenase FDH2 enzyme from *Granulicella mallensis* MP5ACTX8 in complex with NAD and azide.  
Authors : Robescu, M.S.; Rubini, R.; Filippini, F.; Bergantino, B.; Cendron, L.  
Deposited on : 2019-10-24  
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	386	<div> <div>0%</div> <div>94%</div> <div>5%</div> <div>• •</div> </div>
1	B	386	<div> <div>0%</div> <div>92%</div> <div>5%</div> <div>• • •</div> </div>
1	C	386	<div> <div>3%</div> <div>91%</div> <div>5%</div> <div>•</div> </div>
1	D	386	<div> <div>3%</div> <div>89%</div> <div>7%</div> <div>•</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 12434 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Formate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	374	Total	C	N	O	S	0	2	0
			2927	1868	500	547	12			
1	B	373	Total	C	N	O	S	0	3	0
			2924	1866	500	546	12			
1	C	374	Total	C	N	O	S	0	0	0
			2911	1858	498	543	12			
1	D	373	Total	C	N	O	S	0	0	0
			2906	1855	497	542	12			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	184	Total	O	0	0
			184	184		
2	B	193	Total	O	0	0
			193	193		
2	C	197	Total	O	0	0
			197	197		
2	D	192	Total	O	0	0
			192	192		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Formate dehydrogenase





- Molecule 1: Formate dehydrogenase



- Molecule 1: Formate dehydrogenase



- Molecule 1: Formate dehydrogenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	99.36Å 112.49Å 147.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	62.10 – 1.97 48.95 – 1.97	Depositor EDS
% Data completeness (in resolution range)	97.8 (62.10-1.97) 99.8 (48.95-1.97)	Depositor EDS
$R_{merge}$	0.20	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.95 (at 1.97Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.189 , 0.235 0.171 , 0.206	Depositor DCC
$R_{free}$ test set	5824 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.4	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 37.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12434	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.54	0/2999	0.81	0/4081
1	B	0.54	0/2996	0.81	0/4076
1	C	0.51	0/2983	0.81	2/4059 (0.0%)
1	D	0.53	0/2978	0.81	0/4052
All	All	0.53	0/11956	0.81	2/16268 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	296	ARG	CG-CD-NE	-7.70	95.62	111.80
1	C	253	THR	CB-CA-C	-5.47	96.83	111.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	125[A]	SER	Mainchain
1	B	125[B]	SER	Mainchain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2927	0	2913	7	0
1	B	2924	0	2913	15	0
1	C	2911	0	2900	14	0
1	D	2906	0	2895	13	0
2	A	184	0	0	2	0
2	B	193	0	0	5	0
2	C	197	0	0	4	0
2	D	192	0	0	1	0
All	All	12434	0	11621	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:123[B]:ILE:HD11	1:B:146:SER:N	2.00	0.77
1:B:123[A]:ILE:CD1	2:B:562:HOH:O	2.36	0.73
1:C:259:HIS:HB2	1:C:260:PRO:HD2	1.75	0.68
1:B:120:THR:HG21	1:B:124[A]:GLY:O	1.96	0.66
1:C:296:ARG:NH1	2:C:401:HOH:O	2.29	0.66
1:A:314:GLN:NE2	2:A:401:HOH:O	2.30	0.65
1:D:373:LYS:HG2	1:D:374:LEU:N	2.12	0.63
1:C:124:GLY:N	1:C:142:GLU:OE2	2.26	0.62
1:C:259:HIS:HB2	1:C:260:PRO:CD	2.32	0.59
1:B:123[B]:ILE:HD11	1:B:146:SER:H	1.66	0.59
1:A:134:ILE:HD13	1:A:374:LEU:HD11	1.87	0.57
1:C:79:GLY:O	1:C:85:GLU:HG3	2.07	0.55
1:C:123:ILE:HD13	2:C:501:HOH:O	2.06	0.54
1:D:140:VAL:HB	1:D:369:VAL:HG13	1.90	0.53
1:D:124:GLY:N	1:D:142:GLU:OE2	2.40	0.53
1:B:123[A]:ILE:HD12	2:B:422:HOH:O	2.09	0.52
1:C:155:VAL:HG11	1:C:253:THR:HG21	1.92	0.51
1:D:81:ASP:O	1:D:86:ARG:NH1	2.44	0.51
1:A:85:GLU:OE2	1:A:111:LYS:HE2	2.11	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:107:GLU:OE2	1:B:107:GLU:N	2.42	0.49
1:D:2:ALA:N	2:D:403:HOH:O	2.46	0.49
1:B:372:GLY:C	1:B:374:LEU:H	2.17	0.48
1:A:2:ALA:N	1:A:69:HIS:HD1	2.12	0.47
1:D:4:ILE:N	1:D:4:ILE:HD12	2.30	0.47
1:C:372:GLY:HA2	2:C:449:HOH:O	2.14	0.47
1:B:123[A]:ILE:HD11	2:B:562:HOH:O	2.10	0.47
1:B:123[B]:ILE:HG21	2:B:562:HOH:O	2.15	0.47
1:D:259:HIS:HD2	1:D:261:GLY:H	1.63	0.46
1:C:281:VAL:HA	1:C:307:ALA:O	2.17	0.45
1:D:7:VAL:HA	1:D:74:THR:O	2.17	0.45
1:A:124:GLY:HA3	2:A:541:HOH:O	2.17	0.44
1:B:123[A]:ILE:HG13	2:B:562:HOH:O	2.16	0.44
1:D:94:VAL:HG23	1:D:115:LEU:HD13	1.99	0.44
1:D:196:GLY:HA2	1:D:219:HIS:O	2.17	0.44
1:D:6:CYS:O	1:D:73:VAL:HA	2.18	0.43
1:D:2:ALA:N	1:D:69:HIS:HD1	2.16	0.43
1:B:123[B]:ILE:HD11	1:B:145:TYR:C	2.38	0.43
1:B:184:VAL:HG11	1:C:337:THR:HG23	2.00	0.43
1:B:6:CYS:O	1:B:73:VAL:HA	2.19	0.42
1:C:309:ASP:O	1:C:333:HIS:HA	2.20	0.42
1:D:16:TYR:CD1	1:D:17:PRO:HD2	2.54	0.42
1:C:124:GLY:C	2:C:451:HOH:O	2.57	0.41
1:C:16:TYR:CD1	1:C:55:SER:HB2	2.55	0.41
1:B:350:ARG:O	1:B:354:GLU:HG3	2.21	0.41
1:A:373:LYS:HE2	1:B:367:LEU:O	2.21	0.40
1:A:113:LYS:HA	1:A:113:LYS:HD2	1.80	0.40
1:C:306:TYR:O	1:C:328:HIS:HA	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/386 (97%)	361 (96%)	13 (4%)	0	100	100
1	B	374/386 (97%)	364 (97%)	9 (2%)	1 (0%)	41	29
1	C	372/386 (96%)	361 (97%)	11 (3%)	0	100	100
1	D	371/386 (96%)	358 (96%)	13 (4%)	0	100	100
All	All	1491/1544 (97%)	1444 (97%)	46 (3%)	1 (0%)	51	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	199	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	315/322 (98%)	311 (99%)	4 (1%)	69	64
1	B	315/322 (98%)	308 (98%)	7 (2%)	52	46
1	C	313/322 (97%)	309 (99%)	4 (1%)	69	64
1	D	313/322 (97%)	308 (98%)	5 (2%)	62	56
All	All	1256/1288 (98%)	1236 (98%)	20 (2%)	65	56

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	ASP
1	A	113	LYS
1	A	145	TYR
1	A	285	ARG
1	B	123[A]	ILE
1	B	123[B]	ILE
1	B	126	ASP
1	B	134	ILE
1	B	145	TYR
1	B	369	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	374	LEU
1	C	131	GLU
1	C	145	TYR
1	C	253	THR
1	C	373	LYS
1	D	27	LYS
1	D	107	GLU
1	D	113	LYS
1	D	145	TYR
1	D	253	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	259	HIS
1	D	259	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	374/386 (96%)	-0.05	3 (0%) 86 87	12, 18, 37, 70	0
1	B	373/386 (96%)	0.06	5 (1%) 77 78	13, 20, 37, 61	0
1	C	374/386 (96%)	0.12	13 (3%) 44 46	14, 21, 38, 64	0
1	D	373/386 (96%)	0.17	13 (3%) 44 46	13, 20, 39, 62	0
All	All	1494/1544 (96%)	0.07	34 (2%) 60 62	12, 20, 38, 70	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	371	GLY	5.3
1	B	123[A]	ILE	5.1
1	B	124[A]	GLY	4.7
1	A	375	ALA	4.1
1	C	260	PRO	3.7
1	D	124	GLY	3.7
1	C	375	ALA	3.6
1	C	221	PHE	3.2
1	C	285	ARG	3.1
1	A	126	ASP	2.7
1	B	374	LEU	2.7
1	C	223	GLN	2.7
1	D	373	LYS	2.6
1	D	80	GLU	2.5
1	D	261	GLY	2.5
1	C	228	GLU	2.5
1	D	224	HIS	2.5
1	D	125	SER	2.4
1	C	374	LEU	2.4
1	D	259	HIS	2.4
1	C	227	PRO	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	238	TYR	2.4
1	C	261	GLY	2.4
1	D	372	GLY	2.4
1	D	79	GLY	2.3
1	D	232	ASN	2.2
1	C	224	HIS	2.2
1	C	222	ASP	2.2
1	B	373	LYS	2.1
1	D	260	PRO	2.1
1	B	155	VAL	2.1
1	D	163	VAL	2.1
1	C	259	HIS	2.0
1	A	80	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.