



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 07:31 am BST

PDB ID : 6TBY  
Title : Phycocyanobilin-adducted PAS-GAF bidomain of Sorghum bicolor phyB  
Authors : Nagano, S.; Guan, K.; Shenkutie, S.M.; Hughes, J.E.  
Deposited on : 2019-11-04  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	353	<div> <div>8%</div> <div>79%</div> <div>12%</div> <div>9%</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2699 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phytochrome.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	320	Total	C	N	O	S	0	1	0
			2483	1580	442	444	17			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
AAA	99	MET	-	initiating methionine	UNP Q6S527
AAA	100	HIS	-	expression tag	UNP Q6S527
AAA	101	HIS	-	expression tag	UNP Q6S527
AAA	102	HIS	-	expression tag	UNP Q6S527
AAA	103	HIS	-	expression tag	UNP Q6S527
AAA	104	HIS	-	expression tag	UNP Q6S527
AAA	105	HIS	-	expression tag	UNP Q6S527
AAA	106	GLU	-	expression tag	UNP Q6S527
AAA	107	ASN	-	expression tag	UNP Q6S527
AAA	108	LEU	-	expression tag	UNP Q6S527
AAA	109	TYR	-	expression tag	UNP Q6S527
AAA	110	PHE	-	expression tag	UNP Q6S527
AAA	111	GLN	-	expression tag	UNP Q6S527
AAA	112	GLY	-	expression tag	UNP Q6S527

- Molecule 2 is PHYCOCYANOBILIN (three-letter code: CYC) (formula:  $C_{33}H_{40}N_4O_6$ ) (labeled as "Ligand of Interest" by author).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	AAA	1	Total	C	N	O	0	0
			43	33	4	6		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $\text{C}_3\text{H}_8\text{O}_3$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	AAA	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula:  $C_2H_6OS$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	AAA	1	Total	C	O	S	0	0
			4	2	1	1		
4	AAA	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	AAA	1	Total	C	O	0	0
			7	4	3		
5	AAA	1	Total	C	O	0	0
			7	4	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	AAA	145	Total 145	O 145	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phytochrome



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	134.74 Å   134.74 Å   46.54 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	44.10 – 1.80 44.10 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (44.10-1.80) 99.3 (44.10-1.80)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.97 (at 1.79 Å)	Xtriage
Refinement program	REFMAC 5.8.0257	Depositor
R, $R_{free}$	0.179 , 0.212 0.187 , 0.217	Depositor DCC
$R_{free}$ test set	2198 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.9	Xtriage
Anisotropy	0.018	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 45.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.030 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2699	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CYC, GOL, PEG, BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	AAA	0.74	0/2541	0.91	3/3450 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	AAA	348	ARG	CG-CD-NE	-6.54	98.06	111.80
1	AAA	306	ARG	CG-CD-NE	-6.16	98.87	111.80
1	AAA	235	ARG	CB-CA-C	5.00	120.41	110.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2483	0	2487	30	1
2	AAA	43	0	37	1	0
3	AAA	6	0	8	1	0
4	AAA	8	0	10	0	0
5	AAA	14	0	20	1	0
6	AAA	145	0	0	3	0
All	All	2699	0	2562	32	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:282:LEU:HD23	1:AAA:430:LEU:HD22	1.55	0.88
1:AAA:210:ARG:NH1	6:AAA:601:HOH:O	1.90	0.81
1:AAA:333:GLN:HG3	1:AAA:334:ASN:ND2	1.99	0.78
1:AAA:282:LEU:CD2	1:AAA:430:LEU:HD22	2.22	0.68
1:AAA:295:GLU:CD	1:AAA:295:GLU:H	1.97	0.67
1:AAA:412:LEU:HB3	3:AAA:501:GOL:H2	1.81	0.61
1:AAA:388:MET:SD	1:AAA:434:CYS:HB3	2.40	0.61
1:AAA:282:LEU:HD23	1:AAA:430:LEU:CD2	2.29	0.57
1:AAA:291:TYR:CZ	1:AAA:299:GLY:HA3	2.40	0.57
1:AAA:119:ILE:HG21	1:AAA:330:LEU:HD12	1.88	0.55
1:AAA:119:ILE:HG21	1:AAA:330:LEU:CD1	2.37	0.54
1:AAA:108:LEU:HD12	1:AAA:328:ARG:NH2	2.22	0.54
1:AAA:355:MET:HE3	6:AAA:692:HOH:O	2.11	0.49
1:AAA:250:GLN:NE2	1:AAA:436:PHE:CE2	2.80	0.49
2:AAA:500:CYC:NB	2:AAA:500:CYC:HMA1	2.29	0.48
1:AAA:266:GLY:N	1:AAA:448:GLU:OE1	2.44	0.47
1:AAA:282:LEU:HD22	1:AAA:434:CYS:SG	2.56	0.46
1:AAA:260:LEU:O	1:AAA:444:GLN:NE2	2.44	0.46
1:AAA:282:LEU:CD2	1:AAA:430:LEU:CD2	2.90	0.46
5:AAA:505:PEG:H31	5:AAA:505:PEG:H12	1.50	0.46
1:AAA:333:GLN:HG3	1:AAA:334:ASN:HD21	1.80	0.45
1:AAA:249:SER:HB2	1:AAA:430:LEU:HD23	1.98	0.45
1:AAA:119:ILE:CG2	1:AAA:330:LEU:CD1	2.95	0.44
1:AAA:355:MET:CE	6:AAA:692:HOH:O	2.66	0.44
1:AAA:106:GLU:OE1	1:AAA:106:GLU:N	2.52	0.43
1:AAA:426:ILE:HD12	1:AAA:430:LEU:HB3	2.01	0.42
1:AAA:152:LEU:HD12	1:AAA:152:LEU:HA	1.83	0.41
1:AAA:109:TYR:CG	1:AAA:110:PHE:N	2.88	0.41
1:AAA:110:PHE:CD1	1:AAA:110:PHE:C	2.94	0.41
1:AAA:174:LEU:HD13	1:AAA:352:ASP:HA	2.02	0.41
1:AAA:385:SER:HA	1:AAA:418:CYS:O	2.22	0.40
1:AAA:125:ILE:HG13	1:AAA:347:VAL:HB	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:250:GLN:OE1	1:AAA:250:GLN:NE2[6_555]	2.15	0.05

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	315/353 (89%)	304 (96%)	11 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	267/294 (91%)	262 (98%)	5 (2%)	57	46

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	117	SER
1	AAA	255	ARG
1	AAA	298	HIS
1	AAA	337	ARG
1	AAA	451	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	BME	AAA	503	1	3,3,3	0.36	0	1,2,2	0.30	0
5	PEG	AAA	505	-	6,6,6	0.23	0	5,5,5	0.15	0
2	CYC	AAA	500	1	36,46,46	1.18	2 (5%)	44,67,67	1.14	5 (11%)
5	PEG	AAA	504	-	6,6,6	0.21	0	5,5,5	0.06	0
3	GOL	AAA	501	-	5,5,5	0.11	0	5,5,5	0.42	0
4	BME	AAA	502	1	3,3,3	0.22	0	1,2,2	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BME	AAA	503	1	-	1/1/1/1	-
5	PEG	AAA	505	-	-	3/4/4/4	-
2	CYC	AAA	500	1	-	3/21/74/74	0/4/4/4

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PEG	AAA	504	-	-	2/4/4/4	-
3	GOL	AAA	501	-	-	4/4/4/4	-
4	BME	AAA	502	1	-	1/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	AAA	500	CYC	CHA-C1A	5.75	1.39	1.35
2	AAA	500	CYC	C1B-C2B	-2.29	1.41	1.45

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	AAA	500	CYC	C4D-CHA-C1A	3.75	133.29	128.81
2	AAA	500	CYC	CAB-C3B-C2B	2.41	131.66	127.53
2	AAA	500	CYC	CAA-CBA-CGA	2.34	116.59	112.67
2	AAA	500	CYC	CBD-CAD-C3D	-2.19	108.45	112.49
2	AAA	500	CYC	CHA-C1A-NA	-2.04	126.00	128.83

There are no chirality outliers.

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	AAA	500	CYC	C3A-C4A-CHB-C1B
2	AAA	500	CYC	ND-C1D-CHD-C4C
4	AAA	502	BME	O1-C1-C2-S2
5	AAA	505	PEG	C1-C2-O2-C3
5	AAA	504	PEG	O1-C1-C2-O2
5	AAA	505	PEG	O1-C1-C2-O2
2	AAA	500	CYC	NA-C4A-CHB-C1B
3	AAA	501	GOL	O1-C1-C2-C3
5	AAA	504	PEG	O2-C3-C4-O4
3	AAA	501	GOL	O1-C1-C2-O2
4	AAA	503	BME	O1-C1-C2-S2
5	AAA	505	PEG	C4-C3-O2-C2
3	AAA	501	GOL	C1-C2-C3-O3
3	AAA	501	GOL	O2-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	AAA	505	PEG	1	0
2	AAA	500	CYC	1	0
3	AAA	501	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	AAA	320/353 (90%)	0.14	28 (8%) 10 7	24, 37, 75, 107	2 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	451	LEU	10.0
1	AAA	450	GLN	5.5
1	AAA	266	GLY	5.0
1	AAA	106	GLU	4.6
1	AAA	449	LEU	4.5
1	AAA	447	MET	4.5
1	AAA	107	ASN	4.2
1	AAA	238	ASP	4.0
1	AAA	409	ALA	3.9
1	AAA	237	GLU	3.4
1	AAA	242	SER	3.3
1	AAA	211	VAL	3.2
1	AAA	267	ASP	3.1
1	AAA	263	LEU	3.1
1	AAA	268	ILE	3.0
1	AAA	410	MET	2.9
1	AAA	110	PHE	2.8
1	AAA	235	ARG	2.8
1	AAA	265	GLY	2.8
1	AAA	236	THR	2.7
1	AAA	264	PRO	2.6
1	AAA	394	SER	2.6
1	AAA	108	LEU	2.5
1	AAA	295	GLU	2.5
1	AAA	262	ALA	2.3
1	AAA	240	ALA	2.3
1	AAA	109	TYR	2.2

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	AAA	169	PRO	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	PEG	AAA	504	7/7	0.61	0.24	42,56,60,60	7
5	PEG	AAA	505	7/7	0.75	0.18	44,47,51,54	7
3	GOL	AAA	501	6/6	0.85	0.38	56,64,68,71	0
4	BME	AAA	502	4/4	0.94	0.12	59,60,60,68	0
2	CYC	AAA	500	43/43	0.97	0.08	24,30,36,47	0
4	BME	AAA	503	4/4	0.97	0.08	41,48,49,56	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around CYC AAA 500:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers ⓘ

There are no such residues in this entry.