



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 14, 2020 – 09:34 AM BST

PDB ID : 6TC9
Title : Crystal structure of MutM from *Neisseria meningitidis*
Authors : Silhan, J.; Landova, B.; Boura, E.
Deposited on : 2019-11-05
Resolution : 2.17 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.4.dev1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.4.dev1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6864 (2.20-2.16)
Clashscore	141614	7689 (2.20-2.16)
Ramachandran outliers	138981	7564 (2.20-2.16)
Sidechain outliers	138945	7564 (2.20-2.16)
RSRZ outliers	127900	6738 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	14	<div> <div>57%</div> <div>29%</div> <div>14%</div> </div>
1	G	14	<div> <div>64%</div> <div>21%</div> <div>14%</div> </div>
2	A	275	<div> <div>4%</div> <div>85%</div> <div>9%</div> <div>6%</div> </div>
2	C	275	<div> <div>3%</div> <div>79%</div> <div>14%</div> <div>6%</div> </div>
3	B	14	<div> <div>86%</div> <div>14%</div> </div>
3	F	14	<div> <div>14%</div> <div>64%</div> <div>36%</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5173 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA containing abasic site analogue.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	12	Total	C	N	O	P	0	0	0
			230	110	36	72	12			
1	G	12	Total	C	N	O	P	0	0	0
			230	110	36	72	12			

- Molecule 2 is a protein called Formamidopyrimidine-DNA glycosylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	259	Total	C	N	O	S	0	0	0
			2037	1296	374	357	10			
2	C	259	Total	C	N	O	S	0	0	0
			2038	1295	376	357	10			

- Molecule 3 is a DNA chain called DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	B	14	Total	C	N	O	P	0	0	0
			288	137	58	80	13			
3	F	14	Total	C	N	O	P	0	0	0
			288	137	58	80	13			

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		
4	C	1	Total	Zn	0	0
			1	1		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total 1	Mg 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	G	1	Total 1	O 1	0	0
6	A	16	Total 16	O 16	0	0
6	C	41	Total 41	O 41	0	0
6	F	1	Total 1	O 1	0	0

- Molecule 1: DNA containing abasic site analogue




- | | | | | | | | | |
|----|----|----|----|----|----|----|--|-----|
| DT | DG | T3 | C4 | C5 | A6 | N7 | | C14 |
|----|----|----|----|----|----|----|--|-----|

-
- | Position | Residue | Information Content (bits) |
|----------|-------------------|----------------------------|
| K198 | Lysine (K) | 0.05 |
| G219 | Glycine (G) | 0.35 |
| G219 | Serine (S) | 0.05 |
| G219 | Threonine (T) | 0.02 |
| G219 | Leucine (L) | 0.02 |
| G219 | Alanine (A) | 0.01 |
| G219 | Arginine (R) | 0.01 |
| G219 | Aspartate (D) | 0.01 |
| G219 | Phenylalanine (F) | 0.01 |
| G219 | Valine (V) | 0.01 |
| G219 | Asparagine (Q) | 0.01 |
| G219 | Serine (S) | 0.01 |
| G219 | Aspartate (D) | 0.01 |
| G219 | Glutamine (N) | 0.01 |
| G219 | Glutamate (E) | 0.01 |
| G219 | Lysine (K) | 0.01 |
| G219 | Serine (S) | 0.01 |
| G234 | Glutamate (E) | 0.05 |
| K275 | Lysine (K) | 0.05 |

- | Y122 | E123 | GLY | ILE | E126 | A146 | D147 | Y150 | R152 | A168 | G172 | N175 | N179 | K198 | A202 | G219 | GLY | SER | THR | LEU | ARG | ASP | PHE | VAL | ASP | SER | ASP | GLY | LNS | S233 | R352 | V258 | C273 | G274 | Z275 |
|------|------|-----|-----|------|------|------|------|------|------|------|------|------|------|------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|------|------|------|------|------|
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |


- 
- WORLD WIDE
PDB
PROTEIN DATA BANK

Chain B:  86% 14%



● Molecule 3: DNA

Chain F:  14% 64% 36%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	130.77Å 80.38Å 83.62Å 90.00° 89.95° 90.00°	Depositor
Resolution (Å)	36.22 – 2.17 36.22 – 2.18	Depositor EDS
% Data completeness (in resolution range)	98.4 (36.22-2.17) 98.5 (36.22-2.18)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 2.18Å)	Xtriage
Refinement program	PHENIX 1.15.2-3472	Depositor
R, R_{free}	0.238 , 0.281 0.238 , 0.281	Depositor DCC
R_{free} test set	2250 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	44.3	Xtriage
Anisotropy	0.633	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.479 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5173	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG, 3DR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	D	0.72	0/242	1.05	0/367
1	G	0.68	0/242	1.04	0/367
2	A	0.47	0/2079	0.64	0/2808
2	C	0.48	0/2080	0.66	2/2809 (0.1%)
3	B	0.74	0/324	0.91	0/499
3	F	0.71	0/324	0.91	0/499
All	All	0.53	0/5291	0.74	2/7349 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	33	LEU	CA-CB-CG	6.47	130.19	115.30
2	C	80	ARG	NE-CZ-NH1	-5.08	117.76	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	230	0	133	2	0
1	G	230	0	133	1	0
2	A	2037	0	2067	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	2038	0	2065	30	0
3	B	288	0	158	1	0
3	F	288	0	158	3	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	C	1	0	0	0	0
6	A	16	0	0	0	0
6	C	41	0	0	1	0
6	F	1	0	0	0	0
6	G	1	0	0	0	0
All	All	5173	0	4714	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:122:TYR:CE1	2:C:123:GLU:OE2	2.25	0.90
2:A:30:GLN:OE1	2:A:31:LEU:N	2.20	0.74
2:C:147:ASP:OD1	2:C:198:LYS:NZ	2.21	0.72
2:C:31:LEU:HD13	2:C:36:GLN:OE1	1.92	0.70
2:A:192:PRO:HB2	2:A:195:ARG:HG3	1.78	0.65
2:A:58:LYS:HE3	2:A:172:GLY:HA2	1.79	0.64
2:C:93:ASP:OD2	2:C:111:ARG:NH1	2.34	0.61
2:A:33:LEU:HA	2:A:115:LYS:HB3	1.83	0.60
2:C:18:ILE:HG21	2:C:108:MET:HE3	1.84	0.60
1:D:9:DT:O2	2:A:34:ARG:NH1	2.35	0.59
2:C:58:LYS:HE3	2:C:172:GLY:HA2	1.84	0.58
2:C:27:VAL:HB	2:C:99:ASP:HB2	1.86	0.57
2:C:104:ASP:OD2	2:C:106:THR:OG1	2.23	0.56
2:A:147:ASP:OD1	2:A:198:LYS:NZ	2.21	0.54
2:C:30:GLN:CD	2:C:31:LEU:H	2.11	0.54
2:C:18:ILE:HD13	2:C:108:MET:HE1	1.90	0.53
2:A:40:ASP:O	2:A:44:ILE:HG13	2.09	0.53
2:A:27:VAL:HB	2:A:99:ASP:HB2	1.91	0.53
2:C:40:ASP:O	2:C:44:ILE:HG13	2.09	0.53
2:A:31:LEU:HD13	2:A:36:GLN:OE1	2.10	0.51
2:C:33:LEU:O	2:C:35:TRP:N	2.42	0.50
2:C:83:THR:HG22	6:C:403:HOH:O	2.13	0.49
2:A:123:GLU:N	2:A:123:GLU:OE2	2.46	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:123:GLU:H	2:C:123:GLU:CD	2.14	0.49
2:C:80:ARG:NH1	2:C:82:PHE:CZ	2.82	0.47
2:A:18:ILE:HG21	2:A:108:MET:HE3	1.96	0.47
2:C:152:ARG:NH1	2:C:168:ALA:O	2.47	0.46
1:G:5:DC:H2'	1:G:6:DA:C8	2.50	0.46
2:C:122:TYR:CD1	2:C:123:GLU:OE2	2.68	0.45
2:C:175:ASN:O	2:C:179:ASN:ND2	2.47	0.45
2:A:104:ASP:OD2	2:A:106:THR:OG1	2.34	0.45
2:C:48:ARG:HD3	2:C:48:ARG:HA	1.83	0.44
3:F:3:DG:H2''	3:F:4:DT:O5'	2.18	0.44
2:C:252:ARG:HD3	2:C:273:CYS:SG	2.58	0.43
2:C:2:PRO:HB3	2:C:6:GLU:HB3	2.00	0.43
1:D:5:DC:H2'	1:D:6:DA:C8	2.54	0.43
2:C:104:ASP:CG	2:C:106:THR:HG1	2.23	0.42
2:C:146:ALA:HB2	2:C:202:ALA:HA	2.00	0.42
3:F:2:DG:C6	3:F:3:DG:C6	3.07	0.42
2:C:80:ARG:NH1	2:C:82:PHE:HZ	2.17	0.42
2:A:152:ARG:NH1	2:A:168:ALA:O	2.53	0.42
3:B:2:DG:C6	3:B:3:DG:C6	3.07	0.42
2:A:104:ASP:CG	2:A:106:THR:HG1	2.22	0.42
2:C:18:ILE:HD13	2:C:108:MET:CE	2.49	0.42
2:C:60:LEU:HD12	2:C:79:LEU:HD12	2.01	0.41
3:F:12:DG:H2''	3:F:13:DA:C8	2.55	0.41
2:C:71:ILE:HD13	2:C:79:LEU:HD11	2.01	0.41
2:C:112:ASP:CG	2:C:115:LYS:HA	2.40	0.41
2:C:30:GLN:NE2	2:C:31:LEU:H	2.17	0.41
2:C:112:ASP:OD1	2:C:115:LYS:HA	2.22	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	253/275 (92%)	244 (96%)	9 (4%)	0	100	100
2	C	253/275 (92%)	239 (94%)	14 (6%)	0	100	100
All	All	506/550 (92%)	483 (96%)	23 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	215/232 (93%)	214 (100%)	1 (0%)	88	94
2	C	215/232 (93%)	210 (98%)	5 (2%)	50	60
All	All	430/464 (93%)	424 (99%)	6 (1%)	67	78

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	94	ARG
2	C	34	ARG
2	C	94	ARG
2	C	103	SER
2	C	115	LYS
2	C	123	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	C	264	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	3DR	D	7	1	8,11,12	0.51	0	9,14,17	1.04	1 (11%)
1	3DR	G	7	1	8,11,12	0.69	0	9,14,17	1.18	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	3DR	D	7	1	-	1/3/15/16	0/1/1/1
1	3DR	G	7	1	-	1/3/15/16	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	7	3DR	O4'-C4'-C3'	2.65	107.63	103.73
1	G	7	3DR	O4'-C4'-C3'	2.65	107.62	103.73

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	D	7	3DR	O4'-C4'-C5'-O5'
1	G	7	3DR	O4'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	D	11/14 (78%)	0.23	0 100 100	60, 93, 139, 140	0
1	G	11/14 (78%)	0.28	0 100 100	66, 92, 126, 133	1 (9%)
2	A	259/275 (94%)	0.50	11 (4%) 36 37	38, 62, 104, 132	0
2	C	259/275 (94%)	0.48	9 (3%) 44 44	36, 61, 106, 120	0
3	B	14/14 (100%)	0.26	0 100 100	83, 94, 125, 132	2 (14%)
3	F	14/14 (100%)	0.30	2 (14%) 2 2	86, 93, 135, 136	3 (21%)
All	All	568/606 (93%)	0.47	22 (3%) 39 40	36, 64, 108, 140	6 (1%)

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	64	PHE	3.9
2	C	64	PHE	3.7
2	C	35	TRP	3.2
2	A	50	VAL	3.2
2	C	45	LEU	2.9
2	A	102	PHE	2.8
2	A	45	LEU	2.8
2	C	31	LEU	2.8
2	A	31	LEU	2.8
2	C	102	PHE	2.7
2	A	150	TYR	2.6
3	F	1	DA	2.5
2	A	219	GLY	2.5
2	C	150	TYR	2.5
2	C	50	VAL	2.5
2	A	128	HIS	2.4
2	A	120	LEU	2.2
2	C	23	VAL	2.1
3	F	14	DC	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	258	VAL	2.0
2	A	35	TRP	2.0
2	A	33	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	3DR	D	7	11/12	0.97	0.16	53,56,58,63	0
1	3DR	G	7	11/12	0.97	0.15	51,55,58,65	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MG	C	302	1/1	0.94	0.23	78,78,78,78	0
4	ZN	A	301	1/1	0.99	0.18	55,55,55,55	0
4	ZN	C	301	1/1	0.99	0.18	58,58,58,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.