



## Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 11:53 pm BST

PDB ID : 1TD2  
Title : Crystal Structure of the PdxY Protein from Escherichia coli  
Authors : Safo, M.K.; Musayev, F.N.; Hunt, S.; di Salvo, M.; Scarsdale, N.; Schirch, V.  
Deposited on : 2004-05-21  
Resolution : 2.22 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	6646 (2.24-2.20)
Ramachandran outliers	138981	6543 (2.24-2.20)
Sidechain outliers	138945	6544 (2.24-2.20)
RSRZ outliers	127900	5797 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	287	<div> <div>2%</div> <div>75%</div> <div>22%</div> <div>.</div> </div>
1	B	287	<div> <div>%</div> <div>71%</div> <div>26%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PXL	A	288	-	X	-	-

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4817 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyridoxamine kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	287	Total	C	N	O	S	0	0	0
			2201	1400	386	401	14			
1	B	287	Total	C	N	O	S	0	0	0
			2201	1400	386	401	14			

- Molecule 2 is 3-HYDROXY-5-(HYDROXYMETHYL)-2-METHYLISONICOTINALDEHYDE (three-letter code: PXL) (formula:  $C_8H_9NO_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			12	8	1	3		
2	B	1	Total	C	N	O	0	0
			12	8	1	3		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	215	Total	O	0	0
			215	215		
4	B	171	Total	O	0	0
			171	171		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Pyridoxamine kinase



#### • Molecule 1: Pyridoxamine kinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.13Å 67.38Å 73.59Å 90.00° 93.67° 90.00°	Depositor
Resolution (Å)	38.20 – 2.22 49.65 – 2.20	Depositor EDS
% Data completeness (in resolution range)	90.4 (38.20-2.22) 90.1 (49.65-2.20)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.16 (at 2.20Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.176 , 0.246 0.175 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.8	Xtriage
Anisotropy	0.426	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 58.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4817	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, PXL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.51	0/2249	0.77	2/3058 (0.1%)
1	B	0.48	0/2249	0.71	2/3058 (0.1%)
All	All	0.50	0/4498	0.74	4/6116 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	217	ARG	NE-CZ-NH2	7.29	123.94	120.30
1	A	217	ARG	NE-CZ-NH2	7.28	123.94	120.30
1	A	216	MET	CG-SD-CE	5.98	109.76	100.20
1	B	216	MET	CG-SD-CE	5.97	109.75	100.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2201	0	2210	63	0
1	B	2201	0	2211	60	0
2	A	12	0	8	2	0
2	B	12	0	9	2	0
3	B	5	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	215	0	0	3	0
4	B	171	0	0	5	0
All	All	4817	0	4438	120	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (120) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:ASN:HD22	1:A:186:ARG:NH2	1.56	1.03
1:A:159:ASN:ND2	1:A:186:ARG:HH22	1.56	1.01
1:A:186:ARG:HG2	4:A:386:HOH:O	1.61	0.98
1:A:8:ILE:N	1:A:8:ILE:HD13	1.85	0.90
1:A:6:LEU:HG	1:A:8:ILE:HD11	1.55	0.88
1:A:229:LEU:HB3	1:A:247:VAL:HG22	1.63	0.80
1:B:3:LYS:HD2	1:B:3:LYS:N	1.98	0.78
1:A:19:ASN:HD22	1:A:37:ASN:HD22	1.36	0.72
1:A:160:ASN:ND2	1:A:163:GLU:HG3	2.06	0.70
1:A:122:CYS:SG	2:A:288:PXL:C4	2.82	0.68
1:A:85:GLY:H	1:A:89:GLN:HE22	1.42	0.68
1:A:7:ALA:C	1:A:8:ILE:HD13	2.15	0.67
1:A:40:GLN:HE22	1:B:37:ASN:HB2	1.59	0.66
1:A:117:HIS:O	1:A:119:GLU:N	2.29	0.66
1:A:85:GLY:H	1:A:89:GLN:NE2	1.95	0.65
1:A:184:LEU:HD12	1:A:195:GLU:HB3	1.79	0.64
1:A:19:ASN:HD22	1:A:37:ASN:ND2	1.95	0.64
1:A:44:HIS:HD2	1:A:46:GLN:H	1.43	0.64
1:B:229:LEU:HB3	1:B:247:VAL:HG22	1.79	0.63
1:B:44:HIS:HD2	1:B:46:GLN:H	1.46	0.63
1:B:85:GLY:H	1:B:89:GLN:NE2	1.97	0.63
1:B:88:GLU:CD	1:B:88:GLU:H	2.01	0.62
1:A:44:HIS:HE1	4:A:330:HOH:O	1.83	0.61
1:A:160:ASN:HD21	1:A:163:GLU:HG3	1.64	0.61
1:B:216:MET:O	1:B:217:ARG:CG	2.49	0.61
1:B:174:GLN:HG3	4:B:620:HOH:O	2.00	0.61
1:B:130:GLU:O	1:B:134:ARG:HG2	2.01	0.60
1:A:241:GLN:O	1:A:245:GLU:HG3	2.02	0.60
1:B:160:ASN:ND2	1:B:162:GLU:HB3	2.17	0.59
1:B:85:GLY:H	1:B:89:GLN:HE22	1.51	0.58
1:A:180:LEU:HG	1:A:230:LEU:HD22	1.85	0.58

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:216:MET:O	1:B:217:ARG:HG3	2.03	0.58
1:A:118:PRO:HA	1:A:123:ILE:HD13	1.86	0.58
1:B:245:GLU:HG2	1:B:282:PHE:O	2.04	0.58
1:A:73:LEU:HD22	1:A:100:VAL:HG22	1.86	0.57
1:B:147:LEU:HD22	1:B:184:LEU:O	2.04	0.57
1:B:210:PRO:HG2	1:B:252:TYR:CE1	2.39	0.57
1:A:117:HIS:N	1:A:118:PRO:HD3	2.20	0.56
1:A:186:ARG:HB3	1:A:191:ARG:NH1	2.21	0.56
1:B:167:ALA:O	1:B:171:LEU:HD13	2.06	0.56
1:A:254:ILE:HG22	1:A:266:LEU:HD22	1.86	0.56
1:B:180:LEU:HD13	1:B:182:LYS:HB3	1.87	0.56
1:B:253:GLU:OE1	1:B:280:HIS:HD2	1.88	0.56
1:A:8:ILE:HD12	1:A:36:LEU:HB3	1.88	0.56
1:B:216:MET:C	1:B:217:ARG:HG3	2.27	0.55
1:A:159:ASN:HD22	1:A:186:ARG:HH22	0.75	0.55
1:A:24:PHE:HB3	1:A:25:PRO:HD3	1.90	0.54
1:A:116:GLY:C	1:A:118:PRO:HD3	2.28	0.54
1:B:180:LEU:HG	1:B:230:LEU:HD22	1.89	0.54
1:A:65:GLN:HA	1:A:65:GLN:OE1	2.08	0.53
1:B:194:PHE:O	1:B:208:SER:HA	2.08	0.53
3:B:288:SO4:O4	2:B:289:PXL:H5A1	2.09	0.53
1:B:144:ALA:HA	1:B:180:LEU:O	2.08	0.53
1:B:24:PHE:HB3	1:B:25:PRO:HD3	1.92	0.52
1:A:94:LEU:HD11	1:A:135:HIS:HB3	1.92	0.52
1:A:274:ARG:NH1	4:A:326:HOH:O	2.24	0.52
1:A:39:VAL:HG12	1:A:54:VAL:HA	1.92	0.52
1:B:160:ASN:ND2	1:B:163:GLU:H	2.06	0.52
1:A:8:ILE:CD1	1:A:8:ILE:N	2.59	0.51
1:A:254:ILE:CG2	1:A:266:LEU:HD22	2.39	0.51
1:A:186:ARG:HB3	1:A:191:ARG:HH11	1.75	0.51
1:A:94:LEU:CD1	1:A:135:HIS:HB3	2.41	0.51
1:B:225:VAL:HG12	1:B:229:LEU:HD22	1.92	0.51
1:B:195:GLU:OE1	1:B:206:HIS:HE1	1.93	0.50
1:B:28:ARG:HD3	1:B:272:GLN:HB2	1.94	0.50
1:A:193:ARG:NH1	1:A:210:PRO:HG3	2.27	0.49
1:B:202:ASP:OD1	1:B:203:GLU:HG2	2.11	0.49
1:B:195:GLU:OE1	1:B:206:HIS:CE1	2.65	0.49
1:A:6:LEU:HG	1:A:8:ILE:CD1	2.36	0.49
1:B:36:LEU:HD13	1:B:63:ILE:HG21	1.95	0.49
1:A:44:HIS:HD2	1:A:46:GLN:N	2.10	0.48
1:A:159:ASN:ND2	1:A:186:ARG:NH2	2.34	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:184:LEU:HD21	1:A:197:LEU:HD22	1.96	0.48
1:B:254:ILE:HG21	1:B:266:LEU:HG	1.96	0.47
1:B:182:LYS:HA	1:B:196:MET:HG2	1.96	0.47
1:B:214:PHE:HB2	1:B:218:GLN:OE1	2.15	0.47
1:B:19:ASN:HD22	1:B:37:ASN:HD22	1.62	0.47
1:A:166:LEU:O	1:A:170:GLU:HG3	2.14	0.47
1:B:44:HIS:HD2	1:B:46:GLN:N	2.13	0.47
1:A:56:PRO:HG2	1:A:59:HIS:HB2	1.97	0.46
1:B:104:ASN:ND2	1:B:106:GLN:H	2.13	0.46
1:B:115:MET:SD	1:B:152:ILE:HD12	2.56	0.46
1:B:172:ILE:HA	1:B:176:PRO:O	2.15	0.46
1:B:46:GLN:HG3	1:B:121:GLY:HA3	1.97	0.45
1:B:86:SER:OG	1:B:88:GLU:HG2	2.16	0.45
1:A:118:PRO:HG3	1:A:124:VAL:H	1.80	0.45
1:A:209:ARG:HD3	1:A:252:TYR:CD1	2.52	0.45
1:B:34:TRP:CZ3	1:B:76:CYS:HB2	2.52	0.45
1:A:144:ALA:HA	1:A:180:LEU:O	2.17	0.45
1:B:134:ARG:HG3	1:B:135:HIS:CD2	2.51	0.45
1:A:27:ARG:HA	1:A:31:ALA:O	2.17	0.45
1:B:148:VAL:HG23	4:B:487:HOH:O	2.16	0.44
1:A:194:PHE:HB3	1:A:209:ARG:CG	2.47	0.44
1:A:38:THR:HB	1:A:89:GLN:HE22	1.83	0.44
1:A:37:ASN:H	1:B:40:GLN:HE22	1.64	0.44
1:B:102:ALA:HB3	4:B:479:HOH:O	2.18	0.43
1:B:195:GLU:HG2	1:B:208:SER:HB3	2.00	0.43
1:B:254:ILE:CG2	1:B:266:LEU:HG	2.48	0.43
1:A:40:GLN:HE22	1:B:37:ASN:CB	2.28	0.43
1:B:39:VAL:HB	1:B:41:PHE:CE2	2.54	0.43
1:B:216:MET:HG2	4:B:667:HOH:O	2.19	0.43
1:B:122:CYS:SG	2:B:289:PXL:C4A	3.07	0.43
1:A:209:ARG:O	1:A:209:ARG:HG3	2.15	0.42
1:B:151:GLU:OE2	1:B:157:ALA:HA	2.19	0.42
1:B:194:PHE:HB3	1:B:209:ARG:HG3	2.01	0.42
1:A:8:ILE:HG22	1:A:38:THR:CG2	2.49	0.42
1:B:184:LEU:O	1:B:185:ALA:HB3	2.20	0.42
1:A:147:LEU:C	1:A:147:LEU:HD13	2.40	0.42
1:A:56:PRO:O	1:A:59:HIS:HB3	2.19	0.42
1:B:33:VAL:O	1:B:35:PRO:HD3	2.19	0.42
1:B:73:LEU:HB2	4:B:631:HOH:O	2.19	0.42
1:A:115:MET:HG3	1:A:149:GLU:HG2	2.03	0.41
1:A:134:ARG:HH11	1:A:134:ARG:HA	1.86	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:224:ASP:OD2	2:A:288:PXL:OP4	2.35	0.41
1:A:24:PHE:HE1	1:A:272:GLN:HB3	1.85	0.41
1:A:125:ALA:HA	1:A:126:PRO:HD3	1.93	0.40
1:B:28:ARG:CD	1:B:272:GLN:HB2	2.51	0.40
1:A:35:PRO:O	1:B:11:HIS:HE1	2.03	0.40
1:B:125:ALA:HA	1:B:126:PRO:HD3	1.95	0.40
1:B:186:ARG:HA	1:B:191:ARG:HG2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/287 (99%)	276 (97%)	7 (2%)	2 (1%)	22	21
1	B	285/287 (99%)	270 (95%)	13 (5%)	2 (1%)	22	21
All	All	570/574 (99%)	546 (96%)	20 (4%)	4 (1%)	22	21

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	118	PRO
1	A	119	GLU
1	B	183	HIS
1	B	216	MET

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/232 (100%)	220 (95%)	12 (5%)	23	27
1	B	232/232 (100%)	220 (95%)	12 (5%)	23	27
All	All	464/464 (100%)	440 (95%)	24 (5%)	23	27

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	ILE
1	A	28	ARG
1	A	110	PHE
1	A	117	HIS
1	A	120	LYS
1	A	134	ARG
1	A	145	PRO
1	A	160	ASN
1	A	171	LEU
1	A	180	LEU
1	A	197	LEU
1	A	229	LEU
1	B	3	LYS
1	B	28	ARG
1	B	73	LEU
1	B	88	GLU
1	B	104	ASN
1	B	150	LEU
1	B	166	LEU
1	B	177	GLN
1	B	180	LEU
1	B	196	MET
1	B	229	LEU
1	B	255	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	ASN
1	A	40	GLN
1	A	44	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	89	GLN
1	A	104	ASN
1	A	106	GLN
1	A	159	ASN
1	A	160	ASN
1	A	236	GLN
1	A	267	GLN
1	A	280	HIS
1	B	11	HIS
1	B	32	ASN
1	B	37	ASN
1	B	40	GLN
1	B	44	HIS
1	B	89	GLN
1	B	104	ASN
1	B	160	ASN
1	B	206	HIS
1	B	267	GLN
1	B	280	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PXL	B	289	-	12,12,12	3.13	7 (58%)	15,16,16	1.58	4 (26%)
2	PXL	A	288	1	12,12,12	3.17	9 (75%)	15,16,16	2.24	2 (13%)
3	SO4	B	288	-	4,4,4	0.50	0	6,6,6	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PXL	B	289	-	-	0/4/4/4	0/1/1/1
2	PXL	A	288	1	-	4/4/4/4	0/1/1/1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	288	PXL	O4A-C4A	6.10	1.42	1.21
2	B	289	PXL	C2-N1	5.14	1.43	1.33
2	B	289	PXL	C6-N1	4.63	1.44	1.34
2	A	288	PXL	C3-C2	4.12	1.45	1.40
2	B	289	PXL	C3-C2	4.08	1.45	1.40
2	B	289	PXL	C4-C3	3.68	1.46	1.40
2	A	288	PXL	C6-N1	3.54	1.41	1.34
2	A	288	PXL	C4-C5	3.54	1.46	1.42
2	B	289	PXL	C4-C5	3.40	1.46	1.42
2	B	289	PXL	C6-C5	3.24	1.44	1.37
2	B	289	PXL	C4-C4A	3.22	1.53	1.46
2	A	288	PXL	C4-C4A	3.15	1.53	1.46
2	A	288	PXL	C2-N1	2.98	1.39	1.33
2	A	288	PXL	C4-C3	2.96	1.45	1.40
2	A	288	PXL	C6-C5	2.19	1.42	1.37
2	A	288	PXL	C2A-C2	2.14	1.54	1.50

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	288	PXL	O4A-C4A-C4	-7.18	109.26	124.91
2	B	289	PXL	O4A-C4A-C4	-3.39	117.53	124.91

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	288	PXL	C5-C6-N1	-2.70	119.32	123.82
2	B	289	PXL	C5-C6-N1	-2.54	119.59	123.82
2	B	289	PXL	C3-C4-C5	2.42	120.12	118.26
2	B	289	PXL	C5A-C5-C4	-2.01	119.52	122.37

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	288	PXL	C3-C4-C4A-O4A
2	A	288	PXL	C6-C5-C5A-OP4
2	A	288	PXL	C4-C5-C5A-OP4
2	A	288	PXL	C5-C4-C4A-O4A

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	289	PXL	2	0
2	A	288	PXL	2	0
3	B	288	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	287/287 (100%)	-0.48	5 (1%) 70 68	10, 20, 40, 94	0
1	B	287/287 (100%)	-0.40	2 (0%) 87 86	12, 24, 48, 74	0
All	All	574/574 (100%)	-0.44	7 (1%) 79 77	10, 21, 45, 94	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	118	PRO	7.7
1	B	217	ARG	4.6
1	A	216	MET	3.8
1	B	216	MET	3.8
1	A	119	GLU	3.5
1	A	120	LYS	2.8
1	A	217	ARG	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PXL	A	288	12/12	0.93	0.13	14,18,35,36	0
3	SO4	B	288	5/5	0.95	0.19	49,54,65,69	0
2	PXL	B	289	12/12	0.96	0.12	13,23,31,40	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.