



# Full wwPDB NMR Structure Validation Report ⓘ

Aug 16, 2020 – 08:18 PM BST

PDB ID : 6TDN  
Title : Bam\_5925cDD 5924nDD docking domains  
Authors : Risser, F.; Chagot, B.  
Deposited on : 2019-11-08

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.13.1  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 92%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	87	
1	B	87	

## 2 Ensemble composition and analysis

This entry contains 25 models. Model 3 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *fewest violations*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:11-A:32, A:56-A:82, B:111-B:132, B:156-B:182 (98)	0.31	3

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 4 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 6, 7, 8, 9, 10, 13, 14, 18, 23, 24
2	15, 22
3	5, 19
4	4, 21
5	17, 20
Single-model clusters	11; 12; 16; 25

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2540 atoms, of which 1258 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Beta-ketoacyl synthase,Beta-ketoacyl synthase.

Mol	Chain	Residues	Atoms						Trace
1	A	87	Total	C	H	N	O	S	0
			1270	392	629	111	136	2	
1	B	87	Total	C	H	N	O	S	0
			1270	392	629	111	136	2	

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP Q0B303
A	2	PRO	-	expression tag	UNP Q0B303
A	3	GLY	-	expression tag	UNP Q0B303
A	4	SER	-	expression tag	UNP Q0B303
A	5	TYR	-	expression tag	UNP Q0B303
A	35	GLY	-	linker	UNP Q0B303
A	36	GLY	-	linker	UNP Q0B303
A	37	GLY	-	linker	UNP Q0B303
A	38	SER	-	linker	UNP Q0B303
A	39	GLY	-	linker	UNP Q0B303
A	40	GLY	-	linker	UNP Q0B303
A	41	GLY	-	linker	UNP Q0B303
A	42	SER	-	linker	UNP Q0B303
A	43	GLY	-	linker	UNP Q0B303
A	44	GLY	-	linker	UNP Q0B303
A	45	GLY	-	linker	UNP Q0B303
A	46	SER	-	linker	UNP Q0B303
B	101	GLY	-	expression tag	UNP Q0B303
B	102	PRO	-	expression tag	UNP Q0B303
B	103	GLY	-	expression tag	UNP Q0B303
B	104	SER	-	expression tag	UNP Q0B303
B	105	TYR	-	expression tag	UNP Q0B303
B	135	GLY	-	linker	UNP Q0B303
B	136	GLY	-	linker	UNP Q0B303
B	137	GLY	-	linker	UNP Q0B303
B	138	SER	-	linker	UNP Q0B303
B	139	GLY	-	linker	UNP Q0B303
B	140	GLY	-	linker	UNP Q0B303
B	141	GLY	-	linker	UNP Q0B303

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	142	SER	-	linker	UNP Q0B303
B	143	GLY	-	linker	UNP Q0B303
B	144	GLY	-	linker	UNP Q0B303
B	145	GLY	-	linker	UNP Q0B303
B	146	SER	-	linker	UNP Q0B303

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble


These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.2 Score per residue for model 2

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.3 Score per residue for model 3 (medoid)

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.4 Score per residue for model 4

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.5 Score per residue for model 5

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.6 Score per residue for model 6

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.7 Score per residue for model 7

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase






### 4.2.8 Score per residue for model 8

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.9 Score per residue for model 9

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.10 Score per residue for model 10

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.11 Score per residue for model 11

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.12 Score per residue for model 12

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.13 Score per residue for model 13

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




#### 4.2.14 Score per residue for model 14

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




#### 4.2.15 Score per residue for model 15

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




#### 4.2.16 Score per residue for model 16

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.17 Score per residue for model 17

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.18 Score per residue for model 18

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.19 Score per residue for model 19

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




#### 4.2.20 Score per residue for model 20

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




#### 4.2.21 Score per residue for model 21

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




#### 4.2.22 Score per residue for model 22

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.23 Score per residue for model 23

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.24 Score per residue for model 24

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase




### 4.2.25 Score per residue for model 25

- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase



- Molecule 1: Beta-ketoacyl synthase,Beta-ketoacyl synthase



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 50 calculated structures, 25 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	refinement	
CYANA	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1938
Number of shifts mapped to atoms	1938
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	92%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality i

### 6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.68±0.01	0±0/400 ( 0.0± 0.0%)	0.97±0.03	1±1/536 ( 0.1± 0.1%)
1	B	0.68±0.01	0±0/400 ( 0.0± 0.0%)	0.97±0.04	1±1/536 ( 0.2± 0.1%)
All	All	0.68	0/20000 ( 0.0%)	0.97	43/26800 ( 0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	B	0.0±0.0	0.1±0.3
All	All	0	2

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	63	ARG	NE-CZ-NH2	9.10	124.85	120.30	22	2
1	A	63	ARG	NE-CZ-NH1	8.36	124.48	120.30	4	10
1	B	163	ARG	NE-CZ-NH1	7.76	124.18	120.30	1	14
1	B	166	ARG	NE-CZ-NH2	7.75	124.18	120.30	3	2
1	B	166	ARG	NE-CZ-NH1	7.34	123.97	120.30	25	8
1	A	66	ARG	NE-CZ-NH2	6.46	123.53	120.30	1	1
1	B	163	ARG	NE-CZ-NH2	-6.39	117.10	120.30	10	2
1	A	66	ARG	NE-CZ-NH1	6.12	123.36	120.30	10	4

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.



Mol	Chain	Res	Type	Group	Models (Total)
1	B	166	ARG	Sidechain	1
1	B	163	ARG	Sidechain	1

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	19800	20500	20500	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	49/87 (56%)	49±0 (100±1%)	0±0 (0±1%)	0±0 (0±0%)	54 85
1	B	49/87 (56%)	49±1 (99±1%)	0±0 (0±1%)	0±0 (0±1%)	50 82
All	All	2450/4350 (56%)	2438 (100%)	8 (0%)	4 (0%)	50 82

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	128	GLY	2
1	A	28	GLY	1
1	B	129	LYS	1

### 6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	43/67 (64%)	41±1 (95±3%)	2±1 (5±3%)	27	76
1	B	43/67 (64%)	41±1 (95±2%)	2±1 (5±2%)	26	75
All	All	2150/3350 (64%)	2036 (95%)	114 (5%)	26	75

All 15 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	119	GLN	25
1	A	19	GLN	25
1	B	126	LEU	13
1	A	26	LEU	12
1	B	177	LEU	8
1	A	77	LEU	8
1	B	157	LYS	4
1	A	57	LYS	4
1	A	75	LEU	3
1	B	129	LYS	3
1	B	111	GLU	3
1	A	29	LYS	3
1	B	175	LEU	1
1	B	130	GLU	1
1	A	30	GLU	1

### 6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 92% for the well-defined parts and 88% for the entire structure.

### 7.1 Chemical shift list 1

File name: `working_cs.cif`

Chemical shift list name: *starch\_output*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1938
Number of shifts mapped to atoms	1938
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	152	$-0.47 \pm 0.06$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}_\beta$	140	$0.06 \pm 0.05$	None needed ( $< 0.5$ ppm)
$^{13}\text{C}'$	152	$-0.47 \pm 0.07$	None needed ( $< 0.5$ ppm)
$^{15}\text{N}$	154	$0.12 \pm 0.09$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 92%, i.e. 1170 atoms were assigned a chemical shift out of a possible 1266. 24 out of 24 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	490/490 (100%)	196/196 (100%)	196/196 (100%)	98/98 (100%)
Sidechain	630/718 (88%)	386/418 (92%)	230/268 (86%)	14/32 (44%)

*Continued on next page...*

Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	50/58 (86%)	30/30 (100%)	18/26 (69%)	2/2 (100%)
Overall	1170/1266 (92%)	612/644 (95%)	444/490 (91%)	114/132 (86%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 88%, i.e. 1744 atoms were assigned a chemical shift out of a possible 1978. 28 out of 28 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	774/858 (90%)	316/342 (92%)	304/348 (87%)	154/168 (92%)
Sidechain	908/1046 (87%)	564/616 (92%)	322/384 (84%)	22/46 (48%)
Aromatic	62/74 (84%)	38/38 (100%)	22/34 (65%)	2/2 (100%)
Overall	1744/1978 (88%)	918/996 (92%)	648/766 (85%)	178/216 (82%)


#### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The images below report *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:

