



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 6, 2020 – 07:04 PM BST

PDB ID : 5TGO  
Title : Crystal structure of H10 hemagglutinin mutant (K158aA-D193T-Q226L-G228S) from Jiangxi-Donghu (2013) H10N8 influenza virus  
Authors : Tzarum, N.; Wilson, I.A.  
Deposited on : 2016-09-28  
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	<div> <div></div> <div>81%17%•</div> </div>
1	C	323	<div> <div>%</div> <div>82%16%•</div> </div>
1	E	323	<div> <div>%</div> <div>77%21%•</div> </div>
2	B	180	<div> <div>4%</div> <div>74%21%5%</div> </div>
2	D	180	<div> <div>3%</div> <div>69%23%• 6%</div> </div>
2	F	180	<div> <div>2%</div> <div>68%26%• 5%</div> </div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	G	3	 100%
4	H	2	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MAN	G	3	-	-	-	X
4	NAG	H	2	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 12147 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	318	Total	C	N	O	S	0	0	0
			2433	1505	447	464	17			
1	C	319	Total	C	N	O	S	0	0	0
			2441	1511	448	465	17			
1	E	318	Total	C	N	O	S	0	0	0
			2433	1505	447	464	17			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	158A	ALA	LYS	engineered mutation	UNP A0A0J9X252
A	193	THR	ASP	engineered mutation	UNP A0A0J9X252
A	226	LEU	GLN	engineered mutation	UNP A0A0J9X252
A	228	SER	GLY	engineered mutation	UNP A0A0J9X252
C	158A	ALA	LYS	engineered mutation	UNP A0A0J9X252
C	193	THR	ASP	engineered mutation	UNP A0A0J9X252
C	226	LEU	GLN	engineered mutation	UNP A0A0J9X252
C	228	SER	GLY	engineered mutation	UNP A0A0J9X252
E	158A	ALA	LYS	engineered mutation	UNP A0A0J9X252
E	193	THR	ASP	engineered mutation	UNP A0A0J9X252
E	226	LEU	GLN	engineered mutation	UNP A0A0J9X252
E	228	SER	GLY	engineered mutation	UNP A0A0J9X252


- Molecule 2 is a protein called Hemagglutinin HA2 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	171	Total	C	N	O	S	0	0	0
			1382	854	239	281	8			
2	D	169	Total	C	N	O	S	0	0	0
			1367	846	236	277	8			
2	F	171	Total	C	N	O	S	0	0	0
			1382	854	239	281	8			

There are 21 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	175	SER	-	expression tag	UNP A0A0J9X253
B	176	GLY	-	expression tag	UNP A0A0J9X253
B	177	ARG	-	expression tag	UNP A0A0J9X253
B	178	LEU	-	expression tag	UNP A0A0J9X253
B	179	VAL	-	expression tag	UNP A0A0J9X253
B	180	PRO	-	expression tag	UNP A0A0J9X253
B	181	ARG	-	expression tag	UNP A0A0J9X253
D	175	SER	-	expression tag	UNP A0A0J9X253
D	176	GLY	-	expression tag	UNP A0A0J9X253
D	177	ARG	-	expression tag	UNP A0A0J9X253
D	178	LEU	-	expression tag	UNP A0A0J9X253
D	179	VAL	-	expression tag	UNP A0A0J9X253
D	180	PRO	-	expression tag	UNP A0A0J9X253
D	181	ARG	-	expression tag	UNP A0A0J9X253
F	175	SER	-	expression tag	UNP A0A0J9X253
F	176	GLY	-	expression tag	UNP A0A0J9X253
F	177	ARG	-	expression tag	UNP A0A0J9X253
F	178	LEU	-	expression tag	UNP A0A0J9X253
F	179	VAL	-	expression tag	UNP A0A0J9X253
F	180	PRO	-	expression tag	UNP A0A0J9X253
F	181	ARG	-	expression tag	UNP A0A0J9X253

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	G	3	Total	C	N	O	0	0	0
			36	20	1	15			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	H	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	158	Total	O	0	0
			158	158		
6	B	55	Total	O	0	0
			55	55		
6	C	133	Total	O	0	0
			133	133		
6	D	56	Total	O	0	0
			56	56		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	E	123	Total 123	O 123	0	0
6	F	64	Total 64	O 64	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hemagglutinin HA1 chain



- Molecule 1: Hemagglutinin HA1 chain




- Molecule 1: Hemagglutinin HA1 chain




- Molecule 2: Hemagglutinin HA2 chain







- Molecule 2: Hemagglutinin HA2 chain



- Molecule 2: Hemagglutinin HA2 chain



- Molecule 3: alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.96 Å   253.93 Å   69.93 Å 90.00°   111.73°   90.00°	Depositor
Resolution (Å)	37.37 – 2.35 48.63 – 2.34	Depositor EDS
% Data completeness (in resolution range)	95.2 (37.37-2.35) 90.1 (48.63-2.34)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.22 (at 2.34 Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.200   ,   0.242 0.202   ,   0.242	Depositor DCC
$R_{free}$ test set	4107 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.2	Xtriage
Anisotropy	0.166	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 33.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	12147	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.61	1/2482 (0.0%)	0.70	0/3365
1	C	0.53	0/2490	0.69	1/3376 (0.0%)
1	E	0.56	0/2482	0.68	2/3365 (0.1%)
2	B	0.50	0/1407	0.67	0/1899
2	D	0.50	0/1391	0.63	0/1875
2	F	0.55	1/1407 (0.1%)	0.67	0/1899
All	All	0.55	2/11659 (0.0%)	0.68	3/15779 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	277	CYS	CB-SG	-5.05	1.73	1.81
2	F	144	CYS	CB-SG	5.02	1.90	1.82

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	220	ARG	C-N-CD	6.77	142.61	128.40
1	E	85	ASP	CB-CG-OD2	-5.48	113.36	118.30
1	C	286	GLY	N-CA-C	5.11	125.86	113.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2433	0	2385	50	0
1	C	2441	0	2396	43	0
1	E	2433	0	2386	49	0
2	B	1382	0	1279	32	0
2	D	1367	0	1265	36	0
2	F	1382	0	1279	42	0
3	G	36	0	31	5	0
4	H	28	0	25	1	0
5	A	28	0	26	1	0
5	C	28	0	25	5	0
6	A	158	0	0	16	0
6	B	55	0	0	3	0
6	C	133	0	0	14	0
6	D	56	0	0	8	0
6	E	123	0	0	6	0
6	F	64	0	0	6	0
All	All	12147	0	11097	221	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 221 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:501:NAG:O4	3:G:1:NAG:C1	1.66	1.42
5:C:501:NAG:C4	3:G:1:NAG:C1	2.44	0.95
5:C:501:NAG:O4	3:G:1:NAG:C2	2.29	0.80
2:B:53:ASN:O	2:B:57:GLU:HG2	1.84	0.78
1:A:276:ASN:O	6:A:501:HOH:O	2.01	0.78

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	316/323 (98%)	309 (98%)	7 (2%)	0	100	100
1	C	317/323 (98%)	311 (98%)	5 (2%)	1 (0%)	41	47
1	E	316/323 (98%)	309 (98%)	6 (2%)	1 (0%)	41	47
2	B	169/180 (94%)	164 (97%)	5 (3%)	0	100	100
2	D	165/180 (92%)	160 (97%)	5 (3%)	0	100	100
2	F	169/180 (94%)	166 (98%)	3 (2%)	0	100	100
All	All	1452/1509 (96%)	1419 (98%)	31 (2%)	2 (0%)	51	63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	286	GLY
1	E	220	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	269/272 (99%)	267 (99%)	2 (1%)	84	91
1	C	270/272 (99%)	269 (100%)	1 (0%)	91	95
1	E	269/272 (99%)	266 (99%)	3 (1%)	73	84
2	B	146/154 (95%)	145 (99%)	1 (1%)	84	91
2	D	144/154 (94%)	141 (98%)	3 (2%)	53	65
2	F	146/154 (95%)	143 (98%)	3 (2%)	53	65
All	All	1244/1278 (97%)	1231 (99%)	13 (1%)	76	85

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	102	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	D	155	ASN
2	F	58	LYS
2	D	71	SER
1	E	223	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	95	GLN
2	D	125	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	G	1	3	14,14,15	0.22	0	17,19,21	0.53	0
3	BMA	G	2	3	11,11,12	1.46	2 (18%)	15,15,17	0.89	1 (6%)
3	MAN	G	3	3	11,11,12	1.57	3 (27%)	15,15,17	1.25	3 (20%)
4	NAG	H	1	1,4	14,14,15	1.17	2 (14%)	17,19,21	1.21	1 (5%)
4	NAG	H	2	4	14,14,15	0.34	0	17,19,21	0.79	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	G	1	3	-	2/6/23/26	0/1/1/1
3	BMA	G	2	3	-	0/2/19/22	0/1/1/1
3	MAN	G	3	3	-	0/2/19/22	0/1/1/1
4	NAG	H	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	H	2	4	-	0/6/23/26	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	3	MAN	O5-C1	3.21	1.48	1.43
3	G	2	BMA	O5-C1	3.05	1.48	1.43
3	G	2	BMA	O5-C5	2.62	1.48	1.43
3	G	3	MAN	O5-C5	2.43	1.48	1.43
4	H	1	NAG	O5-C1	-2.39	1.39	1.43

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	3	MAN	C1-O5-C5	2.74	115.90	112.19
3	G	3	MAN	C1-C2-C3	2.59	112.85	109.67
4	H	1	NAG	O5-C5-C6	-2.23	103.70	107.20
3	G	2	BMA	C1-O5-C5	2.14	115.09	112.19
3	G	3	MAN	C6-C5-C4	2.01	117.71	113.00

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	H	1	NAG	O5-C5-C6-O6
4	H	1	NAG	C4-C5-C6-O6
3	G	1	NAG	O5-C5-C6-O6
3	G	1	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	1	NAG	1	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	1	NAG	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	C	500	1	14,14,15	0.85	1 (7%)	17,19,21	1.11	1 (5%)
5	NAG	A	401	1	14,14,15	0.77	1 (7%)	17,19,21	0.88	0
5	NAG	C	501	1	14,14,15	1.06	2 (14%)	17,19,21	1.47	3 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	A	400	1	14,14,15	0.84	1 (7%)	17,19,21	1.15	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	C	500	1	-	5/6/23/26	0/1/1/1
5	NAG	A	401	1	-	0/6/23/26	0/1/1/1
5	NAG	C	501	1	-	0/6/23/26	0/1/1/1
5	NAG	A	400	1	-	4/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	501	NAG	O5-C1	-2.46	1.39	1.43
5	A	401	NAG	O5-C1	-2.46	1.39	1.43
5	C	500	NAG	O5-C1	-2.39	1.39	1.43
5	A	400	NAG	O5-C1	-2.26	1.40	1.43
5	C	501	NAG	O7-C7	-2.09	1.18	1.23

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	501	NAG	C1-C2-N2	-3.36	104.75	110.49
5	C	501	NAG	C6-C5-C4	-3.02	105.93	113.00
5	A	400	NAG	C2-N2-C7	-2.80	118.91	122.90
5	C	500	NAG	C1-C2-N2	-2.54	106.14	110.49
5	C	501	NAG	O3-C3-C2	2.48	114.61	109.47

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	500	NAG	C8-C7-N2-C2
5	C	500	NAG	O7-C7-N2-C2
5	C	500	NAG	O5-C5-C6-O6
5	A	400	NAG	C8-C7-N2-C2
5	A	400	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	501	NAG	5	0
5	A	400	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	318/323 (98%)	-0.33	0 100 100	22, 32, 52, 75	0
1	C	319/323 (98%)	-0.27	2 (0%) 89 93	24, 35, 53, 75	0
1	E	318/323 (98%)	-0.19	4 (1%) 77 84	23, 36, 56, 76	0
2	B	171/180 (95%)	0.32	7 (4%) 37 49	26, 47, 65, 72	0
2	D	169/180 (93%)	0.24	5 (2%) 50 61	26, 48, 66, 73	0
2	F	171/180 (95%)	0.11	3 (1%) 68 77	25, 43, 69, 85	0
All	All	1466/1509 (97%)	-0.09	21 (1%) 75 83	22, 38, 62, 85	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	58	LYS	4.6
2	F	2	LEU	4.5
2	D	156	THR	4.0
1	C	326	LEU	3.6
2	F	59	THR	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAG	H	2	14/15	0.73	0.43	72,85,92,99	0
3	MAN	G	3	11/12	0.74	0.47	89,93,101,107	0
3	BMA	G	2	11/12	0.75	0.19	83,89,96,96	0
4	NAG	H	1	14/15	0.87	0.24	29,54,67,79	0
3	NAG	G	1	14/15	0.91	0.14	68,80,92,95	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain G:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	NAG	C	500	14/15	0.62	0.39	55,78,82,85	0
5	NAG	A	400	14/15	0.84	0.25	52,64,72,76	0
5	NAG	A	401	14/15	0.85	0.18	26,51,59,67	0
5	NAG	C	501	14/15	0.88	0.12	30,50,59,69	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.