



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 05:18 am BST

PDB ID : 4TK1  
Title : Geph E in complex with a GABA receptor alpha3 subunit derived peptide in space group P21212  
Authors : Kasaragod, V.B.; Maric, H.M.; Schindelin, H.  
Deposited on : 2014-05-25  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	419	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>13%</div> <div>..</div> </div> </div>
1	B	419	<div> <div>5%</div> <div> <div></div> <div>84%</div> <div>11%</div> <div>.</div> </div> </div>
2	C	11	<div> <div>9%</div> <div> <div></div> <div>100%</div> </div> </div>
2	D	11	<div> <div></div> <div> <div>64%</div> <div>18%</div> <div>18%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	BME	A	802	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6500 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Gephyrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	414	Total	C	N	O	S	0	0	0
			3155	1990	547	599	19			
1	B	401	Total	C	N	O	S	0	0	0
			3054	1933	528	574	19			

- Molecule 2 is a protein called Gamma-aminobutyric acid receptor subunit alpha-3.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	11	Total	C	N	O	0	0	0
			88	58	13	17			
2	D	9	Total	C	N	O	0	0	0
			71	48	10	13			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C<sub>2</sub>H<sub>6</sub>OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			4	2	1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	55	Total	O	0	0
			55	55		
5	B	58	Total	O	0	0
			58	58		
5	C	1	Total	O	0	0
			1	1		
5	D	2	Total	O	0	0
			2	2		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Gephyrin



- Molecule 1: Gephyrin



- Molecule 2: Gamma-aminobutyric acid receptor subunit alpha-3



- Molecule 2: Gamma-aminobutyric acid receptor subunit alpha-3





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	110.44Å 157.69Å 51.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.54 – 2.70 48.54 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.5 (48.54-2.70) 99.6 (48.54-2.70)	Depositor EDS
$R_{merge}$	0.21	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.91 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.215 , 0.264 0.219 , 0.271	Depositor DCC
$R_{free}$ test set	1255 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.9	Xtriage
Anisotropy	0.488	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 40.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6500	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.43	0/3214	0.67	2/4369 (0.0%)
1	B	0.45	0/3113	0.66	1/4234 (0.0%)
2	C	0.54	0/90	0.74	0/122
2	D	0.47	0/73	0.48	0/100
All	All	0.44	0/6490	0.67	3/8825 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	564	ARG	NE-CZ-NH1	6.22	123.41	120.30
1	B	564	ARG	NE-CZ-NH1	6.11	123.35	120.30
1	A	438	ASP	N-CA-C	-5.34	96.59	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3155	0	3197	44	0
1	B	3054	0	3104	31	0
2	C	88	0	84	0	0
2	D	71	0	67	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	4	0	6	5	0
5	A	55	0	0	0	0
5	B	58	0	0	0	0
5	C	1	0	0	0	0
5	D	2	0	0	0	0
All	All	6500	0	6474	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:448:LEU:O	1:A:449:VAL:HG23	1.51	1.10
1:A:383:GLY:HA2	1:A:450:GLN:OE1	1.71	0.90
1:B:447:ILE:HG23	1:B:449:VAL:CG2	2.03	0.88
1:A:429:ASP:HB3	1:A:449:VAL:HG21	1.54	0.87
1:A:699:ARG:HH11	1:A:699:ARG:HG2	1.40	0.86
1:A:419:CYS:HG	4:A:802:BME:HS2	0.96	0.85
1:A:448:LEU:O	1:A:449:VAL:CG2	2.25	0.84
1:A:448:LEU:C	1:A:449:VAL:CG2	2.47	0.83
1:B:447:ILE:HG23	1:B:449:VAL:HG21	1.68	0.74
1:A:699:ARG:NH1	1:A:699:ARG:HG2	1.98	0.74
1:B:458:ARG:NH1	1:B:463:ASP:OD2	2.21	0.73
1:B:449:VAL:HG12	1:B:451:ALA:H	1.54	0.72
1:B:389:ILE:HD11	1:B:445:VAL:HG23	1.74	0.70
1:A:389:ILE:HD11	1:A:445:VAL:HG23	1.73	0.68
1:A:698:SER:O	1:A:700:LEU:N	2.27	0.67
1:B:534:ILE:CD1	1:B:539:TYR:HB2	2.27	0.65
1:A:429:ASP:HB3	1:A:449:VAL:CG2	2.28	0.64
1:A:699:ARG:HH11	1:A:699:ARG:CG	2.12	0.63
1:A:448:LEU:C	1:A:449:VAL:HG22	2.18	0.63
1:A:437:ASP:HA	1:A:440:THR:HG23	1.82	0.62
1:A:564:ARG:HG2	1:A:564:ARG:HH11	1.65	0.59
1:B:447:ILE:HG23	1:B:449:VAL:HG23	1.84	0.59
1:B:564:ARG:HH11	1:B:564:ARG:HG2	1.68	0.59
1:A:419:CYS:SG	4:A:802:BME:S2	2.77	0.58
1:A:389:ILE:HD12	1:A:442:GLU:HB3	1.86	0.58
1:A:437:ASP:HA	1:A:440:THR:CG2	2.34	0.57

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:683:GLN:HE22	2:D:373:THR:HG21	1.71	0.56
1:A:448:LEU:CD1	1:A:448:LEU:N	2.72	0.52
1:A:389:ILE:HD11	1:A:445:VAL:CG2	2.40	0.51
1:A:383:GLY:CA	1:A:450:GLN:OE1	2.54	0.50
1:B:449:VAL:HG12	1:B:450:GLN:N	2.25	0.50
1:A:369:PRO:HG3	4:A:802:BME:H12	1.94	0.49
1:A:604:GLY:HA2	1:A:625:PRO:HG2	1.95	0.49
1:A:369:PRO:CG	4:A:802:BME:H12	2.42	0.49
1:A:608:THR:HB	1:A:623:ALA:HB3	1.94	0.49
1:A:369:PRO:HG3	4:A:802:BME:C1	2.43	0.48
1:B:389:ILE:HD11	1:B:445:VAL:CG2	2.41	0.48
1:B:604:GLY:HA2	1:B:625:PRO:HG2	1.96	0.48
1:A:683:GLN:NE2	2:D:373:THR:HG21	2.28	0.47
1:B:506:THR:HG21	1:B:554:LEU:HD21	1.95	0.47
1:A:371:SER:O	1:A:413:THR:O	2.32	0.47
1:B:534:ILE:HD12	1:B:539:TYR:HB2	1.96	0.47
1:A:602:LYS:O	1:A:672:GLU:HA	2.15	0.47
1:A:491:THR:HG21	1:B:514:GLU:HG3	1.96	0.46
1:B:608:THR:HB	1:B:623:ALA:HB3	1.96	0.46
1:B:671:PRO:HB3	2:D:371:VAL:HG13	1.97	0.46
1:A:389:ILE:CD1	1:A:442:GLU:HB3	2.46	0.46
1:B:602:LYS:O	1:B:672:GLU:HA	2.16	0.46
1:A:699:ARG:HA	1:A:699:ARG:HD3	1.51	0.46
1:A:655:THR:OG1	1:B:736:LEU:O	2.21	0.44
1:A:699:ARG:HH21	1:B:465:LYS:HG3	1.82	0.44
1:B:378:VAL:HG23	1:B:408:VAL:HG12	1.98	0.44
1:B:509:GLU:OE1	1:B:509:GLU:N	2.40	0.43
1:B:680:TRP:CE2	1:B:687:PRO:HB3	2.54	0.43
1:B:582:LEU:HD21	1:B:623:ALA:HB2	2.01	0.42
1:B:596:PHE:CE2	1:B:609:PHE:HB3	2.53	0.42
1:A:430:THR:HA	1:A:448:LEU:HD13	2.01	0.42
1:B:506:THR:CG2	1:B:554:LEU:HD21	2.50	0.42
1:A:582:LEU:HD21	1:A:623:ALA:HB2	2.01	0.42
1:A:596:PHE:CE2	1:A:609:PHE:HB3	2.55	0.42
1:A:702:SER:C	1:A:704:ARG:H	2.24	0.41
1:A:735:ARG:HG2	1:B:735:ARG:HB2	2.01	0.41
1:B:389:ILE:CD1	1:B:445:VAL:HG23	2.48	0.41
1:B:352:GLY:O	1:B:647:GLN:NE2	2.54	0.41
1:A:448:LEU:H	1:A:448:LEU:HD13	1.86	0.41
1:A:698:SER:OG	1:A:699:ARG:N	2.54	0.40
1:A:612:LEU:HG	1:A:614:ILE:HG12	2.03	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:353:ARG:HD2	1:B:498:PHE:CE1	2.56	0.40
1:B:612:LEU:HG	1:B:614:ILE:HG12	2.03	0.40
1:B:447:ILE:HG12	1:B:449:VAL:HG22	2.03	0.40
1:A:389:ILE:CD1	1:A:445:VAL:HG23	2.47	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	408/419 (97%)	398 (98%)	9 (2%)	1 (0%)	47	73
1	B	395/419 (94%)	385 (98%)	9 (2%)	1 (0%)	41	66
2	C	9/11 (82%)	9 (100%)	0	0	100	100
2	D	7/11 (64%)	7 (100%)	0	0	100	100
All	All	819/860 (95%)	799 (98%)	18 (2%)	2 (0%)	47	73

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	449	VAL
1	B	449	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	351/356 (99%)	325 (93%)	26 (7%)	13	32
1	B	339/356 (95%)	324 (96%)	15 (4%)	28	56
2	C	10/10 (100%)	10 (100%)	0	100	100
2	D	8/10 (80%)	8 (100%)	0	100	100
All	All	708/732 (97%)	667 (94%)	41 (6%)	20	43

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	363	ASP
1	A	428	GLU
1	A	435	GLU
1	A	447	ILE
1	A	448	LEU
1	A	449	VAL
1	A	452	ARG
1	A	484	LEU
1	A	498	PHE
1	A	506	THR
1	A	528	SER
1	A	536	GLU
1	A	564	ARG
1	A	599	VAL
1	A	605	LEU
1	A	647	GLN
1	A	661	LEU
1	A	666	LYS
1	A	697	SER
1	A	699	ARG
1	A	704	ARG
1	A	705	SER
1	A	717	GLU
1	A	718	GLN
1	A	722	LEU
1	A	726	GLU
1	B	363	ASP
1	B	484	LEU
1	B	528	SER
1	B	534	ILE
1	B	564	ARG
1	B	599	VAL
1	B	605	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	661	LEU
1	B	666	LYS
1	B	704	ARG
1	B	705	SER
1	B	707	ASN
1	B	715	LYS
1	B	718	GLN
1	B	722	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	394	GLN
1	B	426	GLN
2	D	369	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	BME	A	802	-	3,3,3	0.21	0	1,2,2	0.02	0
3	GOL	B	801	-	5,5,5	0.37	0	5,5,5	0.51	0
3	GOL	A	801	-	5,5,5	0.49	0	5,5,5	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BME	A	802	-	-	0/1/1/1	-
3	GOL	B	801	-	-	0/4/4/4	-
3	GOL	A	801	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	801	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	802	BME	5	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	414/419 (98%)	0.05	10 (2%) 59 60	24, 45, 89, 123	0
1	B	401/419 (95%)	0.23	22 (5%) 25 24	25, 46, 92, 112	0
2	C	11/11 (100%)	0.43	1 (9%) 9 7	39, 46, 58, 66	0
2	D	9/11 (81%)	0.93	0 100 100	60, 66, 85, 93	0
All	All	835/860 (97%)	0.15	33 (3%) 38 37	24, 46, 91, 123	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	381	ALA	4.5
1	B	423	ALA	4.2
1	B	700	LEU	4.0
1	B	450	GLN	3.8
1	B	449	VAL	3.8
1	B	576	MET	3.4
1	B	442	GLU	3.3
1	B	718	GLN	3.2
1	B	386	ASP	3.0
1	A	438	ASP	3.0
1	B	389	ILE	2.9
1	A	451	ALA	2.7
1	B	408	VAL	2.7
1	A	577	GLY	2.7
1	A	448	LEU	2.7
1	B	424	VAL	2.7
2	C	378	ASN	2.7
1	B	428	GLU	2.7
1	B	452	ARG	2.5
1	B	388	PHE	2.5
1	B	385	GLY	2.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	663	CYS	2.4
1	B	378	VAL	2.4
1	A	427	VAL	2.3
1	A	699	ARG	2.3
1	B	380	ALA	2.3
1	B	443	LEU	2.2
1	A	428	GLU	2.1
1	B	447	ILE	2.1
1	A	700	LEU	2.1
1	A	320	PRO	2.0
1	B	453	PRO	2.0
1	B	376	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	B	801	6/6	0.90	0.21	55,56,57,58	0
4	BME	A	802	4/4	0.96	0.18	37,38,38,39	0
3	GOL	A	801	6/6	0.96	0.34	51,52,54,54	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.