



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 08:45 pm BST

PDB ID : 1TLA
Title : HYDROPHOBIC CORE REPACKING AND AROMATIC-AROMATIC INTERACTION IN THE THERMOSTABLE MUTANT OF T4 LYSOZYME SER 117 (RIGHT ARROW) PHE
Authors : Anderson, D.E.; Hurley, J.H.; Nicholson, H.; Baase, W.A.; Matthews, B.W.
Deposited on : 1993-03-22
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	164	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1442 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called T4 LYSOZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	0	0	0
			1297	820	235	237	5			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	54	THR	CYS	CONFLICT	UNP P00720
A	97	ALA	CYS	CONFLICT	UNP P00720
A	117	PHE	SER	CONFLICT	UNP P00720

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total 2	Cl 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	138	Total 138	O 138	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: T4 LYSOZYME



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	61.60 Å 61.60 Å 96.20 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	(Not available) – 2.00 22.21 – 1.62	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.00) 77.2 (22.21-1.62)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.26 (at 1.62 Å)	Xtriage
Refinement program	TNT	Depositor
R, R_{free}	0.167 , (Not available) 0.174 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	12.7	Xtriage
Anisotropy	0.171	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 117.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.052 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1442	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	3.60	8/1318 (0.6%)	1.66	25/1775 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	162	LYS	C-O	97.46	3.08	1.23
1	A	162	LYS	CA-C	59.27	3.07	1.52
1	A	162	LYS	N-CA	52.41	2.51	1.46
1	A	5	GLU	CD-OE2	8.22	1.34	1.25
1	A	11	GLU	CD-OE1	6.26	1.32	1.25
1	A	64	GLU	CD-OE2	5.19	1.31	1.25
1	A	22	GLU	CD-OE1	5.06	1.31	1.25
1	A	108	GLU	CD-OE1	5.05	1.31	1.25

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	162	LYS	CA-C-O	-31.35	54.27	120.10
1	A	162	LYS	N-CA-C	-16.34	66.89	111.00
1	A	70	ASP	CB-CG-OD2	-8.20	110.92	118.30
1	A	70	ASP	CB-CG-OD1	7.61	125.14	118.30
1	A	114	PHE	CB-CG-CD1	7.56	126.09	120.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	96	ARG	NE-CZ-NH2	-7.24	116.68	120.30
1	A	96	ARG	NE-CZ-NH1	7.21	123.91	120.30
1	A	80	ARG	NE-CZ-NH1	7.17	123.89	120.30
1	A	153	PHE	CA-CB-CG	6.95	130.57	113.90
1	A	47	ASP	CB-CG-OD1	6.77	124.39	118.30
1	A	92	ASP	CB-CG-OD2	-6.71	112.26	118.30
1	A	80	ARG	NE-CZ-NH2	-6.38	117.11	120.30
1	A	47	ASP	CB-CG-OD2	-6.12	112.79	118.30
1	A	148	ARG	NE-CZ-NH2	6.09	123.35	120.30
1	A	61	ASP	CB-CG-OD1	-6.08	112.82	118.30
1	A	10	ASP	CB-CG-OD1	6.03	123.72	118.30
1	A	10	ASP	CB-CG-OD2	-5.84	113.04	118.30
1	A	61	ASP	CB-CG-OD2	5.73	123.46	118.30
1	A	25	TYR	CB-CG-CD1	-5.62	117.63	121.00
1	A	119	ARG	NE-CZ-NH1	5.51	123.05	120.30
1	A	72	ASP	CB-CG-OD2	5.46	123.21	118.30
1	A	125	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	A	89	ASP	CB-CG-OD2	5.15	122.94	118.30
1	A	89	ASP	CB-CG-OD1	-5.12	113.69	118.30
1	A	162	LYS	CB-CA-C	5.06	120.53	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	5	GLU	Sidechain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1297	0	1323	37	0
2	A	5	0	0	0	0
3	A	2	0	0	0	0
4	A	138	0	0	13	0
All	All	1442	0	1323	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:PRO:HA	4:A:191:HOH:O	1.77	0.84
1:A:14:ARG:HG3	4:A:298:HOH:O	1.86	0.75
1:A:16:LYS:HE2	1:A:56:GLY:O	1.87	0.74
1:A:162:LYS:N	1:A:162:LYS:CA	2.51	0.73
1:A:133:LEU:HG	1:A:138:TRP:HZ3	1.53	0.71
1:A:154:ARG:HD3	4:A:271:HOH:O	1.91	0.70
1:A:81:ASN:HB2	1:A:108:GLU:OE2	1.92	0.70
1:A:85:LYS:NZ	1:A:89:ASP:OD2	2.28	0.66
1:A:49:ALA:HB2	4:A:282:HOH:O	1.96	0.65
1:A:65:LYS:HG3	4:A:320:HOH:O	1.97	0.63
1:A:117:PHE:CE1	1:A:133:LEU:HD22	2.35	0.62
1:A:14:ARG:NH2	4:A:322:HOH:O	2.30	0.62
1:A:102:MET:HE1	1:A:133:LEU:HD11	1.83	0.60
1:A:128:GLU:HG3	4:A:311:HOH:O	2.02	0.59
1:A:52:ARG:NH2	4:A:317:HOH:O	2.36	0.57
1:A:16:LYS:HD2	1:A:57:VAL:HG22	1.86	0.57
1:A:8:ARG:O	1:A:8:ARG:HG3	2.05	0.57
1:A:83:LYS:NZ	4:A:276:HOH:O	2.38	0.57
1:A:133:LEU:HG	1:A:138:TRP:CZ3	2.39	0.56
1:A:80:ARG:NH1	4:A:289:HOH:O	2.39	0.56
1:A:32:LEU:HB3	4:A:266:HOH:O	2.08	0.54
1:A:16:LYS:HD2	1:A:57:VAL:CG2	2.40	0.52
1:A:80:ARG:NH2	4:A:242:HOH:O	2.44	0.50
1:A:11:GLU:CD	1:A:145:ARG:HH12	2.13	0.50
1:A:119:ARG:O	1:A:123:GLN:HG3	2.11	0.50
1:A:13:LEU:HD12	1:A:29:ILE:HG13	1.94	0.49
1:A:39:LEU:HG	1:A:39:LEU:O	2.13	0.49
1:A:65:LYS:HA	4:A:320:HOH:O	2.13	0.48
1:A:137:ARG:NH1	1:A:141:GLN:HG3	2.29	0.47
1:A:13:LEU:O	1:A:14:ARG:HG2	2.16	0.46
1:A:130:ALA:HB1	1:A:150:ILE:CG2	2.46	0.45
1:A:130:ALA:HB1	1:A:150:ILE:HG23	1.99	0.45
1:A:162:LYS:HE2	1:A:162:LYS:HB2	1.72	0.43
1:A:19:LYS:HA	1:A:24:TYR:O	2.19	0.42
1:A:8:ARG:HH11	1:A:8:ARG:HG3	1.84	0.42
1:A:97:ALA:HA	1:A:100:ILE:HD12	2.01	0.42
1:A:50:ILE:HG22	1:A:52:ARG:HG2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/164 (98%)	155 (97%)	5 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	135/137 (98%)	127 (94%)	8 (6%)	19	15

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	LYS
1	A	20	ASP
1	A	36	SER
1	A	53	ASN
1	A	83	LYS
1	A	127	ASP
1	A	128	GLU
1	A	154	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	140	ASN
1	A	141	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	323	-	4,4,4	1.17	0	6,6,6	1.28	1 (16%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	323	PO4	O4-P-O2	-2.80	98.98	107.97

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers ⓘ

Unable to reproduce the depositors R factor - this section is therefore empty.