



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 06:02 am BST

PDB ID : 4TN3  
Title : Structure of the BBox-Coiled-coil region of Rhesus Trim5alpha  
Authors : Kirkpatrick, J.J.; Stoye, J.P.; Taylor, I.A.; Goldstone, D.C.  
Deposited on : 2014-06-03  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	400	<div> <div>2%</div> <div> <div></div> <div>79%</div> <div>10%</div> <div>10%</div> </div> </div>
1	B	400	<div> <div>6%</div> <div> <div></div> <div>82%</div> <div>9%</div> <div>10%</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 11512 atoms, of which 5712 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRIM5/cyclophilin A fusion protein/T4 Lysozyme chimera.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	359	Total	C	H	N	O	S	0	0	0
			5740	1806	2854	508	554	18			
1	B	362	Total	C	H	N	O	S	0	0	0
			5768	1819	2858	518	557	16			

There are 50 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	68	MET	-	expression tag	UNP G9MAP5
A	69	ALA	-	expression tag	UNP G9MAP5
A	70	HIS	-	expression tag	UNP G9MAP5
A	71	HIS	-	expression tag	UNP G9MAP5
A	72	HIS	-	expression tag	UNP G9MAP5
A	73	HIS	-	expression tag	UNP G9MAP5
A	74	HIS	-	expression tag	UNP G9MAP5
A	75	HIS	-	expression tag	UNP G9MAP5
A	76	SER	-	expression tag	UNP G9MAP5
A	77	ALA	-	expression tag	UNP G9MAP5
A	78	ALA	-	expression tag	UNP G9MAP5
A	79	LEU	-	expression tag	UNP G9MAP5
A	80	GLU	-	expression tag	UNP G9MAP5
A	81	VAL	-	expression tag	UNP G9MAP5
A	82	LEU	-	expression tag	UNP G9MAP5
A	83	PHE	-	expression tag	UNP G9MAP5
A	84	GLN	-	expression tag	UNP G9MAP5
A	85	GLY	-	expression tag	UNP G9MAP5
A	86	PRO	-	expression tag	UNP G9MAP5
A	87	GLY	-	expression tag	UNP G9MAP5
A	315	GLY	ARG	conflict	UNP P00720
A	323	ASN	ASP	conflict	UNP P00720
A	357	THR	CYS	conflict	UNP P00720
A	400	ALA	CYS	conflict	UNP P00720
A	440	ARG	ILE	conflict	UNP P00720

*Continued on next page...*

*Continued from previous page...*



Chain	Residue	Modelled	Actual	Comment	Reference
B	68	MET	-	expression tag	UNP G9MAP5
B	69	ALA	-	expression tag	UNP G9MAP5
B	70	HIS	-	expression tag	UNP G9MAP5
B	71	HIS	-	expression tag	UNP G9MAP5
B	72	HIS	-	expression tag	UNP G9MAP5
B	73	HIS	-	expression tag	UNP G9MAP5
B	74	HIS	-	expression tag	UNP G9MAP5
B	75	HIS	-	expression tag	UNP G9MAP5
B	76	SER	-	expression tag	UNP G9MAP5
B	77	ALA	-	expression tag	UNP G9MAP5
B	78	ALA	-	expression tag	UNP G9MAP5
B	79	LEU	-	expression tag	UNP G9MAP5
B	80	GLU	-	expression tag	UNP G9MAP5
B	81	VAL	-	expression tag	UNP G9MAP5
B	82	LEU	-	expression tag	UNP G9MAP5
B	83	PHE	-	expression tag	UNP G9MAP5
B	84	GLN	-	expression tag	UNP G9MAP5
B	85	GLY	-	expression tag	UNP G9MAP5
B	86	PRO	-	expression tag	UNP G9MAP5
B	87	GLY	-	expression tag	UNP G9MAP5
B	315	GLY	ARG	conflict	UNP P00720
B	323	ASN	ASP	conflict	UNP P00720
B	357	THR	CYS	conflict	UNP P00720
B	400	ALA	CYS	conflict	UNP P00720
B	440	ARG	ILE	conflict	UNP P00720

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total Zn 2 2	0	0
2	A	2	Total Zn 2 2	0	0



- Molecule 1: TRIM5/cyclophilin A fusion protein/T4 Lysozyme chimera



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	124.12Å 59.74Å 146.96Å 90.00° 94.77° 90.00°	Depositor
Resolution (Å)	33.93 – 3.20 33.93 – 3.20	Depositor EDS
% Data completeness (in resolution range)	98.9 (33.93-3.20) 98.9 (33.93-3.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.35 (at 3.18Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, $R_{free}$	0.258 , 0.316 0.269 , 0.325	Depositor DCC
$R_{free}$ test set	903 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	79.7	Xtriage
Anisotropy	0.463	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 76.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	11512	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	102.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/2926	0.41	0/3933
1	B	0.25	0/2955	0.39	0/3974
All	All	0.25	0/5881	0.40	0/7907

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2886	2854	2861	16	0
1	B	2910	2858	2867	14	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
All	All	5800	5712	5728	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 28 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:144:LEU:O	1:B:148:LEU:N	2.30	0.64
1:A:155:GLN:NE2	1:A:155:GLN:O	2.33	0.62
1:B:125:HIS:O	1:B:127:GLY:N	2.33	0.61
1:A:249:LEU:O	1:A:251:GLY:N	2.37	0.58
1:A:423:MET:O	1:A:427:LYS:N	2.38	0.56

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	353/400 (88%)	312 (88%)	35 (10%)	6 (2%)	9	42
1	B	358/400 (90%)	331 (92%)	26 (7%)	1 (0%)	41	74
All	All	711/800 (89%)	643 (90%)	61 (9%)	7 (1%)	15	54

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	250	GLN
1	B	126	ARG
1	A	137	ALA
1	A	241	LEU
1	A	294	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/349 (88%)	295 (96%)	13 (4%)	30	65
1	B	310/349 (89%)	296 (96%)	14 (4%)	27	63
All	All	618/698 (88%)	591 (96%)	27 (4%)	28	64

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	462	ASP
1	B	119	CYS
1	B	262	MET
1	B	93	LYS
1	A	184	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	359/400 (89%)	0.08	7 (1%) 66 53	10, 83, 164, 234	0
1	B	362/400 (90%)	0.27	22 (6%) 21 12	26, 105, 186, 265	0
All	All	721/800 (90%)	0.17	29 (4%) 38 25	10, 93, 179, 265	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	320	ILE	6.9
1	B	343	ASN	4.1
1	B	330	ILE	3.9
1	B	165	ILE	3.6
1	B	322	LYS	3.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	A	701	1/1	0.94	0.09	76,76,76,76	0
2	ZN	B	700	1/1	0.95	0.12	66,66,66,66	0
2	ZN	B	701	1/1	0.98	0.17	57,57,57,57	0
2	ZN	A	702	1/1	0.99	0.12	68,68,68,68	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.