



# Full wwPDB X-ray Structure Validation Report ⓘ

May 18, 2020 – 08:58 am BST

PDB ID : 3TPB  
Title : Structure of HipA(S150A)  
Authors : schumacher, M.A.  
Deposited on : 2011-09-07  
Resolution : 1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	9470 (1.90-1.86)
Clashscore	141614	10282 (1.90-1.86)
Ramachandran outliers	138981	10152 (1.90-1.86)
Sidechain outliers	138945	10152 (1.90-1.86)
RSRZ outliers	127900	9303 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	<div> <div>3%</div> <div> <div></div> <div>72%</div> <div>18%</div> <div>•</div> <div>9%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	A	834	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3434 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase HipA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	401	Total	C	N	O	S	0	0	0
			3178	2040	554	572	12			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	150	ALA	SER	ENGINEERED MUTATION	UNP P23874
A	243	ARG	GLU	CONFLICT	UNP P23874
A	309	GLN	ASP	CONFLICT	UNP P23874

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	245	Total	O	0	0
			245	245		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein kinase HipA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.43 Å   84.50 Å   50.00 Å 90.00°   89.94°   90.00°	Depositor
Resolution (Å)	50.00 – 1.88 50.00 – 1.88	Depositor EDS
% Data completeness (in resolution range)	89.4 (50.00-1.88) 89.6 (50.00-1.88)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.38 (at 1.88 Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.192 , 0.243 0.191 , 0.240	Depositor DCC
$R_{free}$ test set	2929 reflections (9.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.0	Xtriage
Anisotropy	0.649	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 53.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.005 for -l,k,h 0.035 for h,-k,-l 0.021 for -l,-k,-h	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3434	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.44% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.56	7/3248 (0.2%)	0.67	0/4398

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	388	MET	CG-SD	6.05	1.96	1.81
1	A	283	MET	CG-SD	5.96	1.96	1.81
1	A	279	MET	CG-SD	5.92	1.96	1.81
1	A	296	MET	CG-SD	5.90	1.96	1.81
1	A	255	MET	CG-SD	5.43	1.95	1.81
1	A	8	MET	CG-SD	5.13	1.94	1.81
1	A	354	MET	CG-SD	5.04	1.94	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3178	0	3226	62	0
2	A	1	0	0	1	0
3	A	10	0	0	0	0
4	A	245	0	0	11	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3434	0	3226	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:389:HIS:HD1	1:A:437:TYR:HE2	1.25	0.85
1:A:225:ALA:O	1:A:228:VAL:HG12	1.81	0.79
1:A:3:LYS:HG2	1:A:17:THR:HG22	1.65	0.77
1:A:124:ARG:HG3	1:A:124:ARG:HH11	1.48	0.77
1:A:306:GLY:HA2	1:A:366:ILE:HD13	1.71	0.72
1:A:416:GLU:HG3	4:A:620:HOH:O	1.90	0.71
1:A:363:LYS:HG3	1:A:373:HIS:CD2	2.24	0.71
1:A:283:MET:HE3	1:A:283:MET:HA	1.72	0.71
1:A:17:THR:HG23	4:A:516:HOH:O	1.97	0.65
1:A:124:ARG:NH1	1:A:124:ARG:HG3	2.13	0.64
1:A:49:ARG:O	1:A:49:ARG:HD3	1.98	0.64
1:A:336:ALA:O	1:A:339:VAL:HG22	1.97	0.64
1:A:283:MET:CE	1:A:283:MET:HA	2.27	0.63
1:A:27:LYS:HE3	1:A:49:ARG:NH1	2.18	0.58
1:A:224:LYS:HE2	1:A:229:ARG:NH1	2.19	0.58
1:A:224:LYS:HE2	1:A:229:ARG:HH12	1.69	0.57
1:A:363:LYS:HE2	1:A:373:HIS:NE2	2.20	0.57
1:A:342:GLY:C	1:A:344:GLY:H	2.08	0.57
1:A:297:LYS:HE3	4:A:616:HOH:O	2.07	0.55
1:A:21:ASN:ND2	1:A:23:ALA:H	2.06	0.54
1:A:194:LEU:HD12	1:A:194:LEU:N	2.23	0.53
1:A:309:GLN:HB2	4:A:506:HOH:O	2.08	0.53
1:A:363:LYS:HD3	2:A:441:CL:CL	2.46	0.53
1:A:83:SER:OG	1:A:85:GLN:HB2	2.09	0.53
1:A:407:LYS:HE3	1:A:423:GLU:OE1	2.09	0.52
1:A:196:LEU:HB2	4:A:629:HOH:O	2.09	0.52
1:A:221:GLU:HG2	4:A:608:HOH:O	2.09	0.51
1:A:385:GLU:HG2	1:A:389:HIS:HD2	1.76	0.51
1:A:119:LYS:HD3	1:A:120:LEU:N	2.26	0.51
1:A:74:ARG:NH2	1:A:156:GLU:HG2	2.27	0.50
1:A:342:GLY:O	1:A:344:GLY:N	2.44	0.50
1:A:126:GLU:O	1:A:130:THR:HG23	2.11	0.50
1:A:433:LEU:O	1:A:437:TYR:HB2	2.12	0.50

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:30:PRO:HB3	1:A:49:ARG:HH21	1.76	0.50
1:A:41:LEU:HD11	1:A:101:VAL:HG12	1.95	0.49
1:A:419:VAL:O	1:A:423:GLU:HB2	2.13	0.48
1:A:375:LEU:HD11	1:A:437:TYR:OH	2.15	0.47
1:A:235:ARG:HB3	1:A:237:ASP:OD1	2.16	0.46
1:A:2:PRO:HB2	1:A:18:LYS:O	2.16	0.46
1:A:162:ARG:HG2	1:A:167:TRP:CZ3	2.51	0.45
1:A:157:LYS:HG2	4:A:592:HOH:O	2.17	0.45
1:A:43:LEU:HD12	1:A:258:THR:OG1	2.16	0.45
1:A:21:ASN:HD22	1:A:21:ASN:N	2.14	0.45
1:A:73:ASP:OD1	1:A:84:ARG:HD2	2.16	0.44
1:A:156:GLU:OE1	1:A:156:GLU:HA	2.18	0.44
1:A:42:SER:HB2	1:A:60:PHE:CE1	2.53	0.44
1:A:309:GLN:NE2	4:A:682:HOH:O	2.51	0.43
1:A:101:VAL:O	1:A:101:VAL:HG23	2.19	0.42
1:A:237:ASP:HB3	1:A:252:GLN:OE1	2.19	0.42
1:A:277:ARG:NH2	4:A:519:HOH:O	2.51	0.42
1:A:68:SER:HA	1:A:69:PRO:HD3	1.93	0.42
1:A:342:GLY:C	1:A:344:GLY:N	2.72	0.42
1:A:348:SER:O	1:A:351:LYS:NZ	2.53	0.41
1:A:42:SER:HB2	1:A:60:PHE:CD1	2.55	0.41
1:A:106:GLU:CD	1:A:107:ASP:N	2.74	0.41
1:A:350:LEU:C	1:A:351:LYS:HD2	2.41	0.41
1:A:113:PRO:HB2	1:A:173:ILE:HD11	2.03	0.41
1:A:290:LYS:HE3	4:A:548:HOH:O	2.21	0.41
1:A:394:ASP:OD2	1:A:397:ARG:NH2	2.54	0.41
1:A:90:LEU:HD23	1:A:90:LEU:HA	1.96	0.41
1:A:119:LYS:C	1:A:119:LYS:HD3	2.41	0.40
1:A:120:LEU:HA	4:A:684:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	393/440 (89%)	383 (98%)	7 (2%)	3 (1%)	19	9

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	436	GLU
1	A	342	GLY
1	A	343	THR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/373 (92%)	329 (96%)	13 (4%)	33	21

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	21	ASN
1	A	49	ARG
1	A	51	ASN
1	A	85	GLN
1	A	106	GLU
1	A	124	ARG
1	A	162	ARG
1	A	171	LYS
1	A	175	PRO
1	A	283	MET
1	A	309	GLN
1	A	362	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	ASN
1	A	21	ASN
1	A	51	ASN
1	A	216	ASN
1	A	320	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PO4	A	833	-	4,4,4	1.75	1 (25%)	6,6,6	0.46	0
3	PO4	A	834	-	4,4,4	1.55	0	6,6,6	0.44	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	833	PO4	P-O4	-2.29	1.47	1.54

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	401/440 (91%)	0.07	15 (3%) 41 43	14, 27, 53, 66	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	343	THR	9.2
1	A	154	ALA	6.5
1	A	437	TYR	5.2
1	A	2	PRO	4.9
1	A	342	GLY	4.4
1	A	80	HIS	4.2
1	A	436	GLU	3.9
1	A	341	GLY	3.2
1	A	106	GLU	2.9
1	A	107	ASP	2.8
1	A	195	ASP	2.6
1	A	49	ARG	2.3
1	A	155	GLN	2.1
1	A	156	GLU	2.1
1	A	10	ASN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PO4	A	834	5/5	0.63	0.64	77,78,80,81	0
3	PO4	A	833	5/5	0.97	0.11	32,41,43,44	0
2	CL	A	441	1/1	0.99	0.10	20,20,20,20	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.