



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 10:56 pm BST

PDB ID : 3TPP  
Title : Crystal structure of BACE1 complexed with an inhibitor  
Authors : Xu, Y.C.; Li, M.J.; Greenblatt, H.; Chen, T.T.; Silman, I.; Sussman, J.L.  
Deposited on : 2011-09-08  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*


The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	URE	A	412	-	X	-	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3427 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-secretase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	375	2993	1930	487	561	15	0	17	0

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-39	MET	-	EXPRESSION TAG	UNP P56817
A	-38	GLY	-	EXPRESSION TAG	UNP P56817
A	-37	SER	-	EXPRESSION TAG	UNP P56817
A	-36	SER	-	EXPRESSION TAG	UNP P56817
A	-35	HIS	-	EXPRESSION TAG	UNP P56817
A	-34	HIS	-	EXPRESSION TAG	UNP P56817
A	-33	HIS	-	EXPRESSION TAG	UNP P56817
A	-32	HIS	-	EXPRESSION TAG	UNP P56817
A	-31	HIS	-	EXPRESSION TAG	UNP P56817
A	-30	HIS	-	EXPRESSION TAG	UNP P56817
A	-29	SER	-	EXPRESSION TAG	UNP P56817
A	-28	ALA	-	EXPRESSION TAG	UNP P56817
A	-27	GLY	-	EXPRESSION TAG	UNP P56817
A	-26	GLU	-	EXPRESSION TAG	UNP P56817
A	-25	ASN	-	EXPRESSION TAG	UNP P56817
A	-24	LEU	-	EXPRESSION TAG	UNP P56817
A	-23	TYR	-	EXPRESSION TAG	UNP P56817
A	-22	PHE	-	EXPRESSION TAG	UNP P56817
A	-21	GLN	-	EXPRESSION TAG	UNP P56817
A	-20	GLY	-	EXPRESSION TAG	UNP P56817
A	-19	THR	-	EXPRESSION TAG	UNP P56817
A	75	ALA	LYS	ENGINEERED MUTATION	UNP P56817
A	77	ALA	GLU	ENGINEERED MUTATION	UNP P56817

- Molecule 2 is N-[(1S,2R)-1-BENZYL-3-(CYCLOPROPYLAMINO)-2-HYDROXYPROPYL]-5-[METHYL(METHYLSULFONYL)AMINO]-N'-[(1R)-1-PHENYLETHYL]ISOPHTHALAMIDE (three-letter code: 5HA) (formula: C<sub>31</sub>H<sub>38</sub>N<sub>4</sub>O<sub>5</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			41	31	4	5	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	8	Total	Cl	0	0
			8	8		

- Molecule 4 is UREA (three-letter code: URE) (formula: CH<sub>4</sub>N<sub>2</sub>O).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 4 1 2 1	0	0
4	A	1	Total C N O 4 1 2 1	0	0
4	A	1	Total C N O 4 1 2 1	0	0
4	A	1	Total C N O 4 1 2 1	0	0
4	A	1	Total C N O 4 1 2 1	0	0
4	A	1	Total C N O 4 1 2 1	0	0
4	A	1	Total C N O 4 1 2 1	0	0
4	A	1	Total C N O 4 1 2 1	0	0
4	A	1	Total C N O 4 1 2 1	0	0
4	A	1	Total C N O 4 1 2 1	0	0
4	A	1	Total C N O 4 1 2 1	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	331	Total	O	0	0
			331	331		



- Molecule 1: Beta-secretase 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.55Å 128.20Å 76.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.17 – 1.60 49.14 – 1.60	Depositor EDS
% Data completeness (in resolution range)	98.7 (43.17-1.60) 98.7 (49.14-1.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.40 (at 1.60Å)	Xtriage
Refinement program	PHENIX 1.7_650	Depositor
R, $R_{free}$	0.154 , 0.180 0.151 , 0.178	Depositor DCC
$R_{free}$ test set	3408 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	12.7	Xtriage
Anisotropy	0.113	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 50.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3427	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.90% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: URE, 5HA, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.31	5/3116 (0.2%)	1.25	10/4237 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	195	ARG	CG-CD	-8.17	1.31	1.51
1	A	104	GLU	CD-OE2	7.72	1.34	1.25
1	A	296	PHE	CE2-CZ	5.45	1.47	1.37
1	A	319	CYS	CB-SG	5.12	1.91	1.82
1	A	286	TYR	CD1-CE1	5.02	1.46	1.39

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	195	ARG	NE-CZ-NH1	-12.89	113.85	120.30
1	A	130	ASP	CB-CG-OD2	7.72	125.25	118.30
1	A	54	ARG	NE-CZ-NH1	-6.83	116.89	120.30
1	A	366	ARG	NE-CZ-NH2	-6.58	117.01	120.30
1	A	61	ARG	NE-CZ-NH2	-6.58	117.01	120.30
1	A	108	PHE	CB-CG-CD1	-6.40	116.32	120.80
1	A	198	TYR	CD1-CE1-CZ	-5.83	114.56	119.80
1	A	223	ASP	CB-CG-OD1	5.23	123.01	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	216	ASP	CB-CG-OD2	5.04	122.83	118.30
1	A	205	ARG	NE-CZ-NH2	-5.03	117.79	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	197	TRP	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2993	0	2907	22	0
2	A	41	0	38	0	0
3	A	8	0	0	0	0
4	A	44	0	44	1	0
5	A	10	0	0	0	0
6	A	331	0	0	7	1
All	All	3427	0	2989	22	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:27:LEU:HD13	1:A:40[B]:VAL:HG21	1.40	1.00
1:A:45:HIS:HD2	1:A:47:PHE:H	1.13	0.96
1:A:27:LEU:CD1	1:A:40[B]:VAL:HG21	1.96	0.95
1:A:27:LEU:HD13	1:A:40[B]:VAL:CG2	2.09	0.83
1:A:45:HIS:CD2	1:A:47:PHE:H	2.01	0.77
1:A:128:ARG:NE	6:A:724:HOH:O	2.23	0.71
1:A:145[B]:HIS:HE1	6:A:636:HOH:O	1.76	0.67
1:A:314:THR:HG22	1:A:316:GLN:H	1.62	0.64

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:8:GLY:HA3	1:A:170:VAL:HG12	1.80	0.62
1:A:314:THR:HB	1:A:317:ASP:OD1	1.99	0.62
1:A:314:THR:CG2	1:A:316:GLN:H	2.13	0.61
1:A:142[B]:LYS:CG	6:A:691:HOH:O	2.59	0.51
1:A:145[B]:HIS:CD2	6:A:630:HOH:O	2.63	0.50
1:A:314:THR:HG22	1:A:316:GLN:N	2.26	0.50
1:A:17:GLU:OE1	6:A:676:HOH:O	2.20	0.49
1:A:8:GLY:CA	1:A:170:VAL:HG12	2.42	0.48
1:A:8:GLY:HA3	1:A:170:VAL:CG1	2.45	0.47
1:A:20[B]:VAL:HG12	1:A:85:VAL:HG22	1.98	0.45
1:A:145[B]:HIS:CE1	6:A:636:HOH:O	2.60	0.45
1:A:45:HIS:HE1	4:A:405:URE:N2	2.16	0.42
1:A:222:TYR:HA	1:A:223:ASP:HA	1.66	0.41
1:A:145[B]:HIS:HD2	6:A:630:HOH:O	2.01	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:731:HOH:O	6:A:731:HOH:O[3_655]	2.02	0.18

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/433 (89%)	381 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	322/368 (88%)	320 (99%)	2 (1%)	86	77

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	111	ASN
1	A	197	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	ASN
1	A	45	HIS
1	A	111	ASN
1	A	181	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 8 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	URE	A	412	-	3,3,3	1.75	1 (33%)	3,3,3	3.88	2 (66%)
4	URE	A	402	-	3,3,3	0.65	0	3,3,3	1.90	1 (33%)
4	URE	A	405	-	3,3,3	0.84	0	3,3,3	1.63	1 (33%)
4	URE	A	409	-	3,3,3	0.63	0	3,3,3	2.41	2 (66%)
4	URE	A	410	-	3,3,3	0.44	0	3,3,3	1.31	0
4	URE	A	404	-	3,3,3	1.87	1 (33%)	3,3,3	2.12	1 (33%)
4	URE	A	407	-	3,3,3	0.42	0	3,3,3	1.02	0
4	URE	A	408	-	3,3,3	0.84	0	3,3,3	2.11	2 (66%)
4	URE	A	403	-	3,3,3	0.52	0	3,3,3	1.16	0
5	SO4	A	413	-	4,4,4	0.58	0	6,6,6	0.42	0
4	URE	A	411	-	3,3,3	0.32	0	3,3,3	2.43	1 (33%)
2	5HA	A	999	-	43,44,44	1.46	8 (18%)	58,62,62	1.91	16 (27%)
4	URE	A	406	-	3,3,3	0.26	0	3,3,3	0.45	0
5	SO4	A	414	-	4,4,4	0.26	0	6,6,6	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5HA	A	999	-	-	5/41/45/45	0/4/4/4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	999	5HA	C14-C9	3.61	1.44	1.39
2	A	999	5HA	C3-C1	3.45	1.61	1.48
4	A	404	URE	O-C	-3.15	1.14	1.26
2	A	999	5HA	C8-C6	3.11	1.60	1.53
2	A	999	5HA	C15-N4	2.60	1.39	1.34
2	A	999	5HA	C12-C11	2.50	1.43	1.39
4	A	412	URE	C-N2	2.31	1.40	1.34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	999	5HA	O6-S1	2.13	1.46	1.43
2	A	999	5HA	C27-C28	2.12	1.43	1.38
2	A	999	5HA	C31-C30	2.03	1.42	1.39

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	999	5HA	C3-C2-C1	5.77	65.62	60.33
4	A	412	URE	N2-C-N1	5.57	128.26	117.82
2	A	999	5HA	C5-C6-N2	4.37	118.34	109.92
2	A	999	5HA	C3-C2-N1	-4.20	112.59	118.61
2	A	999	5HA	C8-C6-N2	-3.90	104.38	110.07
4	A	412	URE	O-C-N1	-3.62	112.83	121.02
4	A	411	URE	N2-C-N1	-3.60	111.06	117.82
2	A	999	5HA	C11-C15-N4	-3.44	110.47	117.06
2	A	999	5HA	C3-C1-C2	-3.20	57.15	59.84
4	A	409	URE	N2-C-N1	-3.17	111.87	117.82
4	A	404	URE	N2-C-N1	3.16	123.74	117.82
2	A	999	5HA	C1-C3-C2	-3.10	57.23	59.84
4	A	409	URE	O-C-N2	2.72	127.18	121.02
4	A	408	URE	O-C-N2	-2.68	114.96	121.02
4	A	402	URE	O-C-N1	-2.58	115.20	121.02
2	A	999	5HA	O2-C7-C9	2.56	125.50	120.94
2	A	999	5HA	C1-C2-N1	-2.42	115.14	118.61
2	A	999	5HA	C14-C13-N3	-2.38	114.27	119.59
4	A	408	URE	N2-C-N1	2.37	122.26	117.82
2	A	999	5HA	C11-C10-C9	-2.36	117.69	120.44
2	A	999	5HA	C33-C32-C31	2.30	123.69	120.19
4	A	405	URE	O-C-N1	2.23	126.07	121.02
2	A	999	5HA	C18-C17-C30	-2.21	107.21	112.25
2	A	999	5HA	O4-C15-C11	2.16	124.79	120.94
2	A	999	5HA	C32-C31-C30	-2.15	117.98	120.65
2	A	999	5HA	C34-C33-C32	-2.04	116.14	119.93

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	999	5HA	C12-C13-N3-S1
2	A	999	5HA	C14-C13-N3-S1
2	A	999	5HA	C24-C29-C8-C6
2	A	999	5HA	C13-N3-S1-O5

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	A	999	5HA	C28-C29-C8-C6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	405	URE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	375/433 (86%)	-0.28	11 (2%) 51 49	7, 11, 29, 46	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	365	PHE	6.1
1	A	254	THR	5.0
1	A	170	VAL	4.1
1	A	314	THR	3.7
1	A	-4	ARG	3.5
1	A	253	SER	3.4
1	A	315	SER	2.7
1	A	273	GLY	2.5
1	A	385	ASN	2.4
1	A	10	SER	2.2
1	A	272	ALA	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	URE	A	406	4/4	0.81	0.17	15,19,28,29	0
4	URE	A	408	4/4	0.83	0.17	15,20,26,26	0
3	CL	A	401	1/1	0.88	0.07	45,45,45,45	0
4	URE	A	409	4/4	0.89	0.10	25,26,26,30	0
4	URE	A	412	4/4	0.90	0.17	19,23,28,29	0
4	URE	A	410	4/4	0.90	0.29	26,30,31,35	0
4	URE	A	411	4/4	0.91	0.32	30,30,32,33	0
3	CL	A	400	1/1	0.93	0.10	31,31,31,31	0
4	URE	A	407	4/4	0.94	0.09	16,18,19,23	0
3	CL	A	399	1/1	0.94	0.06	44,44,44,44	0
4	URE	A	405	4/4	0.95	0.08	13,15,19,23	0
3	CL	A	397	1/1	0.95	0.04	45,45,45,45	0
4	URE	A	404	4/4	0.95	0.09	10,17,17,18	0
5	SO4	A	414	5/5	0.96	0.12	30,39,43,44	0
4	URE	A	402	4/4	0.96	0.06	10,12,12,12	0
3	CL	A	395	1/1	0.97	0.09	37,37,37,37	0
3	CL	A	398	1/1	0.97	0.05	30,30,30,30	0
4	URE	A	403	4/4	0.97	0.06	13,13,13,14	0
5	SO4	A	413	5/5	0.97	0.13	12,20,26,30	5
3	CL	A	396	1/1	0.99	0.06	20,20,20,20	0
2	5HA	A	999	41/41	0.99	0.08	6,9,12,14	0
3	CL	A	394	1/1	1.00	0.09	7,7,7,7	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 5HA A 999:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.