



wwPDB EM Validation Summary Report ⓘ

Nov 20, 2022 – 01:36 AM EST

PDB ID : 7TRJ
EMDB ID : EMD-26098
Title : The eukaryotic translation initiation factor 2B from Homo sapiens with a H160D mutation in the beta subunit
Authors : Wang, L.; Schoof, M.; Lawrence, R.; Boone, M.; Frost, A.; Walter, P.
Deposited on : 2022-01-29
Resolution : 2.80 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev43
MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	FAILED
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	721	44% 12% • 43%
1	B	721	44% 12% • 43%
2	C	351	70% 22% • 6%
2	D	351	69% 24% • 6%
3	E	523	52% 14% • 33%
3	F	523	53% 14% • 33%
4	G	305	66% 24% • 7%
4	H	305	66% 25% • 7%
5	I	452	38% 13% • 48%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	J	452	<div> <div></div> <div>37%</div> <div>13%</div> <div>48%</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 25190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Translation initiation factor eIF-2B subunit epsilon.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	414	Total	C	N	O	S	0	0
			3210	2030	570	595	15		
1	B	414	Total	C	N	O	S	0	0
			3210	2030	570	595	15		

- Molecule 2 is a protein called Translation initiation factor eIF-2B subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	D	329	Total	C	N	O	S	0	0
			2571	1629	450	477	15		
2	C	329	Total	C	N	O	S	0	0
			2571	1629	450	477	15		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	160	ASP	HIS	engineered mutation	UNP P49770
C	160	ASP	HIS	engineered mutation	UNP P49770

- Molecule 3 is a protein called Translation initiation factor eIF-2B subunit delta.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	E	353	Total	C	N	O	S	0	0
			2750	1741	490	505	14		
3	F	353	Total	C	N	O	S	0	0
			2750	1741	490	505	14		

- Molecule 4 is a protein called Translation initiation factor eIF-2B subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	G	285	Total	C	N	O	S	0	0
			2203	1419	364	409	11		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
4	H	285	Total	C	N	O	S	0	0
			2203	1419	364	409	11		


- Molecule 5 is a protein called Translation initiation factor eIF-2B subunit gamma.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	I	236	Total	C	N	O	S	0	0
			1861	1212	312	321	16		
5	J	236	Total	C	N	O	S	0	0
			1861	1212	312	321	16		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Translation initiation factor eIF-2B subunit epsilon




- Molecule 1: Translation initiation factor eIF-2B subunit epsilon






- Molecule 2: Translation initiation factor eIF-2B subunit beta





- Molecule 2: Translation initiation factor eIF-2B subunit beta



- Molecule 3: Translation initiation factor eIF-2B subunit delta





ASP	SER	LEU	GLY	PRO	GLU	THR	GLN	ILE	GLY	GLU	LYS	SER	SER	ILE	ARG	LYS	ARG	SER	VAL	ILE	GLY	SER	SER	CYS	LEU	ILE	LYS	ASP	ARG	VAL	THR	ILE	THR	ASN	CYS	LEU	LEU	MET	ASN	SER	SER	VAL	THR	VAL	GLU	GLU	GLY	SER	SER	VAL	VAL	ILE	ILE	GLN	GLY	SER	ASN	ASN	ALA
VAL	ASP	GLU	LYS	ALA	ASP	ILE	LYS	ASP	CYS	LEU	ILE	GLY	SER	GLY	GLN	ARG	ILE	GLU	ALA	LYS	ALA	LYS	ARG	VAL	ASN	GLU	VAL	ILE	VAL	VAL	GLY	ASN	ASP	GLN	LEU	MET	GLU	ILE																					

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	170244	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	67	Depositor
Minimum defocus (nm)	0.6	Depositor
Maximum defocus (nm)	2.0	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/3277	0.52	2/4465 (0.0%)
1	B	0.26	0/3277	0.52	2/4465 (0.0%)
2	C	0.27	0/2619	0.50	0/3540
2	D	0.27	0/2619	0.50	0/3540
3	E	0.26	0/2802	0.46	0/3808
3	F	0.26	0/2802	0.46	0/3808
4	G	0.27	0/2239	0.52	1/3027 (0.0%)
4	H	0.27	0/2239	0.52	0/3027
5	I	0.26	0/1890	0.59	1/2546 (0.0%)
5	J	0.27	0/1890	0.59	1/2546 (0.0%)
All	All	0.26	0/25654	0.52	7/34772 (0.0%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	J	328	LYS	CA-CB-CG	5.35	125.16	113.40
1	B	396	LEU	CA-CB-CG	5.34	127.59	115.30
5	I	328	LYS	CA-CB-CG	5.34	125.15	113.40
1	A	396	LEU	CA-CB-CG	5.32	127.52	115.30
1	B	127	LEU	CA-CB-CG	5.32	127.52	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3210	0	3160	54	0
1	B	3210	0	3160	51	0
2	C	2571	0	2581	54	0
2	D	2571	0	2581	54	0
3	E	2750	0	2816	50	0
3	F	2750	0	2816	49	0
4	G	2203	0	2271	56	0
4	H	2203	0	2271	54	0
5	I	1861	0	1924	42	0
5	J	1861	0	1924	41	0
All	All	25190	0	25504	458	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 458 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:271:ASP:OD1	1:A:274:ARG:NH2	2.08	0.85
1:B:271:ASP:OD1	1:B:274:ARG:NH2	2.08	0.85
4:H:303:LEU:HD23	4:H:305:LEU:HD22	1.61	0.82
4:G:303:LEU:HD23	4:G:305:LEU:HD22	1.61	0.82
4:G:235:PHE:HB3	4:G:300:LEU:HD21	1.66	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	410/721 (57%)	388 (95%)	21 (5%)	1 (0%)	47 78
1	B	410/721 (57%)	389 (95%)	20 (5%)	1 (0%)	47 78

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	323/351 (92%)	309 (96%)	14 (4%)	0	100	100
2	D	323/351 (92%)	309 (96%)	14 (4%)	0	100	100
3	E	351/523 (67%)	348 (99%)	3 (1%)	0	100	100
3	F	351/523 (67%)	348 (99%)	3 (1%)	0	100	100
4	G	281/305 (92%)	272 (97%)	9 (3%)	0	100	100
4	H	281/305 (92%)	272 (97%)	9 (3%)	0	100	100
5	I	222/452 (49%)	210 (95%)	11 (5%)	1 (0%)	29	61
5	J	222/452 (49%)	210 (95%)	11 (5%)	1 (0%)	29	61
All	All	3174/4704 (68%)	3055 (96%)	115 (4%)	4 (0%)	54	81

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	I	310	CYS
5	J	310	CYS
1	A	52	PHE
1	B	52	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	355/626 (57%)	326 (92%)	29 (8%)	11	33
1	B	355/626 (57%)	326 (92%)	29 (8%)	11	33
2	C	279/298 (94%)	269 (96%)	10 (4%)	35	69
2	D	279/298 (94%)	269 (96%)	10 (4%)	35	69
3	E	309/444 (70%)	290 (94%)	19 (6%)	18	48
3	F	309/444 (70%)	290 (94%)	19 (6%)	18	48
4	G	241/260 (93%)	212 (88%)	29 (12%)	5	15
4	H	241/260 (93%)	212 (88%)	29 (12%)	5	15

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
5	I	203/398 (51%)	187 (92%)	16 (8%)	12	34
5	J	203/398 (51%)	187 (92%)	16 (8%)	12	34
All	All	2774/4052 (68%)	2568 (93%)	206 (7%)	17	37

5 of 206 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	250	SER
3	F	169	THR
5	J	220	SER
1	B	328	THR
1	B	447	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 46 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	371	ASN
2	C	331	ASN
2	C	49	HIS
2	C	162	ASN
3	F	213	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Map visualisation

This section contains visualisations of the EMDB entry EMD-26098. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections

This section was not generated.

6.2 Central slices

This section was not generated.

6.3 Largest variance slices

This section was not generated.

6.4 Orthogonal surface views

This section was not generated.

6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis ⓘ

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution ⓘ

This section was not generated.

7.2 Volume estimate versus contour level ⓘ

This section was not generated.

7.3 Rotationally averaged power spectrum ⓘ

This section was not generated. The rotationally averaged power spectrum had issues being displayed.

8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit

This section was not generated.