



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 5, 2022 – 04:16 PM EDT

PDB ID : 7TSC  
Title : Structure of rat neuronal nitric oxide synthase R349A mutant heme domain in complex with 4-methyl-6-(3-(4-methylpiperazin-1-yl)propyl)pyridin-2-amine  
Authors : Li, H.; Poulos, T.L.  
Deposited on : 2022-01-31  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.29
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.29


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div>27%</div> <div>80%</div> <div>17%</div> <div>..</div> </div>

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 3650 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	411	Total	C	N	O	S	0	1	0
			3343	2140	571	611	21			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	349	ALA	ARG	engineered mutation	UNP P29476

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula:  $C_9H_{15}N_5O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is 4-methyl-6-[3-(4-methylpiperazin-1-yl)propyl]pyridin-2-amine (three-letter code: KMR) (formula:  $C_{14}H_{24}N_4$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N		0	0
			18	14	4			

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	224	Total	O	0	0
			224	224		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.90Å 114.29Å 164.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.53 – 1.80 39.52 – 1.80	Depositor EDS
% Data completeness (in resolution range)	94.0 (39.53-1.80) 96.0 (39.52-1.80)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.04 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.11.1 _2575	Depositor
R, $R_{free}$	0.211 , 0.253 0.214 , 0.253	Depositor DCC
$R_{free}$ test set	2104 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.7	Xtriage
Anisotropy	1.247	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 73.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3650	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	84.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN, H4B, ACT, KMR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.33	0/3439	0.49	0/4665

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3343	0	3258	39	0
2	A	43	0	30	2	0
3	A	17	0	15	1	0
4	A	18	0	0	0	0
5	A	4	0	3	0	0
6	A	1	0	0	0	0
7	A	224	0	0	1	0
All	All	3650	0	3306	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:716:TRP:O	1:A:717:LYS:HB2	1.99	0.61
2:A:801:HEM:HBB2	2:A:801:HEM:HHC	1.82	0.60
2:A:801:HEM:HMC2	2:A:801:HEM:HBC2	1.84	0.59
1:A:359:LYS:HE2	1:A:381:LEU:HD21	1.85	0.59
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.87	0.57
1:A:475:TRP:HB2	1:A:523:LEU:HB3	1.88	0.54
1:A:351:LYS:HD3	1:A:351:LYS:H	1.74	0.52
1:A:361:PHE:O	1:A:364:GLN:HG2	2.10	0.52
1:A:409:TRP:CE3	1:A:421:TRP:HA	2.45	0.51
1:A:359:LYS:HG3	1:A:381:LEU:HD11	1.93	0.50
1:A:525:GLN:HG3	1:A:529:ASN:O	2.12	0.49
1:A:590:GLY:HA2	1:A:648:ILE:HD11	1.95	0.48
1:A:321:THR:HG23	1:A:322:LEU:H	1.79	0.48
1:A:467:ASP:OD1	1:A:468:GLY:N	2.46	0.48
1:A:425:GLN:OE1	7:A:901:HOH:O	2.20	0.48
1:A:496:PRO:HA	1:A:499:VAL:HG23	1.96	0.48
1:A:485:TYR:CZ	1:A:514:ARG:HA	2.49	0.47
1:A:464:GLN:HB3	1:A:579:PHE:CE2	2.49	0.47
1:A:351:LYS:HD2	1:A:392:SER:HB2	1.98	0.46
1:A:548:HIS:CG	1:A:549:PRO:HD2	2.52	0.45
1:A:610:VAL:HG21	1:A:633:ILE:HD11	1.98	0.45
1:A:350:THR:HG22	1:A:351:LYS:H	1.83	0.44
1:A:523:LEU:HD23	1:A:523:LEU:HA	1.84	0.44
1:A:478:GLN:HB2	1:A:481:ARG:CG	2.48	0.44
1:A:716:TRP:HB3	1:A:717:LYS:H	1.43	0.44
1:A:469:LYS:HE3	1:A:470:HIS:CE1	2.53	0.43
1:A:478:GLN:HB2	1:A:481:ARG:HG2	2.00	0.43
1:A:510:TRP:CD1	1:A:521:PRO:HG3	2.54	0.42
1:A:513:PRO:HG2	1:A:518:ASP:OD2	2.19	0.42
1:A:595:VAL:O	1:A:599:CYS:HB2	2.20	0.42
1:A:375:LYS:HG3	1:A:379:ASP:OD1	2.20	0.42
1:A:355:PHE:N	1:A:356:PRO:HD2	2.35	0.41
1:A:480:ILE:HD13	1:A:541:VAL:HG13	2.02	0.41
1:A:612:LYS:HE2	1:A:612:LYS:HB3	1.68	0.41
1:A:336:MET:HE2	3:A:802:H4B:H9	2.01	0.41
1:A:445:HIS:CD2	1:A:445:HIS:C	2.94	0.41
1:A:525:GLN:HE21	1:A:528:GLY:HA2	1.85	0.41
1:A:666:TYR:CD2	1:A:671:GLY:HA2	2.56	0.41
1:A:537:PRO:HA	1:A:538:PRO:HD3	1.97	0.40
1:A:317:HIS:O	1:A:320:SER:HB3	2.22	0.40
1:A:561:TRP:CD1	1:A:593:ILE:HG12	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	408/422 (97%)	388 (95%)	20 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/376 (98%)	349 (95%)	18 (5%)	25	11

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	299	ARG
1	A	321	THR
1	A	337	LEU
1	A	351	LYS
1	A	352	ASP
1	A	371	ARG
1	A	390	SER
1	A	391	THR
1	A	476	ASN
1	A	516	ARG
1	A	523	LEU
1	A	547	ARG
1	A	556	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	622	SER
1	A	653	SER
1	A	662	MET
1	A	715	VAL
1	A	717	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	HEM	A	801	1	41,50,50	1.44	6 (14%)	45,82,82	1.68	6 (13%)
3	H4B	A	802	-	16,18,18	0.84	0	11,26,26	2.72	6 (54%)
5	ACT	A	804	-	3,3,3	0.79	0	3,3,3	0.68	0
4	KMR	A	803	-	19,19,19	0.51	0	25,25,25	1.77	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	801	1	-	6/12/54/54	-
3	H4B	A	802	-	-	0/8/17/17	0/2/2/2
4	KMR	A	803	-	-	4/6/16/16	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	HEM	C3C-C2C	-3.50	1.35	1.40
2	A	801	HEM	C3C-CAC	3.37	1.54	1.47
2	A	801	HEM	CAB-C3B	2.68	1.54	1.47
2	A	801	HEM	CMD-C2D	2.31	1.55	1.50
2	A	801	HEM	CMB-C2B	2.11	1.55	1.50
2	A	801	HEM	C2C-C1C	2.02	1.47	1.42

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	803	KMR	C02-N01-C06	6.03	122.67	118.10
2	A	801	HEM	CAD-CBD-CGD	-5.75	101.22	113.60
3	A	802	H4B	C8A-C4A-C4	5.47	119.43	114.57
2	A	801	HEM	C4B-CHC-C1C	4.42	128.39	122.56
2	A	801	HEM	CBA-CAA-C2A	-4.36	105.18	112.62
3	A	802	H4B	C2-N3-C4	4.10	122.44	115.93
4	A	803	KMR	C05-C06-N01	-3.48	119.21	122.90
3	A	802	H4B	N1-C2-N3	-3.30	120.25	125.42
2	A	801	HEM	C1B-NB-C4B	2.59	107.75	105.07
3	A	802	H4B	C2-N1-C8A	2.42	119.97	114.54
4	A	803	KMR	N02-C02-N01	2.33	120.17	116.49
2	A	801	HEM	C3B-C2B-C1B	2.27	108.17	106.49
3	A	802	H4B	C4-C4A-N5	2.12	120.90	119.12
3	A	802	H4B	C4A-C4-N3	-2.11	118.00	124.01
2	A	801	HEM	CAB-C3B-C2B	-2.10	121.70	128.60

There are no chirality outliers.

All (10) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	801	HEM	C4D-C3D-CAD-CBD
2	A	801	HEM	C2D-C3D-CAD-CBD
4	A	803	KMR	C09-C10-N11-C12
4	A	803	KMR	C09-C10-N11-C16
2	A	801	HEM	C3D-CAD-CBD-CGD
2	A	801	HEM	C4B-C3B-CAB-CBB
4	A	803	KMR	N01-C06-C08-C09
4	A	803	KMR	C05-C06-C08-C09
2	A	801	HEM	CAD-CBD-CGD-O1D
2	A	801	HEM	CAD-CBD-CGD-O2D

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	HEM	2	0
3	A	802	H4B	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	411/422 (97%)	1.47	114 (27%) 0 0	42, 78, 137, 174	0

All (114) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	299	ARG	7.7
1	A	611	ALA	7.2
1	A	369	ILE	6.9
1	A	553	TRP	6.8
1	A	551	PHE	6.4
1	A	355	PHE	5.8
1	A	677	VAL	5.7
1	A	493	LEU	5.4
1	A	506	ILE	5.4
1	A	554	PHE	5.4
1	A	715	VAL	5.3
1	A	616	LEU	5.2
1	A	351	LYS	5.1
1	A	491	SER	5.1
1	A	566	ALA	5.1
1	A	680	VAL	5.1
1	A	488	PRO	5.0
1	A	679	ILE	5.0
1	A	373	GLY	4.8
1	A	370	LYS	4.8
1	A	691	PHE	4.7
1	A	567	VAL	4.7
1	A	604	TYR	4.6
1	A	676	TRP	4.6
1	A	678	TRP	4.4
1	A	348	VAL	4.4
1	A	588	TYR	4.4

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	479	LEU	4.3
1	A	443	CYS	4.3
1	A	374	SER	4.1
1	A	442	ILE	3.9
1	A	492	THR	3.9
1	A	354	LEU	3.8
1	A	716	TRP	3.7
1	A	388	ILE	3.7
1	A	318	LEU	3.7
1	A	384	VAL	3.7
1	A	505	CYS	3.7
1	A	490	GLY	3.7
1	A	372	PHE	3.6
1	A	681	PRO	3.5
1	A	615	ASP	3.5
1	A	508	GLN	3.4
1	A	565	PRO	3.4
1	A	350	THR	3.4
1	A	683	MET	3.3
1	A	322	LEU	3.3
1	A	446	VAL	3.3
1	A	562	TYR	3.3
1	A	515	GLY	3.3
1	A	593	ILE	3.3
1	A	591	THR	3.3
1	A	300	PHE	3.2
1	A	711	TRP	3.2
1	A	607	LEU	3.2
1	A	682	PRO	3.2
1	A	480	ILE	3.1
1	A	481	ARG	3.1
1	A	503	GLU	3.1
1	A	610	VAL	3.1
1	A	352	ASP	3.0
1	A	486	LYS	3.0
1	A	583	PRO	3.0
1	A	485	TYR	3.0
1	A	494	GLY	2.9
1	A	685	GLY	2.9
1	A	487	GLN	2.9
1	A	706	TYR	2.9
1	A	510	TRP	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	614	MET	2.8
1	A	564	LEU	2.8
1	A	670	GLY	2.8
1	A	375	LYS	2.7
1	A	619	ARG	2.7
1	A	618	MET	2.7
1	A	478	GLN	2.7
1	A	666	TYR	2.7
1	A	441	TYR	2.6
1	A	563	GLY	2.6
1	A	590	GLY	2.6
1	A	339	SER	2.6
1	A	714	HIS	2.6
1	A	561	TRP	2.6
1	A	589	MET	2.6
1	A	371	ARG	2.5
1	A	609	GLU	2.5
1	A	507	GLN	2.5
1	A	667	ARG	2.5
1	A	514	ARG	2.5
1	A	582	CYS	2.5
1	A	581	ALA	2.4
1	A	381	LEU	2.4
1	A	540	LEU	2.4
1	A	717	LYS	2.4
1	A	391	THR	2.4
1	A	617	ASP	2.4
1	A	477	SER	2.4
1	A	557	LEU	2.3
1	A	571	LEU	2.3
1	A	511	LYS	2.3
1	A	301	LEU	2.3
1	A	606	ILE	2.2
1	A	439	PHE	2.2
1	A	353	GLN	2.2
1	A	386	LYS	2.2
1	A	690	VAL	2.1
1	A	630	LEU	2.1
1	A	385	ASN	2.1
1	A	499	VAL	2.1
1	A	389	GLU	2.1
1	A	592	GLU	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	568	SER	2.1
1	A	460	THR	2.0
1	A	489	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	KMR	A	803	18/18	0.88	0.36	50,71,97,100	0
5	ACT	A	804	4/4	0.91	0.20	86,92,93,94	0
3	H4B	A	802	17/17	0.96	0.22	43,54,63,64	0
2	HEM	A	801	43/43	0.97	0.20	34,48,57,66	0
6	ZN	A	805	1/1	0.99	0.15	52,52,52,52	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around KMR A 803:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HEM A 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers ⓘ

There are no such residues in this entry.