



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 03:59 pm BST

PDB ID : 3TUF  
Title : Structure of the SpoIIQ-SpoIIAH pore forming complex.  
Authors : Levnikov, V.M.; Blagova, E.V.; Wilkinson, A.J.  
Deposited on : 2011-09-16  
Resolution : 2.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	245	<div> <div>5%</div> <div> <div></div> <div>44%</div> <div>18%</div> <div>•</div> <div>36%</div> </div> </div>
2	A	197	<div> <div>2%</div> <div> <div></div> <div>43%</div> <div>11%</div> <div>•</div> <div>42%</div> </div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2200 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Stage II sporulation protein Q.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	158	Total	C	N	O	S	0	0	0
			1202	748	196	254	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	39	GLY	-	EXPRESSION TAG	UNP P71044
B	40	PRO	-	EXPRESSION TAG	UNP P71044
B	41	ALA	-	EXPRESSION TAG	UNP P71044
B	42	MET	-	EXPRESSION TAG	UNP P71044

- Molecule 2 is a protein called Stage III sporulation protein AH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	115	Total	C	N	O	S	0	0	0
			894	552	145	195	2			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	GLY	-	EXPRESSION TAG	UNP P49785
A	23	PRO	-	EXPRESSION TAG	UNP P49785
A	24	ALA	-	EXPRESSION TAG	UNP P49785

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	63	Total	O	0	0
			63	63		
3	A	41	Total	O	0	0
			41	41		



- Molecule 1: Stage II sporulation protein Q



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.80Å 111.09Å 136.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.72 – 2.26 46.72 – 2.26	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.72-2.26) 84.2 (46.72-2.26)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.01 (at 2.27Å)	Xtriage
Refinement program	REFMAC 5.6.0086	Depositor
R, $R_{free}$	0.165 , 0.226 0.178 , 0.229	Depositor DCC
$R_{free}$ test set	1217 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.6	Xtriage
Anisotropy	0.693	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 50.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2200	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	B	1.14	2/1218 (0.2%)	1.04	2/1644 (0.1%)
2	A	1.02	1/900 (0.1%)	1.00	2/1210 (0.2%)
All	All	1.09	3/2118 (0.1%)	1.02	4/2854 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	120	LYS	C-O	5.70	1.34	1.23
1	B	232	THR	CA-CB	5.25	1.67	1.53
2	A	150	GLU	CG-CD	5.11	1.59	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	148	ASP	CB-CG-OD2	-5.91	112.99	118.30
1	B	193	ASN	CB-CA-C	-5.63	99.14	110.40
2	A	148	LEU	CB-CG-CD2	5.44	120.25	111.00
2	A	162	ILE	CB-CA-C	-5.00	101.59	111.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1202	0	1177	47	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	894	0	893	25	0
3	A	41	0	0	2	0
3	B	63	0	0	1	0
All	All	2200	0	2070	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:75:SER:O	1:B:76:MET:HG3	1.43	1.17
2:A:155:GLU:O	2:A:159:GLU:HG3	1.52	1.08
1:B:75:SER:HA	1:B:224:VAL:HG11	1.14	1.06
1:B:219:PHE:HB3	1:B:227:ILE:HD11	1.41	1.03
1:B:219:PHE:CG	1:B:227:ILE:CD1	2.41	1.03
1:B:219:PHE:CG	1:B:227:ILE:HD13	1.94	1.02
1:B:219:PHE:CB	1:B:227:ILE:HD11	1.93	0.99
1:B:75:SER:C	1:B:76:MET:HG3	1.75	0.97
1:B:219:PHE:CD2	1:B:227:ILE:HD13	2.12	0.84
1:B:219:PHE:CG	1:B:227:ILE:HD11	2.09	0.83
2:A:158:LEU:O	2:A:162:ILE:HG12	1.81	0.80
1:B:75:SER:O	1:B:76:MET:CG	2.30	0.77
1:B:219:PHE:CD2	1:B:227:ILE:CD1	2.69	0.76
1:B:75:SER:C	1:B:76:MET:CG	2.54	0.74
1:B:75:SER:HA	1:B:224:VAL:CG1	2.08	0.73
2:A:151:VAL:O	2:A:155:GLU:HG3	1.90	0.72
1:B:228:GLU:O	1:B:228:GLU:HG3	1.89	0.71
1:B:207:ILE:CD1	1:B:216:PRO:HG3	2.20	0.71
2:A:113:LEU:HD11	2:A:117:ARG:NH1	2.05	0.70
2:A:134:THR:HG23	2:A:137:GLU:OE2	1.92	0.69
2:A:191:SER:O	2:A:194:THR:HG22	1.92	0.69
2:A:203:GLU:CD	2:A:203:GLU:H	1.97	0.67
1:B:207:ILE:HD11	1:B:216:PRO:CB	2.26	0.66
1:B:207:ILE:HD12	1:B:216:PRO:HG3	1.76	0.65
2:A:192:LYS:NZ	3:A:331:HOH:O	2.29	0.65
1:B:159:HIS:HE1	1:B:165:THR:OG1	1.79	0.65
1:B:182:LYS:HG2	1:B:185:GLN:HE21	1.62	0.65
1:B:207:ILE:HD12	1:B:207:ILE:N	2.12	0.65
2:A:124:LEU:HB3	2:A:145:MET:HE2	1.79	0.63
2:A:200:VAL:O	2:A:205:LYS:HE2	2.00	0.62

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:219:PHE:CD1	1:B:227:ILE:HD13	2.34	0.61
1:B:139:SER:O	1:B:159:HIS:HD2	1.85	0.60
1:B:207:ILE:HD11	1:B:216:PRO:HA	1.84	0.60
2:A:113:LEU:O	2:A:117:ARG:HG2	2.01	0.60
1:B:153:TYR:HB2	1:B:170:LEU:HB2	1.85	0.59
1:B:207:ILE:HD11	1:B:216:PRO:CA	2.34	0.58
1:B:159:HIS:CE1	1:B:165:THR:OG1	2.57	0.57
1:B:96:TYR:OH	1:B:105:LYS:HD2	2.06	0.56
1:B:207:ILE:CD1	1:B:216:PRO:HA	2.36	0.56
2:A:113:LEU:HD11	2:A:117:ARG:HH11	1.68	0.55
1:B:83:VAL:HG13	3:B:324:HOH:O	2.07	0.54
2:A:201:ALA:HB3	2:A:204:ILE:HG12	1.89	0.53
1:B:207:ILE:HD12	1:B:207:ILE:H	1.73	0.52
2:A:155:GLU:O	2:A:159:GLU:CG	2.43	0.51
1:B:182:LYS:HG2	1:B:185:GLN:NE2	2.26	0.50
1:B:168:GLN:HG2	1:B:204:HIS:HB3	1.92	0.50
2:A:158:LEU:HD12	2:A:162:ILE:HD11	1.93	0.49
1:B:133:ASP:OD1	1:B:189:LYS:NZ	2.45	0.49
1:B:182:LYS:H	1:B:185:GLN:NE2	2.11	0.49
1:B:207:ILE:HD11	1:B:216:PRO:HB3	1.95	0.48
2:A:134:THR:OG1	2:A:137:GLU:HG3	2.16	0.46
2:A:117:ARG:HH22	2:A:155:GLU:CD	2.18	0.46
1:B:207:ILE:CD1	1:B:216:PRO:CG	2.93	0.45
2:A:150:GLU:HG3	3:A:305:HOH:O	2.17	0.43
1:B:103:GLU:HA	1:B:103:GLU:OE1	2.18	0.43
1:B:163:LEU:HD23	1:B:208:ARG:O	2.18	0.43
2:A:114:GLU:O	2:A:115:ASP:C	2.57	0.43
2:A:201:ALA:HB1	2:A:203:GLU:OE1	2.19	0.43
1:B:131:ASP:HB3	1:B:189:LYS:HD3	2.01	0.43
2:A:160:THR:O	2:A:164:THR:HG23	2.20	0.42
1:B:144:LYS:O	1:B:155:VAL:HA	2.20	0.42
1:B:182:LYS:H	1:B:185:GLN:HE21	1.68	0.42
2:A:162:ILE:H	2:A:162:ILE:HG12	1.64	0.42
1:B:207:ILE:CD1	1:B:216:PRO:CA	2.96	0.42
2:A:203:GLU:N	2:A:203:GLU:CD	2.68	0.41
1:B:167:TYR:N	1:B:167:TYR:CD1	2.88	0.41
1:B:166:VAL:HB	1:B:206:GLU:HB2	2.03	0.41
2:A:184:LYS:HD2	2:A:217:SER:HA	2.03	0.41
1:B:116:TYR:CD2	1:B:116:TYR:N	2.89	0.41
1:B:118:LEU:HD21	2:A:197:ILE:HG21	2.03	0.40
1:B:207:ILE:HD11	1:B:216:PRO:CG	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	156/245 (64%)	143 (92%)	13 (8%)	0	100	100
2	A	113/197 (57%)	109 (96%)	4 (4%)	0	100	100
All	All	269/442 (61%)	252 (94%)	17 (6%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	136/211 (64%)	115 (85%)	21 (15%)	2	1
2	A	100/165 (61%)	92 (92%)	8 (8%)	12	10
All	All	236/376 (63%)	207 (88%)	29 (12%)	4	3

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	76	MET
1	B	86	SER
1	B	103	GLU
1	B	112	TYR
1	B	114	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	117	SER
1	B	127	LYS
1	B	146	GLU
1	B	147	LYS
1	B	158	GLU
1	B	163	LEU
1	B	168	GLN
1	B	169	SER
1	B	177	GLN
1	B	180	LYS
1	B	196	SER
1	B	197	GLU
1	B	207	ILE
1	B	208	ARG
1	B	228	GLU
1	B	229	LYS
2	A	114	GLU
2	A	117	ARG
2	A	122	GLU
2	A	124	LEU
2	A	136	LYS
2	A	148	LEU
2	A	158	LEU
2	A	191	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	88	ASN
1	B	159	HIS
1	B	185	GLN
2	A	188	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	158/245 (64%)	0.14	13 (8%) 11 12	29, 49, 85, 100	0
2	A	115/197 (58%)	0.08	4 (3%) 44 46	29, 55, 80, 98	0
All	All	273/442 (61%)	0.12	17 (6%) 20 22	29, 51, 84, 100	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	232	THR	8.2
1	B	75	SER	6.1
1	B	229	LYS	4.8
1	B	231	ALA	4.8
1	B	227	ILE	4.6
1	B	230	ALA	4.4
2	A	103	ASP	3.9
2	A	217	SER	3.8
1	B	161	ASP	3.2
1	B	225	SER	3.0
1	B	228	GLU	2.9
1	B	112	TYR	2.7
2	A	105	LEU	2.6
1	B	76	MET	2.6
2	A	186	ASP	2.3
1	B	160	ALA	2.3
1	B	219	PHE	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.