



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 23, 2021 – 12:58 PM GMT

PDB ID : 6TUU
Title : Leishmania infantum Rad51 surrogate LiRadA10 in complex with 5,6,7,8-tetrahydro-2-naphthoic acid
Authors : Pantelejevs, T.; Hyvonen, M.
Deposited on : 2020-01-08
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.16
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.16


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3764 (1.76-1.72)
Clashscore	141614	3923 (1.76-1.72)
Ramachandran outliers	138981	3878 (1.76-1.72)
Sidechain outliers	138945	3878 (1.76-1.72)
RSRZ outliers	127900	3705 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	
1	B	243	
1	C	243	
1	D	243	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7558 atoms, of which 44 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA repair and recombination protein RadA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	214	Total	C	N	O	S	2	3	0
			1665	1042	299	320	4			
1	B	222	Total	C	N	O	S	2	3	0
			1727	1080	311	332	4			
1	C	225	Total	C	N	O	S	1	2	0
			1740	1087	317	332	4			
1	D	223	Total	C	N	O	S	0	1	0
			1719	1076	312	327	4			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	107	MET	-	initiating methionine	UNP O74036
A	168	ALA	VAL	conflict	UNP O74036
A	169	LEU	ILE	conflict	UNP O74036
A	170	TYR	TRP	conflict	UNP O74036
A	182	LEU	ILE	conflict	UNP O74036
A	198	ALA	LYS	conflict	UNP O74036
A	199	ASN	HIS	conflict	UNP O74036
A	200	VAL	ILE	conflict	UNP O74036
A	201	ALA	TYR	conflict	UNP O74036
A	202	CYS	VAL	conflict	UNP O74036
A	221	MET	LYS	conflict	UNP O74036
A	288	ASN	ARG	conflict	UNP O74036
B	107	MET	-	initiating methionine	UNP O74036
B	168	ALA	VAL	conflict	UNP O74036
B	169	LEU	ILE	conflict	UNP O74036
B	170	TYR	TRP	conflict	UNP O74036
B	182	LEU	ILE	conflict	UNP O74036
B	198	ALA	LYS	conflict	UNP O74036
B	199	ASN	HIS	conflict	UNP O74036
B	200	VAL	ILE	conflict	UNP O74036
B	201	ALA	TYR	conflict	UNP O74036

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	202	CYS	VAL	conflict	UNP O74036
B	221	MET	LYS	conflict	UNP O74036
B	288	ASN	ARG	conflict	UNP O74036
C	107	MET	-	initiating methionine	UNP O74036
C	168	ALA	VAL	conflict	UNP O74036
C	169	LEU	ILE	conflict	UNP O74036
C	170	TYR	TRP	conflict	UNP O74036
C	182	LEU	ILE	conflict	UNP O74036
C	198	ALA	LYS	conflict	UNP O74036
C	199	ASN	HIS	conflict	UNP O74036
C	200	VAL	ILE	conflict	UNP O74036
C	201	ALA	TYR	conflict	UNP O74036
C	202	CYS	VAL	conflict	UNP O74036
C	221	MET	LYS	conflict	UNP O74036
C	288	ASN	ARG	conflict	UNP O74036
D	107	MET	-	initiating methionine	UNP O74036
D	168	ALA	VAL	conflict	UNP O74036
D	169	LEU	ILE	conflict	UNP O74036
D	170	TYR	TRP	conflict	UNP O74036
D	182	LEU	ILE	conflict	UNP O74036
D	198	ALA	LYS	conflict	UNP O74036
D	199	ASN	HIS	conflict	UNP O74036
D	200	VAL	ILE	conflict	UNP O74036
D	201	ALA	TYR	conflict	UNP O74036
D	202	CYS	VAL	conflict	UNP O74036
D	221	MET	LYS	conflict	UNP O74036
D	288	ASN	ARG	conflict	UNP O74036

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is 5,6,7,8-tetrahydronaphthalene-2-carboxylic acid (three-letter code: NY5) (formula: C₁₁H₁₂O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			24	11	11	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	H	O	0	0
			24	11	11	2		
3	C	1	Total	C	H	O	0	0
			24	11	11	2		
3	D	1	Total	C	H	O	0	0
			24	11	11	2		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	149	Total	O	0	0
			149	149		
5	B	159	Total	O	0	0
			159	159		
5	C	147	Total	O	0	0
			147	147		
5	D	145	Total	O	0	0
			145	145		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA repair and recombination protein RadA




- Molecule 1: DNA repair and recombination protein RadA



- Molecule 1: DNA repair and recombination protein RadA



- Molecule 1: DNA repair and recombination protein RadA



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	40.27Å 118.98Å 83.79Å 90.00° 91.42° 90.00°	Depositor
Resolution (Å)	41.88 – 1.74 48.50 – 1.73	Depositor EDS
% Data completeness (in resolution range)	99.1 (41.88-1.74) 99.1 (48.50-1.73)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.44 (at 1.74Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.196 , 0.230 0.198 , 0.236	Depositor DCC
R_{free} test set	3999 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	25.7	Xtriage
Anisotropy	0.395	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 36.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.107 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7558	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.74 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.1399e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, CL, NY5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/1683	0.64	0/2265
1	B	0.48	0/1749	0.65	0/2358
1	C	0.50	0/1762	0.66	0/2373
1	D	0.50	0/1741	0.64	0/2344
All	All	0.50	0/6935	0.65	0/9340

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1665	0	1691	10	0
1	B	1727	0	1747	8	0
1	C	1740	0	1752	7	0
1	D	1719	0	1737	11	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	13	11	0	1	0
3	B	13	11	0	0	0
3	C	13	11	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	13	11	0	0	0
4	C	1	0	0	0	0
5	A	149	0	0	0	0
5	B	159	0	0	1	0
5	C	147	0	0	0	0
5	D	145	0	0	4	0
All	All	7514	44	6927	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:349:ASP:HA	5:D:501:HOH:O	1.77	0.84
1:D:206[A]:PHE:H	1:D:210:HIS:HD2	1.26	0.83
1:D:206[B]:PHE:H	1:D:210:HIS:HD2	1.32	0.75
1:B:330:ALA:HB1	1:B:333:LEU:HD12	1.84	0.60
1:B:136:VAL:HG13	1:B:315:LEU:HD23	1.87	0.56
1:B:257:ARG:HE	1:B:258[A]:GLN:HE21	1.54	0.56
1:A:136[B]:VAL:HG12	1:A:313:VAL:HB	1.90	0.51
1:D:206[A]:PHE:H	1:D:210:HIS:CD2	2.17	0.51
1:D:233:LYS:HG3	5:D:521:HOH:O	2.10	0.50
1:A:203:ALA:HB2	3:A:402:NY5:C3	2.42	0.49
1:A:131:GLN:O	1:C:307:HIS:HD2	1.95	0.49
1:C:245:ARG:HH22	1:C:284:GLN:HB3	1.77	0.49
1:D:183:ARG:HG3	1:D:193:PRO:HB2	1.95	0.49
1:A:136[A]:VAL:HG22	1:A:313:VAL:HB	1.95	0.49
1:C:136:VAL:HG22	1:C:313:VAL:HB	1.96	0.48
1:C:254:LEU:HD23	5:D:583:HOH:O	2.14	0.47
1:D:136:VAL:HG22	1:D:313:VAL:HB	1.96	0.47
1:B:275:TYR:HB2	1:B:277[A]:ILE:HD12	1.96	0.47
1:D:210:HIS:HE1	5:D:503:HOH:O	1.99	0.46
1:B:146:GLN:HG2	5:B:501:HOH:O	2.15	0.46
1:B:136:VAL:HG22	1:B:313:VAL:HB	1.97	0.46
1:D:206[B]:PHE:H	1:D:210:HIS:CD2	2.21	0.45
1:B:344:GLU:H	1:B:344:GLU:CD	2.20	0.45
1:A:137:PHE:HZ	1:A:328:ILE:HG13	1.83	0.44
1:C:136:VAL:HG12	1:C:144:LYS:HG2	2.00	0.44
1:A:133:ILE:HD12	1:A:269:HIS:CD2	2.54	0.43
1:B:315:LEU:HD11	1:B:340:PHE:CE2	2.55	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:322:LYS:HD2	1:C:339:VAL:HG12	2.01	0.42
1:D:340:PHE:CE1	1:D:347:ILE:HD12	2.56	0.41
1:A:315[A]:LEU:HD11	1:A:323:ARG:HB3	2.03	0.41
1:D:146:GLN:HG3	1:D:342:ILE:HG21	2.02	0.41
1:A:236:ILE:HA	1:A:280:PHE:O	2.21	0.40
1:A:237:VAL:HB	1:A:281:VAL:HG12	2.03	0.40
1:A:137:PHE:CZ	1:A:328:ILE:HG13	2.56	0.40
1:C:136:VAL:HG13	1:C:315:LEU:HG	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/243 (87%)	209 (99%)	1 (0%)	1 (0%)	29	12
1	B	221/243 (91%)	218 (99%)	2 (1%)	1 (0%)	29	12
1	C	221/243 (91%)	219 (99%)	2 (1%)	0	100	100
1	D	220/243 (90%)	216 (98%)	4 (2%)	0	100	100
All	All	873/972 (90%)	862 (99%)	9 (1%)	2 (0%)	47	29

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	251	ARG
1	A	251	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	177/197 (90%)	175 (99%)	2 (1%)	73	59
1	B	183/197 (93%)	183 (100%)	0	100	100
1	C	183/197 (93%)	181 (99%)	2 (1%)	73	59
1	D	180/197 (91%)	179 (99%)	1 (1%)	86	79
All	All	723/788 (92%)	718 (99%)	5 (1%)	84	75

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	187	GLN
1	A	230	ARG
1	C	283	ASN
1	C	347	ILE
1	D	269	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	187	GLN
1	A	217	GLN
1	A	227	ASN
1	A	258	GLN
1	B	199	ASN
1	C	211	GLN
1	C	307	HIS
1	D	199	ASN
1	D	209	ASN
1	D	210	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	PO4	B	401	-	4,4,4	2.39	1 (25%)	6,6,6	0.79	0
2	PO4	A	401	-	4,4,4	2.46	1 (25%)	6,6,6	0.75	0
3	NY5	C	401	-	12,14,14	1.82	2 (16%)	16,19,19	0.92	0
3	NY5	B	402	-	12,14,14	2.81	3 (25%)	16,19,19	0.95	0
3	NY5	A	402	-	12,14,14	2.51	2 (16%)	16,19,19	0.76	0
3	NY5	D	401	-	12,14,14	1.34	2 (16%)	16,19,19	1.15	3 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NY5	B	402	-	-	0/0/11/11	0/2/2/2
3	NY5	C	401	-	-	0/0/11/11	0/2/2/2
3	NY5	A	402	-	-	0/0/11/11	0/2/2/2
3	NY5	D	401	-	-	0/0/11/11	0/2/2/2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	402	NY5	C5-C6	8.67	1.55	1.47
3	A	402	NY5	C5-C6	7.74	1.54	1.47
3	C	401	NY5	C5-C6	5.14	1.52	1.47
2	A	401	PO4	P-O1	4.07	1.60	1.50
2	B	401	PO4	P-O1	3.98	1.60	1.50
3	D	401	NY5	C5-C6	2.90	1.50	1.47
3	B	402	NY5	C4-C3	2.82	1.44	1.39
3	C	401	NY5	C3-C2	2.75	1.45	1.40
3	B	402	NY5	C3-C2	2.61	1.45	1.40
3	D	401	NY5	C3-C2	2.57	1.45	1.40
3	A	402	NY5	C3-C2	2.36	1.44	1.40

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	401	NY5	C5-C4-C3	-2.50	118.15	121.90
3	D	401	NY5	C-C5-C6	-2.39	117.16	120.37
3	D	401	NY5	C-C5-C4	2.13	121.18	118.16

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	NY5	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand NY5 C 401



Ligand NY5 B 402





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/243 (88%)	-0.08	6 (2%) 53 58	19, 29, 51, 65	2 (0%)
1	B	222/243 (91%)	0.01	7 (3%) 47 53	19, 31, 54, 66	2 (0%)
1	C	225/243 (92%)	-0.00	10 (4%) 34 39	17, 28, 53, 73	1 (0%)
1	D	223/243 (91%)	-0.22	2 (0%) 84 88	18, 28, 46, 68	1 (0%)
All	All	884/972 (90%)	-0.08	25 (2%) 53 58	17, 29, 53, 73	6 (0%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	108	ALA	4.8
1	C	334	PRO	4.8
1	B	332	HIS	4.7
1	A	251	ARG	4.4
1	A	108	ALA	4.2
1	C	254	LEU	3.7
1	C	140	PHE	3.6
1	A	249	ILE	3.3
1	B	229	ASP	3.2
1	C	252	GLY	3.1
1	C	302	GLY	2.8
1	B	331	PRO	2.8
1	C	251	ARG	2.7
1	B	308	SER	2.7
1	A	252	GLY	2.6
1	D	110	ILE	2.5
1	C	303	HIS	2.4
1	B	252	GLY	2.4
1	A	109	THR	2.3
1	C	285	VAL	2.2
1	C	206	PHE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	140	PHE	2.0
1	C	332	HIS	2.0
1	B	249	ILE	2.0
1	D	329	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NY5	B	402	13/13	0.80	0.26	38,38,53,55	0
3	NY5	D	401	13/13	0.84	0.18	36,38,53,57	0
3	NY5	A	402	13/13	0.88	0.12	37,38,47,51	0
3	NY5	C	401	13/13	0.88	0.12	34,38,47,50	0
2	PO4	B	401	5/5	0.93	0.11	32,33,37,41	0
4	CL	C	402	1/1	0.98	0.08	49,49,49,49	0
2	PO4	A	401	5/5	0.99	0.06	32,34,37,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around NY5 B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NY5 D 401:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around NY5 A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.