



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 08:42 PM BST

PDB ID : 6TVI
Title : Salmonella typhimurium mutant neuraminidase (D100S)+ DANA
Authors : Garman, E.F.; Salinger, M.T.; Murray, J.W.; Laver, W.G.; Kuhn, P.; Vimr, E.R.
Deposited on : 2020-01-09
Resolution : 1.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1050 (1.06-0.94)
Clashscore	141614	1117 (1.06-0.94)
Ramachandran outliers	138981	1043 (1.06-0.94)
Sidechain outliers	138945	1045 (1.06-0.94)
RSRZ outliers	127900	1023 (1.06-0.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	379	<div> <div>8%</div> <div>88%</div> <div>11% ..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	AAA	406	-	X	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7194 atoms, of which 3342 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Sialidase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	AAA	379	6523	2034	3262	575	635	17	198	41	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	100	SER	ASP	engineered mutation	UNP P29768

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	AAA	1	Total	C	H	O	2	0
			14	3	8	3		
2	AAA	1	Total	C	H	O	4	1
			28	6	16	6		
2	AAA	1	Total	C	H	O	2	0
			14	3	8	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	AAA	1	Total	C	H	O	4	1
			28	6	16	6		
2	AAA	1	Total	C	H	O	2	0
			14	3	8	3		
2	AAA	1	Total	C	H	O	2	0
			14	3	8	3		

- Molecule 3 is 2-DEOXY-2,3-DEHYDRO-N-ACETYL-NEURAMINIC ACID (three-letter code: DAN) (formula: $C_{11}H_{17}NO_8$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	AAA	1	Total	C	H	N	O	4	0
			36	11	16	1	8		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	487	Total	O	0	36
			523	523		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sialidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	47.07Å 81.78Å 91.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	21.96 – 1.00 21.95 – 1.00	Depositor EDS
% Data completeness (in resolution range)	99.6 (21.96-1.00) 99.6 (21.95-1.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.22 (at 1.00Å)	Xtriage
Refinement program	REFMAC 5.8.0257	Depositor
R, R_{free}	0.118 , 0.133 0.123 , 0.138	Depositor DCC
R_{free} test set	5722 reflections (3.03%)	wwPDB-VP
Wilson B-factor (Å ²)	6.1	Xtriage
Anisotropy	0.080	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.50 , 50.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	7194	wwPDB-VP
Average B, all atoms (Å ²)	9.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, DAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.15	9/3322 (0.3%)	1.06	14/4478 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	AAA	0	1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	4	GLU	CD-OE1	42.10	1.72	1.25
1	AAA	323	GLU	CD-OE1	11.65	1.38	1.25
1	AAA	255	GLU	CD-OE2	-10.73	1.13	1.25
1	AAA	255	GLU	CD-OE1	-7.07	1.17	1.25
1	AAA	195[A]	MET	CG-SD	6.71	1.98	1.81
1	AAA	195[B]	MET	CG-SD	6.71	1.98	1.81
1	AAA	354	GLU	CD-OE2	-5.86	1.19	1.25
1	AAA	4	GLU	CG-CD	5.61	1.60	1.51
1	AAA	380	SER	CB-OG	5.06	1.48	1.42

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	4	GLU	CG-CD-OE2	-10.14	98.03	118.30
1	AAA	373[A]	ARG	NE-CZ-NH1	9.81	125.21	120.30
1	AAA	373[B]	ARG	NE-CZ-NH1	9.81	125.21	120.30
1	AAA	111	ARG	NE-CZ-NH1	-9.31	115.64	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	5	LYS	N-CA-CB	8.52	125.94	110.60
1	AAA	156[A]	LYS	CD-CE-NZ	-8.47	92.22	111.70
1	AAA	156[B]	LYS	CD-CE-NZ	-8.47	92.22	111.70
1	AAA	4	GLU	CG-CD-OE1	-7.96	102.39	118.30
1	AAA	156[A]	LYS	CG-CD-CE	-7.80	88.50	111.90
1	AAA	156[B]	LYS	CG-CD-CE	-7.80	88.50	111.90
1	AAA	254	PHE	CB-CG-CD1	7.06	125.74	120.80
1	AAA	4	GLU	OE1-CD-OE2	7.00	131.70	123.30
1	AAA	329	ASP	CB-CG-OD2	-5.58	113.27	118.30
1	AAA	5	LYS	CB-CA-C	-5.41	99.57	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	AAA	4	GLU	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	3261	3262	3217	55	0
2	AAA	48	64	64	8	1
3	AAA	20	16	16	0	0
4	AAA	523	0	0	13	1
All	All	3852	3342	3297	58	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:4:GLU:CD	1:AAA:4:GLU:OE1	1.72	1.28
2:AAA:403:GOL:O2	4:AAA:688[B]:HOH:O	1.76	1.03

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:287[A]:ILE:HD12	1:AAA:378:ILE:HG21	1.39	1.03
1:AAA:287[A]:ILE:CD1	1:AAA:378:ILE:HG21	1.92	0.98
1:AAA:10[B]:LYS:HE3	4:AAA:915:HOH:O	1.66	0.95
1:AAA:19:GLN:H	1:AAA:19:GLN:HE21	1.03	0.93
1:AAA:117[B]:MET:CE	1:AAA:142:LEU:HD11	2.03	0.89
1:AAA:195[A]:MET:HE1	1:AAA:224:TYR:HE1	1.39	0.85
1:AAA:287[A]:ILE:HD12	1:AAA:378:ILE:CG2	2.06	0.84
1:AAA:287[A]:ILE:HD11	1:AAA:296:ALA:HB2	1.60	0.84
1:AAA:117[B]:MET:HE2	1:AAA:142:LEU:HD11	1.60	0.81
1:AAA:195[A]:MET:CE	1:AAA:224:TYR:HE1	1.93	0.81
1:AAA:171:ILE:HG12	1:AAA:195[A]:MET:HE3	1.63	0.81
1:AAA:377:VAL:HG23	4:AAA:728:HOH:O	1.80	0.80
1:AAA:163:ASP:CB	4:AAA:539:HOH:O	2.34	0.75
1:AAA:128:TRP:CZ3	1:AAA:175[B]:LEU:HD22	2.22	0.75
1:AAA:156[B]:LYS:NZ	4:AAA:501:HOH:O	2.21	0.74
1:AAA:163:ASP:HB3	4:AAA:539:HOH:O	1.86	0.73
1:AAA:146:LYS:HG3	4:AAA:579:HOH:O	1.91	0.71
1:AAA:195[A]:MET:CE	1:AAA:224:TYR:CE1	2.74	0.69
1:AAA:19:GLN:HE21	1:AAA:19:GLN:N	1.86	0.68
2:AAA:405:GOL:H32	4:AAA:719:HOH:O	1.94	0.67
1:AAA:148:THR:HG23	4:AAA:579:HOH:O	1.98	0.62
1:AAA:163:ASP:CG	4:AAA:539:HOH:O	2.37	0.62
1:AAA:121:TRP:HH2	1:AAA:175[B]:LEU:HD21	1.65	0.61
1:AAA:117[B]:MET:CE	1:AAA:142:LEU:HD21	2.33	0.59
1:AAA:128:TRP:CH2	1:AAA:175[B]:LEU:HD22	2.37	0.59
1:AAA:195[A]:MET:HE2	1:AAA:224:TYR:CE1	2.38	0.58
1:AAA:82[B]:LYS:NZ	4:AAA:504:HOH:O	2.37	0.57
1:AAA:19:GLN:H	1:AAA:19:GLN:NE2	1.88	0.56
1:AAA:117[B]:MET:HE1	1:AAA:142:LEU:HD21	1.87	0.55
1:AAA:132:ARG:NH1	1:AAA:204:VAL:HG22	2.23	0.54
1:AAA:128:TRP:CE3	1:AAA:175[B]:LEU:HD22	2.42	0.54
1:AAA:171:ILE:CG1	1:AAA:195[A]:MET:HE3	2.36	0.54
1:AAA:205:LEU:HD22	4:AAA:513:HOH:O	2.10	0.52
1:AAA:4:GLU:CG	1:AAA:4:GLU:OE1	2.58	0.51
1:AAA:195[A]:MET:HE1	1:AAA:224:TYR:CE1	2.31	0.49
1:AAA:8:VAL:O	1:AAA:82[B]:LYS:HE2	2.13	0.49
1:AAA:117[B]:MET:CE	1:AAA:142:LEU:CD1	2.85	0.47
2:AAA:405:GOL:C3	4:AAA:719:HOH:O	2.60	0.47
1:AAA:287[A]:ILE:CD1	1:AAA:378:ILE:CG2	2.77	0.46
1:AAA:351:VAL:H	2:AAA:406:GOL:H2	1.83	0.44
1:AAA:131:TYR:O	1:AAA:172[B]:SER:OG	2.30	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:174:MET:O	1:AAA:175[B]:LEU:HD13	2.17	0.44
1:AAA:351:VAL:H	2:AAA:406:GOL:C2	2.30	0.44
1:AAA:287[A]:ILE:HD13	1:AAA:378:ILE:HG21	1.93	0.43
1:AAA:351:VAL:HA	2:AAA:406:GOL:H2	2.00	0.43
1:AAA:171:ILE:HD11	1:AAA:195[A]:MET:HE1	2.00	0.43
1:AAA:287[A]:ILE:CD1	1:AAA:296:ALA:HB2	2.38	0.43
1:AAA:277[A]:ASN:HB2	1:AAA:305:ASN:OD1	2.20	0.42
1:AAA:156[A]:LYS:HE3	1:AAA:156[A]:LYS:HB3	1.75	0.41
1:AAA:207:THR:OG1	1:AAA:225:CYS:HB3	2.20	0.41
1:AAA:4:GLU:HB3	1:AAA:4:GLU:OE1	2.20	0.41
1:AAA:237:PHE:O	1:AAA:238:ASN:C	2.59	0.41
1:AAA:117[B]:MET:HE3	1:AAA:142:LEU:HD21	2.01	0.41
1:AAA:351:VAL:CA	2:AAA:406:GOL:H2	2.51	0.40
1:AAA:298:SER:HA	1:AAA:313:SER:O	2.22	0.40
1:AAA:82[A]:LYS:HE3	2:AAA:405:GOL:O3	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:AAA:406:GOL:O2	4:AAA:889:HOH:O[4_456]	1.94	0.26

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	AAA	418/379 (110%)	401 (96%)	14 (3%)	3 (1%)	22 4

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	178	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	AAA	351	VAL
1	AAA	5	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	AAA	366/325 (113%)	361 (99%)	5 (1%)	67 34

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	4	GLU
1	AAA	5	LYS
1	AAA	19	GLN
1	AAA	195[A]	MET
1	AAA	195[B]	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	AAA	405	-	5,5,5	0.39	0	5,5,5	0.64	0
2	GOL	AAA	401	-	5,5,5	0.32	0	5,5,5	0.72	0
2	GOL	AAA	404[B]	-	5,5,5	0.34	0	5,5,5	0.48	0
2	GOL	AAA	402[B]	-	5,5,5	0.59	0	5,5,5	0.46	0
2	GOL	AAA	402[A]	-	5,5,5	0.28	0	5,5,5	0.47	0
3	DAN	AAA	407	-	17,20,20	0.95	1 (5%)	18,28,28	0.70	0
2	GOL	AAA	403	-	5,5,5	0.30	0	5,5,5	0.85	0
2	GOL	AAA	406	-	5,5,5	1.57	1 (20%)	5,5,5	2.46	3 (60%)
2	GOL	AAA	404[A]	-	5,5,5	0.64	0	5,5,5	1.18	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	AAA	405	-	-	3/4/4/4	-
2	GOL	AAA	401	-	-	0/4/4/4	-
2	GOL	AAA	404[B]	-	-	0/4/4/4	-
2	GOL	AAA	402[B]	-	-	0/4/4/4	-
2	GOL	AAA	402[A]	-	-	2/4/4/4	-
3	DAN	AAA	407	-	-	0/14/34/34	0/1/1/1
2	GOL	AAA	403	-	-	0/4/4/4	-
2	GOL	AAA	406	-	-	2/4/4/4	-
2	GOL	AAA	404[A]	-	-	2/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	AAA	407	DAN	C8-C7	-2.75	1.48	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	AAA	406	GOL	C3-C2	2.09	1.60	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	AAA	406	GOL	O1-C1-C2	3.84	128.60	110.20
2	AAA	406	GOL	O3-C3-C2	2.79	123.57	110.20
2	AAA	406	GOL	C3-C2-C1	-2.63	101.49	111.70
2	AAA	404[A]	GOL	O3-C3-C2	2.40	121.69	110.20

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	AAA	402[A]	GOL	C1-C2-C3-O3
2	AAA	406	GOL	O1-C1-C2-C3
2	AAA	404[A]	GOL	C1-C2-C3-O3
2	AAA	402[A]	GOL	O2-C2-C3-O3
2	AAA	405	GOL	C1-C2-C3-O3
2	AAA	406	GOL	O1-C1-C2-O2
2	AAA	404[A]	GOL	O2-C2-C3-O3
2	AAA	405	GOL	O2-C2-C3-O3
2	AAA	405	GOL	O1-C1-C2-O2


There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	AAA	405	GOL	3	0
2	AAA	403	GOL	1	0
2	AAA	406	GOL	4	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	379/379 (100%)	1.12	29 (7%) 13 14	4, 7, 15, 81	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	5	LYS	5.4
1	AAA	4	GLU	5.0
1	AAA	238	ASN	4.9
1	AAA	291	ASN	4.6
1	AAA	382	ASN	4.4
1	AAA	186	ASP	4.3
1	AAA	20	LYS	3.9
1	AAA	381	TYR	3.6
1	AAA	42[A]	CYS	3.1
1	AAA	133	ASP	3.1
1	AAA	94[A]	LYS	3.0
1	AAA	175[A]	LEU	3.0
1	AAA	202[A]	THR	2.7
1	AAA	103[A]	CYS	2.6
1	AAA	163	ASP	2.5
1	AAA	78	LYS	2.5
1	AAA	23[A]	THR	2.5
1	AAA	185[A]	ASN	2.4
1	AAA	200	ASN	2.4
1	AAA	125	ASP	2.4
1	AAA	239	ALA	2.4
1	AAA	121	TRP	2.3
1	AAA	353	LYS	2.2
1	AAA	209	PHE	2.2
1	AAA	344	CYS	2.1
1	AAA	156[A]	LYS	2.1
1	AAA	333[A]	LYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	AAA	261[A]	LYS	2.0
1	AAA	167	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	AAA	404[B]	6/6	0.83	0.27	1,1,1,1	14
2	GOL	AAA	404[A]	6/6	0.83	0.27	10,12,14,14	14
2	GOL	AAA	406	6/6	0.89	0.27	9,12,33,34	14
2	GOL	AAA	403	6/6	0.92	0.20	5,10,11,11	14
2	GOL	AAA	402[B]	6/6	0.92	0.22	5,6,7,7	14
2	GOL	AAA	402[A]	6/6	0.92	0.22	21,29,42,44	14
2	GOL	AAA	405	6/6	0.94	0.30	13,25,38,39	2
2	GOL	AAA	401	6/6	0.97	0.14	8,11,12,12	2
3	DAN	AAA	407	20/20	0.98	0.10	4,5,6,7	4

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around DAN AAA 407:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.