



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 01:33 PM BST

PDB ID : 6TX0  
Title : Crystal structure of tetrameric human D137N-SAMHD1 (residues 109-626) with XTP, dAMPNPP and Mg  
Authors : Morris, E.R.; Kunzelmann, S.; Caswell, S.J.; Arnold, L.H.; Purkiss, A.G.; Kelly, G.; Taylor, I.A.  
Deposited on : 2020-01-13  
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	520	<div> <div>3%</div> <div> <div></div> <div>86%</div> <div>6%</div> <div>8%</div> </div> </div>
1	B	520	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>7%</div> <div>8%</div> </div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8040 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Deoxynucleoside triphosphate triphosphohydrolase SAMHD1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	478	Total	C	N	O	S	0	0	0
			3859	2471	673	695	20			
1	B	479	Total	C	N	O	S	0	0	0
			3861	2473	670	698	20			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	107	GLY	-	expression tag	UNP Q9Y3Z3
A	108	SER	-	expression tag	UNP Q9Y3Z3
A	137	ASN	ASP	engineered mutation	UNP Q9Y3Z3
B	107	GLY	-	expression tag	UNP Q9Y3Z3
B	108	SER	-	expression tag	UNP Q9Y3Z3
B	137	ASN	ASP	engineered mutation	UNP Q9Y3Z3


- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Fe	0	0
			1	1		
2	A	1	Total	Fe	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by author).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	3	Total	Mg	0	0
			3	3		
3	A	3	Total	Mg	0	0
			3	3		

- Molecule 4 is 2'-deoxy-5'-O-[(R)-hydroxy{[(R)-hydroxy(phosphonooxy)phosphoryl]amino}phosphoryl]adenosine (three-letter code: DZ4) (formula:  $C_{10}H_{17}N_6O_{11}P_3$ ) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			30	10	6	11	3		
4	A	1	Total	C	N	O	P	0	0
			30	10	6	11	3		
4	B	1	Total	C	N	O	P	0	0
			30	10	6	11	3		
4	B	1	Total	C	N	O	P	0	0
			30	10	6	11	3		

- Molecule 5 is [[(2 {R},3 {S},4 {R},5 {R})-5-[2,6-bis(oxidanylidene)-3 {H}-purin-9-yl]-3,4-bis(oxidanyl)oxolan-2-yl]methoxy-oxidanyl-phosphoryl] phosphono hydrogen phosphate (three-letter code: CZF) (formula:  $C_{10}H_{15}N_4O_{15}P_3$ ) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			32	10	4	15	3		
5	B	1	Total	C	N	O	P	0	0
			32	10	4	15	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	60	Total	O	0	0
			60	60		
6	B	68	Total	O	0	0
			68	68		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Deoxynucleoside triphosphate triphosphohydrolase SAMHD1



- Molecule 1: Deoxynucleoside triphosphate triphosphohydrolase SAMHD1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.92Å 105.92Å 198.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.31 – 2.01 72.34 – 2.00	Depositor EDS
% Data completeness (in resolution range)	60.1 (72.31-2.01) 59.2 (72.34-2.00)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.83 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.184 , 0.228 0.192 , 0.234	Depositor DCC
$R_{free}$ test set	2306 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.1	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 33.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8040	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CZF, FE, DZ4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.01	2/3952 (0.1%)	1.05	12/5347 (0.2%)
1	B	1.03	5/3954 (0.1%)	1.05	11/5352 (0.2%)
All	All	1.02	7/7906 (0.1%)	1.05	23/10699 (0.2%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	161	SER	CB-OG	6.16	1.50	1.42
1	B	328	ASN	CG-ND2	5.89	1.47	1.32
1	A	161	SER	CB-OG	5.75	1.49	1.42
1	B	388	ASP	CB-CG	-5.53	1.40	1.51
1	B	166	GLU	CG-CD	5.41	1.60	1.51
1	A	368	SER	CB-OG	-5.12	1.35	1.42
1	B	583	ASP	CB-CG	5.06	1.62	1.51

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	442	ARG	NE-CZ-NH1	8.71	124.65	120.30
1	A	366	ARG	NE-CZ-NH1	8.48	124.54	120.30
1	B	239	MET	CG-SD-CE	7.73	112.56	100.20
1	A	239	MET	CG-SD-CE	7.69	112.51	100.20
1	B	442	ARG	NE-CZ-NH1	7.47	124.03	120.30
1	A	226	ARG	NE-CZ-NH1	7.19	123.90	120.30
1	A	164	ARG	NE-CZ-NH1	-6.69	116.95	120.30
1	A	353	ASP	CB-CG-OD1	6.52	124.17	118.30
1	B	348	ARG	NE-CZ-NH2	6.36	123.48	120.30
1	A	576	ARG	NE-CZ-NH1	6.31	123.45	120.30
1	A	472	ASP	CB-CG-OD2	6.20	123.88	118.30
1	B	576	ARG	NE-CZ-NH1	6.08	123.34	120.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	143	ARG	NE-CZ-NH2	-5.98	117.31	120.30
1	B	348	ARG	NE-CZ-NH1	-5.82	117.39	120.30
1	B	120	ASP	CB-CG-OD2	5.78	123.51	118.30
1	B	226	ARG	NE-CZ-NH2	-5.77	117.41	120.30
1	A	120	ASP	CB-CG-OD2	5.74	123.47	118.30
1	B	383	ASP	CB-CG-OD1	5.53	123.27	118.30
1	A	143	ARG	NE-CZ-NH1	5.37	122.99	120.30
1	B	339	ARG	NE-CZ-NH1	5.33	122.97	120.30
1	B	353	ASP	CB-CG-OD1	5.32	123.09	118.30
1	A	134	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	B	353	ASP	CB-CG-OD2	-5.16	113.65	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3859	0	3795	12	0
1	B	3861	0	3788	14	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
4	A	60	0	26	1	0
4	B	60	0	26	0	0
5	A	32	0	0	0	0
5	B	32	0	0	0	0
6	A	60	0	0	1	0
6	B	68	0	0	0	0
All	All	8040	0	7635	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:425:ASN:OD1	1:B:425:ASN:ND2	2.04	0.91
1:A:463:THR:HG21	1:A:577:ASN:O	1.80	0.81
1:B:475:SER:HA	1:B:478:LYS:HE2	1.75	0.68
1:B:593:PRO:HA	1:B:599:ASN:HD21	1.64	0.63
1:A:486:LYS:O	1:A:488:LEU:HD12	2.01	0.61
1:B:488:LEU:O	1:B:489:LEU:HD23	2.01	0.61
4:A:705:DZ4:PA	6:A:808:HOH:O	2.63	0.57
1:A:472:ASP:N	1:A:472:ASP:OD1	2.36	0.56
1:B:472:ASP:OD1	1:B:472:ASP:N	2.36	0.55
1:A:463:THR:CG2	1:A:577:ASN:O	2.54	0.55
1:B:487:VAL:HG12	1:B:487:VAL:O	2.10	0.52
1:A:463:THR:O	1:A:466:ILE:HG13	2.09	0.51
1:B:402:ALA:O	1:B:405:LYS:HG3	2.14	0.48
1:A:285:TRP:CH2	1:A:295:SER:HB3	2.53	0.44
1:B:291:PRO:HG2	1:B:293:ASN:OD1	2.16	0.44
1:A:194:ARG:CZ	1:A:260:ILE:HD12	2.49	0.42
1:B:194:ARG:CZ	1:B:260:ILE:HD12	2.48	0.42
1:A:475:SER:HA	1:A:478:LYS:CE	2.50	0.42
1:A:483:ALA:O	1:A:485:PRO:HD3	2.20	0.41
1:B:489:LEU:CD1	1:B:568:TYR:CZ	3.03	0.41
1:A:475:SER:HA	1:A:478:LYS:HE2	2.02	0.41
1:B:175:ALA:HB1	1:B:199:VAL:HG12	2.02	0.41
1:B:285:TRP:CH2	1:B:295:SER:HB3	2.55	0.41
1:B:225:ALA:O	1:B:226:ARG:HG3	2.21	0.40
1:B:381:ILE:HD12	1:B:381:ILE:HA	1.96	0.40
1:A:175:ALA:HB1	1:A:199:VAL:HG12	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	474/520 (91%)	455 (96%)	19 (4%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	475/520 (91%)	456 (96%)	18 (4%)	1 (0%)	47	44
All	All	949/1040 (91%)	911 (96%)	37 (4%)	1 (0%)	51	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	486	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	412/464 (89%)	406 (98%)	6 (2%)	65	69
1	B	412/464 (89%)	406 (98%)	6 (2%)	65	69
All	All	824/928 (89%)	812 (98%)	12 (2%)	65	69

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	315	TYR
1	A	331	TYR
1	A	368	SER
1	A	388	ASP
1	A	472	ASP
1	A	543	GLU
1	B	315	TYR
1	B	326	GLN
1	B	331	TYR
1	B	368	SER
1	B	388	ASP
1	B	472	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	375	GLN
1	A	380	ASN
1	B	375	GLN
1	B	380	ASN
1	B	594	GLN
1	B	599	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 8 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	DZ4	A	707	3	29,32,32	1.28	4 (13%)	33,50,50	1.72	6 (18%)
5	CZF	A	706	3	25,34,34	1.54	3 (12%)	29,54,54	2.56	5 (17%)
5	CZF	B	707	3	25,34,34	1.16	1 (4%)	29,54,54	2.34	3 (10%)
4	DZ4	B	701	3	29,32,32	1.34	4 (13%)	33,50,50	2.02	7 (21%)
4	DZ4	A	705	3,2	29,32,32	1.41	2 (6%)	33,50,50	1.43	5 (15%)
4	DZ4	B	706	3,2	29,32,32	1.54	6 (20%)	33,50,50	1.54	5 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DZ4	A	707	3	-	2/15/34/34	0/3/3/3
5	CZF	A	706	3	-	5/18/38/38	0/3/3/3
5	CZF	B	707	3	-	3/18/38/38	0/3/3/3
4	DZ4	B	701	3	-	3/15/34/34	0/3/3/3
4	DZ4	A	705	3,2	-	2/15/34/34	0/3/3/3
4	DZ4	B	706	3,2	-	2/15/34/34	0/3/3/3

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	706	DZ4	PA-O1A	5.36	1.54	1.46
4	A	705	DZ4	PB-O1B	4.81	1.53	1.46
5	A	706	CZF	C5-N2	4.41	1.38	1.33
5	A	706	CZF	C9-N1	4.30	1.40	1.33
5	B	707	CZF	C9-N1	3.81	1.39	1.33
4	A	705	DZ4	PA-O1A	3.43	1.51	1.46
4	B	701	DZ4	PB-O1B	3.25	1.51	1.46
4	B	706	DZ4	PA-O2A	-3.12	1.48	1.56
4	A	707	DZ4	PA-O2A	-3.02	1.48	1.56
4	A	707	DZ4	PA-N3A	2.97	1.71	1.63
4	B	701	DZ4	PB-O3B	2.79	1.62	1.59
4	A	707	DZ4	PA-O1A	2.60	1.50	1.46
4	B	706	DZ4	C8-N7	-2.59	1.30	1.34
4	B	706	DZ4	PB-O1B	2.54	1.50	1.46
5	A	706	CZF	P2-O11	-2.52	1.43	1.55
4	A	707	DZ4	PB-O1B	2.43	1.50	1.46
4	B	701	DZ4	PB-N3A	2.42	1.69	1.63
4	B	706	DZ4	PG-O2G	-2.34	1.45	1.54
4	B	706	DZ4	PB-O2B	-2.20	1.50	1.56
4	B	701	DZ4	PG-O3G	-2.15	1.46	1.54

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	706	CZF	C9-N1-C2	10.59	124.08	115.14
5	B	707	CZF	C9-N1-C2	9.70	123.33	115.14
5	B	707	CZF	C7-C9-N1	-6.50	114.54	123.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	706	CZF	C7-C9-N1	-6.26	114.87	123.43
4	B	701	DZ4	O1B-PB-N3A	-6.19	102.66	111.77
4	B	701	DZ4	O2B-PB-O1B	5.80	122.09	109.92
4	A	707	DZ4	O2A-PA-O1A	5.80	122.07	109.92
4	B	706	DZ4	O2B-PB-O1B	4.38	119.11	109.92
4	A	705	DZ4	O2B-PB-O1B	4.36	119.06	109.92
4	B	706	DZ4	O1B-PB-N3A	4.30	118.10	111.77
4	A	707	DZ4	O2B-PB-O1B	4.05	118.41	109.92
4	A	705	DZ4	O2A-PA-O1A	3.83	117.95	109.92
4	B	701	DZ4	C5-C6-N6	3.67	125.93	120.35
4	A	707	DZ4	O2A-PA-O5'	-3.32	97.76	106.75
4	B	701	DZ4	O5'-PA-O1A	-3.15	102.13	114.24
4	A	707	DZ4	O1A-PA-N3A	-2.97	107.40	111.77
4	B	706	DZ4	O5'-PA-O1A	-2.79	103.50	114.24
4	B	706	DZ4	O2A-PA-O1A	2.79	115.78	109.92
4	B	701	DZ4	O2A-PA-O1A	2.58	115.33	109.92
4	B	706	DZ4	C5-C6-N6	2.57	124.26	120.35
5	A	706	CZF	C1-N4-C5	2.35	130.77	126.64
4	A	705	DZ4	O1A-PA-N3A	-2.26	108.45	111.77
4	A	705	DZ4	O2G-PG-O3B	2.25	112.17	104.64
4	B	701	DZ4	O1A-PA-N3A	-2.21	108.52	111.77
5	A	706	CZF	O3-C4-C6	-2.20	104.69	111.05
4	A	707	DZ4	C5-C6-N6	2.15	123.61	120.35
4	B	701	DZ4	C5-C6-N1	-2.15	115.49	120.35
4	A	707	DZ4	O5'-PA-O1A	-2.14	106.02	114.24
4	A	705	DZ4	O5'-PA-O1A	-2.13	106.05	114.24
5	B	707	CZF	O4-C6-C4	-2.12	100.92	105.11
5	A	706	CZF	O10-P3-O15	-2.07	99.72	111.19

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	701	DZ4	PB-N3A-PA-O1A
4	B	701	DZ4	PA-N3A-PB-O1B
4	B	701	DZ4	PG-O3B-PB-O1B
4	A	707	DZ4	PB-N3A-PA-O1A
4	A	707	DZ4	PA-N3A-PB-O1B
5	A	706	CZF	P2-O10-P3-O14
4	A	705	DZ4	PA-N3A-PB-O1B
4	B	706	DZ4	PA-N3A-PB-O1B
5	B	707	CZF	C6-C8-O5-P1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
4	A	705	DZ4	PB-N3A-PA-O5'
5	A	706	CZF	P2-O7-P1-O9
5	A	706	CZF	C6-C8-O5-P1
4	B	706	DZ4	PG-O3B-PB-O2B
5	A	706	CZF	P2-O10-P3-O15
5	A	706	CZF	P3-O10-P2-O11
5	B	707	CZF	P2-O7-P1-O8
5	B	707	CZF	P2-O7-P1-O9


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	705	DZ4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

## Ligand DZ4 A 707




## Ligand CZF A 706







## Ligand CZF B 707




## Ligand DZ4 B 701



## Ligand DZ4 A 705



## Ligand DZ4 B 706



## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	478/520 (91%)	-0.19	14 (2%) 51 50	18, 41, 79, 128	0
1	B	479/520 (92%)	-0.14	10 (2%) 63 62	20, 43, 77, 125	0
All	All	957/1040 (92%)	-0.16	24 (2%) 57 56	18, 42, 79, 128	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	488	LEU	7.0
1	B	487	VAL	6.3
1	B	488	LEU	6.2
1	A	487	VAL	5.5
1	A	491	VAL	4.6
1	A	490	ASP	4.3
1	A	464	GLY	4.1
1	B	114	THR	3.8
1	A	599	ASN	3.6
1	A	598	TRP	3.5
1	B	599	ASN	3.4
1	A	489	LEU	3.4
1	A	592	THR	3.1
1	B	594	GLN	2.8
1	A	590	LEU	2.7
1	B	344	ASP	2.7
1	B	489	LEU	2.6
1	A	466	ILE	2.6
1	B	593	PRO	2.4
1	A	593	PRO	2.4
1	A	284	LEU	2.3
1	A	563	TYR	2.2
1	B	343	VAL	2.2
1	B	486	LYS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	B	705	1/1	0.96	0.04	35,35,35,35	0
3	MG	B	703	1/1	0.97	0.06	36,36,36,36	0
4	DZ4	B	701	30/30	0.98	0.10	20,24,35,35	0
3	MG	A	703	1/1	0.98	0.04	32,32,32,32	0
5	CZF	B	707	32/32	0.98	0.08	27,32,38,42	0
3	MG	B	704	1/1	0.98	0.07	37,37,37,37	0
4	DZ4	A	705	30/30	0.98	0.08	23,26,31,38	0
4	DZ4	B	706	30/30	0.98	0.11	23,27,30,31	0
3	MG	A	704	1/1	0.99	0.09	32,32,32,32	0
5	CZF	A	706	32/32	0.99	0.10	24,28,33,34	0
4	DZ4	A	707	30/30	0.99	0.07	24,26,33,35	0
2	FE	B	702	1/1	1.00	0.12	28,28,28,28	0
3	MG	A	702	1/1	1.00	0.05	29,29,29,29	0
2	FE	A	701	1/1	1.00	0.10	22,22,22,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around MG B 705:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MG B 703:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around DZ4 B 701:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around MG A 703:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around CZF B 707:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MG B 704:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around DZ4 A 705:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around DZ4 B 706:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MG A 704:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around CZF A 706:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around DZ4 A 707:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around FE B 702:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MG A 702:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers ⓘ

There are no such residues in this entry.