



wwPDB Geometry-Only Validation Summary Report ⓘ

May 21, 2020 – 07:13 am BST

PDB ID : 5TY5
Title : Neutron structure from microgravity-grown crystals of Inorganic Pyrophosphatase from *Thermococcus theoreducens*
Authors : Inoguchi, N.; Coates, L.; Morris, M.L.; Singhal, A.; Monaco, D.A.; Garcia-Ruiz, J.M.; Pusey, M.L.; Ng, J.D.
Deposited on : 2016-11-18
Resolution : 2.30 Å(reported)

This is a wwPDB Geometry-Only Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

NEUTRON DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	178	96%
1	B	178	97%
1	C	178	95%
1	D	178	96%
1	E	178	94%
1	F	178	97%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 19178 atoms, of which 8119 are hydrogens and 2034 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inorganic pyrophosphatase.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	174	Total	C	D	H	N	O	S	0	152	0
			3000	941	225	1337	226	266	5			
1	B	175	Total	C	D	H	N	O	S	0	155	0
			3047	946	233	1368	227	267	6			
1	C	174	Total	C	D	H	N	O	S	0	155	0
			3038	944	234	1362	226	266	6			
1	D	174	Total	C	D	H	N	O	S	0	151	0
			3017	941	228	1351	226	266	5			
1	E	174	Total	C	D	H	N	O	S	0	151	0
			3034	944	231	1361	226	266	6			
1	F	173	Total	C	D	H	N	O	S	0	153	0
			3009	939	235	1340	225	265	5			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	67	Total	D	O	0	0
			171	104	67		
2	B	73	Total	D	O	0	0
			187	114	73		
2	C	71	Total	D	O	0	0
			201	130	71		
2	D	52	Total	D	O	0	0
			148	96	52		
2	E	58	Total	D	O	0	0
			152	94	58		
2	F	64	Total	D	O	0	0
			174	110	64		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Inorganic pyrophosphatase

Chain A:  96%



- Molecule 1: Inorganic pyrophosphatase

Chain B:  97%



- Molecule 1: Inorganic pyrophosphatase

Chain C:  95%



- Molecule 1: Inorganic pyrophosphatase

Chain D:  96%



- Molecule 1: Inorganic pyrophosphatase

Chain E:  94%




- Molecule 1: Inorganic pyrophosphatase

Chain F:

97%

..



4 Model quality [i](#)

4.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DOD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	2/2393 (0.1%)	0.44	0/3237
1	B	0.26	0/2478	0.43	0/3351
1	C	0.26	0/2453	0.41	0/3321
1	D	0.27	0/2455	0.45	0/3320
1	E	0.27	0/2528	0.42	0/3424
1	F	0.25	0/2460	0.45	0/3333
All	All	0.29	2/14767 (0.0%)	0.43	0/19986

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	48[A]	SER	C-N	9.24	1.51	1.34
1	A	48[B]	SER	C-N	9.24	1.51	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1663	1337	162	1	0
1	B	1679	1368	144	1	0
1	C	1676	1362	137	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1666	1351	173	0	0
1	E	1673	1361	168	2	0
1	F	1669	1340	142	0	0
2	A	171	0	0	4	0
2	B	187	0	0	4	0
2	C	201	0	0	1	0
2	D	148	0	0	1	0
2	E	152	0	0	2	0
2	F	174	0	0	5	0
All	All	11059	8119	926	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 18 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:12:GLU:OE2	2:E:201:DOD:O	1.99	0.80
1:C:76:MET:O	2:C:201:DOD:O	2.02	0.77
1:A:90:PRO:O	2:A:202:DOD:O	2.21	0.55
1:B:167:ILE:HG23	2:B:259:DOD:O	2.06	0.51

There are no symmetry-related clashes.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/178 (154%)	264 (96%)	11 (4%)	0	100	100
1	B	286/178 (161%)	278 (97%)	8 (3%)	0	100	100
1	C	280/178 (157%)	273 (98%)	7 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	283/178 (159%)	272 (96%)	11 (4%)	0	100	100
1	E	286/178 (161%)	277 (97%)	9 (3%)	0	100	100
1	F	281/178 (158%)	276 (98%)	5 (2%)	0	100	100
All	All	1691/1068 (158%)	1640 (97%)	51 (3%)	0	100	100

There are no Ramachandran outliers to report.

4.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/159 (157%)	246 (98%)	4 (2%)	62	78
1	B	257/159 (162%)	254 (99%)	3 (1%)	71	84
1	C	255/159 (160%)	248 (97%)	7 (3%)	44	61
1	D	256/159 (161%)	249 (97%)	7 (3%)	44	61
1	E	264/159 (166%)	256 (97%)	8 (3%)	41	57
1	F	255/159 (160%)	253 (99%)	2 (1%)	81	91
All	All	1537/954 (161%)	1506 (98%)	31 (2%)	59	72

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	55[B]	ASP
1	D	70[A]	PHE
1	E	172[B]	GLU
1	D	65[A]	ASP
1	D	70[B]	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

4.3.3 RNA [i](#)

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

4.6 Ligand geometry [i](#)

There are no ligands in this entry.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.