



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 05:59 pm BST

PDB ID : 4U0L  
Title : Structure of the *Vibrio cholerae* di-nucleotide cyclase (DncV) mutant D131A-D133A  
Authors : Xiang, Y.; Zhu, D.Y.  
Deposited on : 2014-07-12  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	419	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>7%</div> <div>8%</div> </div> </div>
1	B	419	<div> <div>8%</div> <div> <div></div> <div>80%</div> <div>10%</div> <div>9%</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6579 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclic AMP-GMP synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	386	Total	C	N	O	S	0	0	0
			3088	1950	536	586	16			
1	B	383	Total	C	N	O	S	0	0	0
			3062	1936	532	578	16			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	ALA	ASP	engineered mutation	UNP Q9KVG7
A	133	ALA	ASP	engineered mutation	UNP Q9KVG7
B	131	ALA	ASP	engineered mutation	UNP Q9KVG7
B	133	ALA	ASP	engineered mutation	UNP Q9KVG7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	215	Total	O	0	0
			215	215		
2	B	214	Total	O	0	0
			214	214		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Cyclic AMP-GMP synthase



#### • Molecule 1: Cyclic AMP-GMP synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.90Å 59.77Å 104.68Å 90.00° 96.00° 90.00°	Depositor
Resolution (Å)	43.90 – 2.10 43.93 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (43.90-2.10) 95.7 (43.93-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.72 (at 2.10Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.187 , 0.233 0.188 , 0.233	Depositor DCC
$R_{free}$ test set	2571 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.1	Xtriage
Anisotropy	0.394	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 48.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6579	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.55% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/3148	0.52	0/4241
1	B	0.41	0/3122	0.54	0/4206
All	All	0.42	0/6270	0.53	0/8447

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3088	0	3091	14	0
1	B	3062	0	3070	32	0
2	A	215	0	0	1	0
2	B	214	0	0	5	0
All	All	6579	0	6161	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:139:PRO:HB3	1:B:201:LYS:HB2	1.63	0.80

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:97:ASP:OD2	1:B:98:ALA:N	2.20	0.73
1:B:96:ASP:N	1:B:96:ASP:OD1	2.21	0.72
1:B:193:ASP:OD2	2:B:696:HOH:O	2.09	0.71
1:A:59:LYS:NZ	1:A:60:SER:OG	2.28	0.67
1:B:175:GLU:OE2	2:B:672:HOH:O	2.12	0.67
1:A:97:ASP:OD1	1:A:98:ALA:N	2.29	0.64
1:A:371:GLU:OE1	2:A:700:HOH:O	2.14	0.64
1:A:208:GLN:HG2	1:B:357:TYR:HB3	1.86	0.58
1:B:140:MET:O	1:B:142:ILE:N	2.36	0.58
1:B:253:ASP:OD2	1:B:253:ASP:N	2.35	0.58
1:B:277:LYS:NZ	2:B:502:HOH:O	2.36	0.57
1:B:123:PRO:HD3	1:B:130:MET:HG3	1.86	0.56
1:B:63:THR:HG23	1:B:66:ILE:H	1.70	0.56
1:B:140:MET:HB2	1:B:200:PRO:HA	1.88	0.56
1:A:212:GLU:O	1:B:339:ARG:NH1	2.38	0.55
1:B:240:LEU:HB2	1:B:241:ASP:HA	1.87	0.55
1:A:27:GLU:O	1:A:31:ASN:HB2	2.07	0.54
1:B:200:PRO:HB2	1:B:203:GLU:HB2	1.91	0.52
1:B:154:LEU:HD21	1:B:196:MET:HB3	1.92	0.52
1:B:31:ASN:ND2	2:B:503:HOH:O	2.42	0.51
1:B:360:GLY:O	1:B:364:MET:HG2	2.10	0.51
1:B:140:MET:C	1:B:142:ILE:H	2.14	0.51
1:B:138:MET:SD	1:B:154:LEU:HD13	2.51	0.50
1:A:28:GLU:OE1	1:A:122:ARG:NH1	2.45	0.50
1:B:140:MET:N	1:B:199:ILE:O	2.45	0.49
1:A:388:GLU:OE2	1:A:391:LYS:NZ	2.30	0.49
1:B:205:GLN:NE2	2:B:505:HOH:O	2.46	0.48
1:B:141:PRO:HB3	1:B:144:GLU:OE2	2.13	0.48
1:B:98:ALA:O	1:B:102:PHE:N	2.37	0.47
1:A:95:ASP:OD1	1:A:95:ASP:N	2.42	0.46
1:A:355:PRO:HG2	1:A:358:LYS:HG3	1.96	0.46
1:B:2:ARG:HE	1:B:2:ARG:HB3	1.70	0.45
1:B:354:PRO:HG2	1:B:363:GLU:HG2	1.99	0.44
1:A:289:TRP:O	1:A:293:GLN:HG2	2.18	0.43
1:A:336:GLU:OE2	1:A:339:ARG:NH1	2.51	0.43
1:B:240:LEU:CB	1:B:241:ASP:HA	2.47	0.43
1:B:145:SER:HA	1:B:146:GLU:HA	1.66	0.43
1:B:254:ARG:NH1	1:B:257:ILE:HG22	2.36	0.41
1:B:101:GLU:HG3	1:B:144:GLU:OE1	2.20	0.41
1:A:409:LEU:HA	1:A:409:LEU:HD23	1.93	0.41
1:B:241:ASP:O	1:B:245:VAL:HG23	2.21	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145:SER:OG	1:A:146:GLU:N	2.54	0.41
1:B:289:TRP:O	1:B:293:GLN:HG2	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	382/419 (91%)	376 (98%)	5 (1%)	1 (0%)	41	41
1	B	379/419 (90%)	358 (94%)	20 (5%)	1 (0%)	41	41
All	All	761/838 (91%)	734 (96%)	25 (3%)	2 (0%)	41	41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	61	ALA
1	B	141	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	338/365 (93%)	331 (98%)	7 (2%)	53	59
1	B	335/365 (92%)	325 (97%)	10 (3%)	41	44

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	673/730 (92%)	656 (98%)	17 (2%)	47 52

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	68	GLN
1	A	85	ARG
1	A	152	SER
1	A	184	LYS
1	A	243	GLU
1	A	368	SER
1	A	372	ARG
1	B	2	ARG
1	B	70	LYS
1	B	96	ASP
1	B	100	ASP
1	B	140	MET
1	B	143	PHE
1	B	201	LYS
1	B	253	ASP
1	B	339	ARG
1	B	347	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	191	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	386/419 (92%)	0.03	10 (2%) 56 61	21, 36, 70, 95	0
1	B	383/419 (91%)	0.36	32 (8%) 11 14	22, 39, 89, 102	1 (0%)
All	All	769/838 (91%)	0.20	42 (5%) 25 31	21, 38, 82, 102	1 (0%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	240	LEU	11.3
1	B	209	ILE	11.1
1	B	211	LEU	10.6
1	B	143	PHE	9.8
1	B	210	ALA	7.9
1	B	207	LYS	6.7
1	B	206	LYS	6.6
1	B	202	ASP	5.6
1	B	212	GLU	5.6
1	B	204	PHE	5.4
1	B	241	ASP	5.4
1	A	239	GLU	5.2
1	B	253	ASP	5.2
1	B	149	ILE	4.8
1	B	242	SER	4.6
1	A	253	ASP	4.4
1	A	240	LEU	4.3
1	B	205	GLN	4.3
1	B	208	GLN	4.1
1	A	3	MET	3.8
1	A	148	LYS	3.8
1	B	144	GLU	3.7
1	B	243	GLU	3.7
1	B	146	GLU	3.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	92	TYR	3.3
1	B	145	SER	3.1
1	B	101	GLU	3.0
1	B	98	ALA	2.8
1	A	145	SER	2.8
1	A	243	GLU	2.7
1	B	2	ARG	2.6
1	B	252	GLY	2.6
1	B	142	ILE	2.5
1	B	201	LYS	2.4
1	A	61	ALA	2.4
1	B	147	PRO	2.3
1	B	148	LYS	2.2
1	A	241	ASP	2.1
1	B	251	GLU	2.0
1	A	144	GLU	2.0
1	B	95	ASP	2.0
1	B	384	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

There are no ligands in this entry.

## 6.5 Other polymers ⓘ

There are no such residues in this entry.