



Full wwPDB X-ray Structure Validation Report ⓘ

May 27, 2020 – 12:48 am BST

PDB ID : 4U0Q
Title : Plasmodium falciparum reticulocyte-binding protein homologue 5 (PfRH5) bound to basigin
Authors : Wright, K.E.; Hjerrild, K.A.; Bartlett, J.; Douglas, A.D.; Jin, J.; Brown, R.E.; Ashfield, R.; Clemmensen, S.B.; de Jongh, W.A.; Draper, S.J.; Higgins, M.K.
Deposited on : 2014-07-14
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	526	 43% 11% • 46%
1	C	526	 40% 12% • 46%
2	B	269	 2% 50% 15% • 33%
2	D	269	 % 51% 15% • 33%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7629 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reticulocyte binding protein 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	286	Total	C	N	O	S	0	0	0
			2430	1569	407	439	15			
1	C	285	Total	C	N	O	S	0	0	0
			2421	1563	405	438	15			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	216	ALA	THR	engineered mutation	UNP B2L3N7
C	216	ALA	THR	engineered mutation	UNP B2L3N7


- Molecule 2 is a protein called Basigin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	181	Total	C	N	O	S	0	0	0
			1389	862	238	281	8			
2	D	181	Total	C	N	O	S	0	0	0
			1389	862	238	281	8			


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Reticulocyte binding protein 5




• Molecule 1: Reticulocyte binding protein 5






• Molecule 2: Basigin



• Molecule 2: Basigin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	75.28Å 109.36Å 151.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	62.40 – 3.10 62.40 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (62.40-3.10) 99.7 (62.40-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 3.13Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.217 , 0.244 0.232 , 0.264	Depositor DCC
R_{free} test set	1186 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	48.0	Xtriage
Anisotropy	0.478	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 47.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	7629	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/2481	0.62	0/3327
1	C	0.41	0/2472	0.62	0/3316
2	B	0.39	0/1416	0.67	0/1918
2	D	0.38	0/1416	0.66	0/1918
All	All	0.40	0/7785	0.64	0/10479

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2430	0	2455	30	0
1	C	2421	0	2442	35	0
2	B	1389	0	1343	26	0
2	D	1389	0	1343	22	0
All	All	7629	0	7583	104	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (104) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:179:TYR:HD2	1:A:471:THR:HG22	1.14	1.11
1:A:447:TRP:HE3	2:B:28:THR:HG22	1.30	0.96
1:A:179:TYR:CD2	1:A:471:THR:HG22	2.05	0.89
1:C:479:LEU:HA	1:C:482:THR:HG23	1.66	0.77
1:C:180:THR:HG22	1:C:471:THR:HG21	1.70	0.73
2:B:54:ARG:HD3	2:B:63:LYS:HD3	1.74	0.70
1:A:447:TRP:CE3	2:B:28:THR:HG22	2.22	0.68
1:C:179:TYR:HD2	1:C:471:THR:HG22	1.57	0.68
2:B:152:ASN:HD22	2:B:160:VAL:H	1.41	0.67
2:B:54:ARG:CD	2:B:63:LYS:HD3	2.28	0.64
1:A:180:THR:HG22	1:A:471:THR:HG21	1.80	0.64
1:C:447:TRP:HE3	2:D:28:THR:HG22	1.64	0.61
1:C:467:SER:O	1:C:471:THR:HG23	2.02	0.60
1:A:446:ILE:HG23	1:A:447:TRP:CD1	2.37	0.59
1:A:192:SER:HA	1:A:343:LEU:HD12	1.85	0.59
2:D:107:VAL:HG12	2:D:128:SER:HB2	1.85	0.58
2:D:52:GLY:HA3	2:D:65:ASP:HA	1.84	0.58
1:C:446:ILE:HG23	1:C:447:TRP:CD1	2.39	0.58
2:B:107:VAL:HG12	2:B:128:SER:HB2	1.85	0.58
1:A:218:ASP:HA	1:A:221:LYS:HG2	1.87	0.56
1:A:168:GLU:OE1	1:A:240:HIS:NE2	2.39	0.56
2:B:37:ILE:HG22	2:B:76:VAL:HB	1.87	0.56
1:C:218:ASP:HA	1:C:221:LYS:HG2	1.88	0.55
2:D:28:THR:HG23	2:D:99:ILE:HG12	1.88	0.55
1:C:165:GLN:HG2	1:C:171:LEU:HD23	1.87	0.55
2:B:140:TYR:HB3	2:B:147:ASP:HB3	1.88	0.55
2:D:140:TYR:HB3	2:D:147:ASP:HB3	1.89	0.55
2:D:40:THR:HG23	2:D:73:GLU:HG2	1.89	0.54
1:A:484:TYR:HA	1:A:487:GLU:HB3	1.89	0.54
2:B:40:THR:HG23	2:B:73:GLU:HG2	1.90	0.54
1:C:368:ILE:HG13	1:C:435:THR:HG22	1.90	0.53
1:C:484:TYR:HA	1:C:487:GLU:HB3	1.90	0.53
1:A:357:ARG:HA	1:A:446:ILE:HD11	1.92	0.51
1:A:394:LEU:HA	1:A:397:LEU:HD12	1.92	0.51
1:C:357:ARG:HA	1:C:446:ILE:HD11	1.92	0.51
2:B:38:LEU:HD11	2:B:73:GLU:HB3	1.93	0.50
2:D:38:LEU:HD11	2:D:73:GLU:HB3	1.93	0.50
2:B:152:ASN:HD22	2:B:160:VAL:N	2.09	0.50
1:A:209:PHE:CE2	1:A:213:ILE:HD12	2.47	0.49
1:C:192:SER:HA	1:C:343:LEU:HD12	1.95	0.48
2:D:110:VAL:HG22	2:D:127:LYS:HE3	1.95	0.48
1:C:500:VAL:HG23	1:C:501:LEU:HD12	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:168:GLU:OE1	1:C:240:HIS:NE2	2.36	0.48
2:B:28:THR:HG23	2:B:99:ILE:HG12	1.94	0.48
2:D:182:GLN:HG2	2:D:199:THR:HG22	1.96	0.48
1:A:496:HIS:HE1	2:D:33:LEU:HD13	1.79	0.48
1:A:447:TRP:HE3	2:B:28:THR:CG2	2.15	0.48
1:C:209:PHE:CE2	1:C:213:ILE:HD12	2.49	0.47
2:B:54:ARG:HB3	2:B:88:VAL:HB	1.96	0.47
1:C:203:TYR:CE1	2:D:131:VAL:HG22	2.50	0.47
2:B:52:GLY:HA3	2:B:65:ASP:HA	1.97	0.47
2:D:54:ARG:HB3	2:D:88:VAL:HB	1.97	0.47
1:A:447:TRP:HB3	2:B:28:THR:O	2.15	0.47
2:B:110:VAL:HG22	2:B:127:LYS:HE3	1.95	0.47
2:B:182:GLN:HG2	2:B:199:THR:HG22	1.96	0.47
2:B:186:ASN:HD22	2:B:195:GLN:HG2	1.80	0.47
2:D:122:ALA:HB2	2:D:174:LEU:HD11	1.97	0.47
2:D:186:ASN:HD22	2:D:195:GLN:HG2	1.81	0.46
2:B:188:THR:HG23	2:B:193:SER:HB3	1.98	0.46
1:C:498:ILE:HG23	1:C:503:MET:HB2	1.98	0.46
1:C:353:THR:HG22	1:C:356:ILE:HD12	1.98	0.46
2:B:43:LEU:HD22	2:B:46:SER:HB2	1.98	0.46
1:A:467:SER:O	1:A:471:THR:HG23	2.16	0.45
1:C:404:TYR:HD2	1:C:407:ILE:HG13	1.81	0.45
2:D:176:MET:HG2	2:D:202:VAL:HG13	1.99	0.45
1:A:173:PHE:CZ	1:A:221:LYS:HB2	2.52	0.45
1:A:438:ILE:O	1:A:442:ILE:HG13	2.17	0.45
2:D:188:THR:HG23	2:D:193:SER:HB3	1.97	0.45
1:C:173:PHE:CZ	1:C:221:LYS:HB2	2.51	0.45
1:A:188:LEU:HD21	1:A:460:LEU:HD23	1.99	0.45
1:A:168:GLU:OE1	1:A:489:HIS:NE2	2.33	0.45
2:D:43:LEU:HD22	2:D:46:SER:HB2	1.98	0.44
1:C:179:TYR:CD2	1:C:471:THR:HG22	2.45	0.44
2:D:142:ILE:HD11	2:D:184:ARG:HB2	1.98	0.44
1:A:200:TYR:HB3	2:B:131:VAL:CG2	2.48	0.44
1:C:229:ASN:HA	1:C:232:ILE:HD12	2.00	0.44
1:C:438:ILE:O	1:C:442:ILE:HG13	2.17	0.44
1:C:334:ASN:O	1:C:338:ASN:HB2	2.18	0.43
1:C:197:SER:OG	2:D:100:GLN:NE2	2.37	0.43
1:A:217:TYR:HA	1:A:328:ILE:HD13	1.99	0.43
1:A:163:ILE:HG23	1:A:171:LEU:HD21	2.00	0.43
1:A:306:GLU:HG2	1:A:310:LYS:HE3	2.00	0.43
1:C:163:ILE:HG23	1:C:171:LEU:HD21	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:306:GLU:HG2	1:C:310:LYS:HE3	2.00	0.43
1:A:353:THR:HG22	1:A:356:ILE:HD12	2.01	0.42
2:B:176:MET:HG2	2:B:202:VAL:HG13	2.00	0.42
1:C:478:MET:O	1:C:482:THR:CG2	2.67	0.42
1:C:347:ASN:HD22	1:C:349:ASN:H	1.67	0.42
2:B:125:VAL:HG22	2:B:168:GLU:HG2	2.02	0.42
1:A:185:TYR:CZ	1:A:333:LYS:HG3	2.54	0.42
1:C:314:LEU:HD22	1:C:318:ILE:HD11	2.02	0.42
2:D:39:LEU:HB3	2:D:99:ILE:HD13	2.00	0.42
2:B:142:ILE:HD11	2:B:184:ARG:HB2	2.01	0.42
2:D:30:VAL:HG23	2:D:39:LEU:HD23	2.03	0.41
1:C:185:TYR:CZ	1:C:333:LYS:HG3	2.56	0.41
1:A:229:ASN:HA	1:A:232:ILE:HD12	2.01	0.41
1:C:168:GLU:OE1	1:C:489:HIS:NE2	2.34	0.41
1:C:193:ILE:HD13	1:C:206:VAL:HG21	2.03	0.41
1:A:383:MET:HG2	1:A:483:PHE:CE2	2.56	0.41
1:C:478:MET:O	1:C:482:THR:HG22	2.21	0.41
1:C:479:LEU:CA	1:C:482:THR:HG23	2.43	0.40
2:B:184:ARG:HG3	2:B:197:ILE:HG13	2.03	0.40
1:A:180:THR:CG2	1:A:471:THR:HG21	2.50	0.40
2:D:184:ARG:HG3	2:D:197:ILE:HG13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/526 (54%)	274 (97%)	6 (2%)	2 (1%)	22	57
1	C	281/526 (53%)	276 (98%)	5 (2%)	0	100	100
2	B	179/269 (66%)	165 (92%)	14 (8%)	0	100	100
2	D	179/269 (66%)	167 (93%)	12 (7%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	921/1590 (58%)	882 (96%)	37 (4%)	2 (0%)	47 79

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	241	PRO
1	A	399	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	277/511 (54%)	253 (91%)	24 (9%)	10 36
1	C	276/511 (54%)	246 (89%)	30 (11%)	6 25
2	B	156/225 (69%)	141 (90%)	15 (10%)	8 31
2	D	156/225 (69%)	144 (92%)	12 (8%)	13 41
All	All	865/1472 (59%)	784 (91%)	81 (9%)	8 32

All (81) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	162	ASP
1	A	163	ILE
1	A	164	LEU
1	A	165	GLN
1	A	168	GLU
1	A	213	ILE
1	A	230	ASP
1	A	305	ASP
1	A	314	LEU
1	A	345	CYS
1	A	347	ASN
1	A	368	ILE
1	A	418	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	419	HIS
1	A	446	ILE
1	A	451	GLN
1	A	453	ASP
1	A	455	LEU
1	A	458	ARG
1	A	465	GLU
1	A	477	GLN
1	A	487	GLU
1	A	490	LEU
1	A	497	LEU
2	B	37	ILE
2	B	40	THR
2	B	43	LEU
2	B	48	THR
2	B	49	GLU
2	B	63	LYS
2	B	64	GLU
2	B	70	GLN
2	B	94	MET
2	B	106	ARG
2	B	141	LYS
2	B	144	ASP
2	B	169	LEU
2	B	175	ASN
2	B	176	MET
1	C	162	ASP
1	C	163	ILE
1	C	164	LEU
1	C	168	GLU
1	C	193	ILE
1	C	213	ILE
1	C	230	ASP
1	C	242	TYR
1	C	314	LEU
1	C	322	GLU
1	C	345	CYS
1	C	347	ASN
1	C	368	ILE
1	C	392	LEU
1	C	393	LEU
1	C	394	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	399	LYS
1	C	400	LYS
1	C	418	LYS
1	C	435	THR
1	C	446	ILE
1	C	451	GLN
1	C	453	ASP
1	C	458	ARG
1	C	465	GLU
1	C	477	GLN
1	C	482	THR
1	C	486	LYS
1	C	487	GLU
1	C	490	LEU
2	D	31	GLU
2	D	37	ILE
2	D	40	THR
2	D	43	LEU
2	D	48	THR
2	D	64	GLU
2	D	70	GLN
2	D	94	MET
2	D	144	ASP
2	D	169	LEU
2	D	173	ASN
2	D	176	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	321	HIS
1	A	347	ASN
1	A	439	GLN
1	A	496	HIS
2	B	115	HIS
2	B	152	ASN
2	B	186	ASN
1	C	321	HIS
1	C	347	ASN
1	C	439	GLN
1	C	496	HIS
2	D	100	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	102	HIS
2	D	186	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	286/526 (54%)	-0.12	0	100	100	26, 43, 77, 116	0
1	C	285/526 (54%)	-0.21	0	100	100	25, 36, 55, 96	0
2	B	181/269 (67%)	0.15	6 (3%)	46	24	23, 57, 97, 111	0
2	D	181/269 (67%)	0.32	4 (2%)	62	41	39, 66, 108, 128	0
All	All	933/1590 (58%)	-0.01	10 (1%)	80	64	23, 45, 93, 128	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	65	ASP	4.3
2	D	66	ALA	4.3
2	B	66	ALA	4.3
2	B	183	TYR	3.8
2	B	65	ASP	3.3
2	D	121	THR	3.0
2	D	144	ASP	2.6
2	B	148	LYS	2.6
2	B	134	VAL	2.4
2	B	121	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.