



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 30, 2020 – 10:27 AM BST

PDB ID : 4U0Z
Title : Eukaryotic Fic Domain containing protein with bound APCPP
Authors : Cole, A.R.; Bunney, T.D.; Katan, M.
Deposited on : 2014-07-14
Resolution : 2.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3104 (3.00-2.92)
Clashscore	141614	3462 (3.00-2.92)
Ramachandran outliers	138981	3340 (3.00-2.92)
Sidechain outliers	138945	3343 (3.00-2.92)
RSRZ outliers	127900	2986 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 18%, green 74%, grey 1%);"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> % 74% 18% • 6% </div> </div>
1	B	344	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 1%, yellow 17%, green 76%, grey 1%);"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> 2% 76% 17% • 5% </div> </div>
1	C	344	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, yellow 18%, green 77%, grey 1%);"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> 77% 18% • • </div> </div>
1	D	344	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 18%, green 75%, grey 1%);"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> % 75% 18% • 6% </div> </div>
1	E	344	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, yellow 17%, green 76%, orange 1%, grey 1%);"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> 76% 17% • 5% </div> </div>
1	F	344	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 3%, orange 1%, yellow 16%, green 76%, grey 1%);"></div> <div style="display: flex; justify-content: space-between; font-size: 0.8em;"> 3% 76% 16% • 6% </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	344	<div><div><div>%</div><div><div></div><div>77%</div><div>16%</div><div>• 6%</div></div></div></div>
1	H	344	<div><div><div>2%</div><div><div></div><div>73%</div><div>19%</div><div>• 8%</div></div></div></div>

2 Entry composition


There are 4 unique types of molecules in this entry. The entry contains 21328 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenosine monophosphate-protein transferase FICD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	322	Total	C	N	O	S	0	0	0
			2541	1622	438	472	9			
1	B	326	Total	C	N	O	S	0	0	0
			2528	1617	432	469	10			
1	C	329	Total	C	N	O	S	0	0	0
			2581	1649	444	476	12			
1	D	325	Total	C	N	O	S	0	0	0
			2528	1616	436	466	10			
1	E	327	Total	C	N	O	S	0	0	0
			2568	1640	441	476	11			
1	F	322	Total	C	N	O	S	0	0	0
			2502	1601	428	461	12			
1	G	322	Total	C	N	O	S	0	0	0
			2509	1607	428	462	12			
1	H	318	Total	C	N	O	S	0	0	0
			2499	1599	429	460	11			

- Molecule 2 is DIPHOSPHOMETHYLPHOSPHONIC ACID ADENOSYL ESTER (three-letter code: APC) (formula: C₁₁H₁₈N₅O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	11	5	12	3		
2	B	1	Total	C	N	O	P	0	0
			31	11	5	12	3		
2	C	1	Total	C	N	O	P	0	0
			31	11	5	12	3		
2	D	1	Total	C	N	O	P	0	0
			31	11	5	12	3		
2	E	1	Total	C	N	O	P	0	0
			31	11	5	12	3		
2	F	1	Total	C	N	O	P	0	0
			31	11	5	12	3		
2	G	1	Total	C	N	O	P	0	0
			31	11	5	12	3		
2	H	1	Total	C	N	O	P	0	0
			31	11	5	12	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		
3	E	1	Total	Mg	0	0
			1	1		
3	H	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Mg 1	0	0
3	C	1	Total 1	Mg 1	0	0
3	A	1	Total 1	Mg 1	0	0
3	F	1	Total 1	Mg 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	96	Total 96	O 96	0	0
4	B	101	Total 101	O 101	0	0
4	C	114	Total 114	O 114	0	0
4	D	86	Total 86	O 86	0	0
4	E	96	Total 96	O 96	0	0
4	F	96	Total 96	O 96	0	0
4	G	128	Total 128	O 128	0	0
4	H	99	Total 99	O 99	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Adenosine monophosphate-protein transferase FICD




- Molecule 1: Adenosine monophosphate-protein transferase FICD





- Molecule 1: Adenosine monophosphate-protein transferase FICD








• Molecule 1: Adenosine monophosphate-protein transferase FICD




• Molecule 1: Adenosine monophosphate-protein transferase FICD




• Molecule 1: Adenosine monophosphate-protein transferase FICD




• Molecule 1: Adenosine monophosphate-protein transferase FICD





• Molecule 1: Adenosine monophosphate-protein transferase FICD



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	77.10Å 83.75Å 130.02Å 89.92° 89.57° 89.43°	Depositor
Resolution (Å)	43.34 – 2.95 43.34 – 2.95	Depositor EDS
% Data completeness (in resolution range)	95.0 (43.34-2.95) 94.9 (43.34-2.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 2.95Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.206 , 0.251 0.209 , 0.235	Depositor DCC
R_{free} test set	3307 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	47.8	Xtriage
Anisotropy	0.873	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 68.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.417 for h,-k,-l 0.387 for -h,k,-l 0.387 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	21328	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 37.12 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.5002e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: APC, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/2596	0.69	0/3529
1	B	0.48	0/2583	0.67	0/3514
1	C	0.53	0/2638	0.72	1/3584 (0.0%)
1	D	0.49	0/2583	0.67	0/3512
1	E	0.54	0/2625	0.70	0/3568
1	F	0.49	0/2557	0.68	0/3477
1	G	0.54	0/2564	0.69	0/3487
1	H	0.50	0/2553	0.67	0/3468
All	All	0.51	0/20699	0.69	1/28139 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	209	PRO	C-N-CA	6.10	136.95	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2541	0	2457	28	0
1	B	2528	0	2412	28	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2581	0	2503	28	0
1	D	2528	0	2419	34	0
1	E	2568	0	2476	26	0
1	F	2502	0	2383	24	0
1	G	2509	0	2407	27	0
1	H	2499	0	2403	30	0
2	A	31	0	14	0	0
2	B	31	0	14	0	0
2	C	31	0	14	0	0
2	D	31	0	14	0	0
2	E	31	0	14	1	0
2	F	31	0	14	0	0
2	G	31	0	14	1	0
2	H	31	0	14	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
4	A	96	0	0	6	0
4	B	101	0	0	3	0
4	C	114	0	0	3	0
4	D	86	0	0	6	0
4	E	96	0	0	3	0
4	F	96	0	0	0	0
4	G	128	0	0	3	0
4	H	99	0	0	1	0
All	All	21328	0	19572	216	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 216 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:209:PRO:HG3	1:C:212:ASN:HB2	1.53	0.90
1:G:141:ASP:HA	4:G:643:HOH:O	1.79	0.81
1:B:258:LEU:HD13	1:H:265:ILE:HD11	1.63	0.78

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:158:GLN:HG3	4:E:626:HOH:O	1.91	0.70
1:G:342:GLU:OE2	4:G:661:HOH:O	2.13	0.67

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/344 (92%)	310 (98%)	7 (2%)	1 (0%)	41	73
1	B	322/344 (94%)	313 (97%)	7 (2%)	2 (1%)	25	60
1	C	327/344 (95%)	315 (96%)	7 (2%)	5 (2%)	10	38
1	D	321/344 (93%)	311 (97%)	8 (2%)	2 (1%)	25	60
1	E	325/344 (94%)	315 (97%)	8 (2%)	2 (1%)	25	60
1	F	318/344 (92%)	312 (98%)	4 (1%)	2 (1%)	25	60
1	G	318/344 (92%)	311 (98%)	6 (2%)	1 (0%)	41	73
1	H	312/344 (91%)	307 (98%)	4 (1%)	1 (0%)	41	73
All	All	2561/2752 (93%)	2494 (97%)	51 (2%)	16 (1%)	25	60

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	208	ILE
1	C	209	PRO
1	C	210	LYS
1	C	213	SER
1	E	213	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/302 (87%)	241 (92%)	22 (8%)	11	35
1	B	255/302 (84%)	236 (92%)	19 (8%)	13	40
1	C	267/302 (88%)	245 (92%)	22 (8%)	11	36
1	D	255/302 (84%)	235 (92%)	20 (8%)	12	38
1	E	264/302 (87%)	237 (90%)	27 (10%)	7	25
1	F	251/302 (83%)	231 (92%)	20 (8%)	12	37
1	G	255/302 (84%)	234 (92%)	21 (8%)	11	36
1	H	255/302 (84%)	234 (92%)	21 (8%)	11	36
All	All	2065/2416 (86%)	1893 (92%)	172 (8%)	11	35

5 of 172 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	359	LEU
1	E	220	GLU
1	H	206	MET
1	D	389	ILE
1	E	133	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 58 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	323	HIS
1	E	225	HIS
1	H	229	HIS
1	D	347	HIS
1	D	401	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	APC	F	501	-	27,33,33	1.68	3 (11%)	31,52,52	1.29	5 (16%)
2	APC	B	501	-	27,33,33	1.54	3 (11%)	31,52,52	1.25	4 (12%)
2	APC	C	501	-	27,33,33	1.59	3 (11%)	31,52,52	1.57	4 (12%)
2	APC	H	501	-	27,33,33	1.61	4 (14%)	31,52,52	1.55	4 (12%)
2	APC	E	501	3	27,33,33	1.68	3 (11%)	31,52,52	1.35	4 (12%)
2	APC	G	501	3	27,33,33	1.65	3 (11%)	31,52,52	1.56	3 (9%)
2	APC	A	501	-	27,33,33	1.47	3 (11%)	31,52,52	1.30	5 (16%)
2	APC	D	501	-	27,33,33	1.69	3 (11%)	31,52,52	1.53	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	APC	F	501	-	-	2/15/38/38	0/3/3/3
2	APC	B	501	-	-	2/15/38/38	0/3/3/3
2	APC	C	501	-	-	2/15/38/38	0/3/3/3
2	APC	H	501	-	-	1/15/38/38	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	APC	E	501	3	-	2/15/38/38	0/3/3/3
2	APC	G	501	3	-	1/15/38/38	0/3/3/3
2	APC	A	501	-	-	2/15/38/38	0/3/3/3
2	APC	D	501	-	-	1/15/38/38	0/3/3/3

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	501	APC	PA-O5'	7.18	1.68	1.57
2	E	501	APC	PA-O5'	6.99	1.67	1.57
2	G	501	APC	PA-O5'	6.98	1.67	1.57
2	F	501	APC	PA-O5'	6.95	1.67	1.57
2	H	501	APC	PA-O5'	6.45	1.67	1.57

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	501	APC	O1B-PB-C3A	5.40	123.35	109.07
2	G	501	APC	O1B-PB-C3A	5.20	122.82	109.07
2	D	501	APC	O1B-PB-C3A	5.05	122.43	109.07
2	C	501	APC	O1B-PB-C3A	5.00	122.29	109.07
2	C	501	APC	O2G-PG-O3B	4.42	119.44	104.64

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	501	APC	PB-C3A-PA-O2A
2	B	501	APC	PB-C3A-PA-O2A
2	E	501	APC	PB-C3A-PA-O2A
2	A	501	APC	PB-C3A-PA-O2A
2	F	501	APC	PB-C3A-PA-O1A


There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	501	APC	1	0
2	G	501	APC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,


bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand APC C 501




Ligand APC H 501




Ligand APC E 501



Ligand APC G 501





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	322/344 (93%)	-0.33	2 (0%) 89 78	16, 48, 94, 121	0
1	B	326/344 (94%)	-0.02	6 (1%) 68 51	38, 71, 116, 126	0
1	C	329/344 (95%)	-0.32	1 (0%) 94 87	16, 47, 97, 118	0
1	D	325/344 (94%)	-0.05	5 (1%) 73 57	40, 69, 111, 138	0
1	E	327/344 (95%)	-0.36	1 (0%) 94 87	20, 46, 95, 114	0
1	F	322/344 (93%)	0.10	10 (3%) 49 32	42, 72, 120, 144	0
1	G	322/344 (93%)	-0.16	5 (1%) 72 55	14, 49, 97, 112	0
1	H	318/344 (92%)	-0.03	6 (1%) 66 49	40, 65, 109, 129	0
All	All	2591/2752 (94%)	-0.15	36 (1%) 75 59	14, 60, 108, 144	0

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	132	ALA	5.1
1	D	214	ALA	4.3
1	D	432	ALA	4.2
1	G	205	VAL	4.1
1	B	109	ALA	3.7

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	A	502	1/1	0.78	0.16	71,71,71,71	0
3	MG	H	502	1/1	0.86	0.12	64,64,64,64	0
3	MG	D	502	1/1	0.88	0.09	86,86,86,86	0
2	APC	F	501	31/31	0.94	0.16	62,82,94,95	0
2	APC	H	501	31/31	0.94	0.19	80,92,95,97	0
3	MG	E	502	1/1	0.94	0.07	59,59,59,59	0
2	APC	D	501	31/31	0.94	0.20	79,92,96,97	0
3	MG	F	502	1/1	0.95	0.20	53,53,53,53	0
3	MG	B	502	1/1	0.96	0.14	52,52,52,52	0
2	APC	B	501	31/31	0.96	0.16	65,97,104,105	0
2	APC	E	501	31/31	0.96	0.18	45,63,72,74	0
2	APC	G	501	31/31	0.96	0.17	40,59,72,75	0
2	APC	C	501	31/31	0.97	0.16	33,55,63,66	0
3	MG	G	502	1/1	0.97	0.14	48,48,48,48	0
3	MG	C	502	1/1	0.97	0.16	32,32,32,32	0
2	APC	A	501	31/31	0.97	0.16	44,60,64,65	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around APC F 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around APC H 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around APC D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around APC B 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around APC E 501:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around APC G 501:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around APC C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around APC A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.