



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 07:11 pm BST

PDB ID : 6U00
Title : Crystal Structure of Fungal RNA Kinase
Authors : Shuman, S.; Goldgur, Y.; Banerjee, A.
Deposited on : 2019-08-13
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	432	
1	B	432	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	B	901	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3629 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	211	Total	C	N	O	S	0	0	0
			1730	1094	290	342	4			
1	B	210	Total	C	N	O	S	0	0	0
			1721	1089	288	340	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	543	LEU	SER	conflict	UNP P43075
A	587	LEU	SER	conflict	UNP P43075
B	543	LEU	SER	conflict	UNP P43075
B	587	LEU	SER	conflict	UNP P43075

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	101	Total	O	0	0
			101	101		
3	B	67	Total	O	0	0
			67	67		

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	55.44Å 122.87Å 150.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.91 – 1.98 47.91 – 1.98	Depositor EDS
% Data completeness (in resolution range)	95.8 (47.91-1.98) 91.8 (47.91-1.98)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.05 (at 1.98Å)	Xtriage
Refinement program	PHENIX 1.15.2 _3472	Depositor
R, R_{free}	0.193 , 0.236 0.193 , 0.236	Depositor DCC
R_{free} test set	2000 reflections (5.76%)	wwPDB-VP
Wilson B-factor (Å ²)	36.5	Xtriage
Anisotropy	0.669	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 55.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3629	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/1759	0.67	2/2383 (0.1%)
1	B	0.47	0/1750	0.83	8/2371 (0.3%)
All	All	0.46	0/3509	0.75	10/4754 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	558	ARG	NE-CZ-NH2	-15.19	112.70	120.30
1	B	558	ARG	NE-CZ-NH1	10.02	125.31	120.30
1	B	484	ARG	NE-CZ-NH2	-9.07	115.76	120.30
1	B	484	ARG	NE-CZ-NH1	8.12	124.36	120.30
1	A	553	LYS	CA-CB-CG	-6.81	98.41	113.40
1	B	558	ARG	CD-NE-CZ	6.07	132.10	123.60
1	B	484	ARG	CB-CG-CD	-6.00	96.00	111.60
1	B	547	LEU	CA-CB-CG	5.71	128.42	115.30
1	B	501	THR	CB-CA-C	-5.19	97.58	111.60
1	A	553	LYS	CB-CG-CD	5.13	124.94	111.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	628	TYR	Peptide
1	B	583	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1730	0	1714	23	1
1	B	1721	0	1706	26	1
2	A	5	0	0	0	0
2	B	5	0	0	2	0
3	A	101	0	0	3	0
3	B	67	0	0	3	0
All	All	3629	0	3420	50	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:577:HIS:ND1	3:B:1002:HOH:O	2.15	0.78
1:B:581:SER:HB2	1:B:583:ASP:HB2	1.69	0.74
1:A:579:LYS:NZ	1:A:584:GLU:OE1	2.23	0.70
2:B:901:PO4:O3	3:B:1001:HOH:O	2.10	0.67
1:A:482:GLU:OE1	3:A:1001:HOH:O	2.13	0.67
1:B:460:LEU:HD21	1:B:490:ILE:HG23	1.81	0.62
1:A:522:TRP:CD2	1:A:553:LYS:HE3	2.35	0.61
1:B:522:TRP:HE3	1:B:556:ILE:HD12	1.65	0.61
1:B:545:GLU:OE1	1:B:545:GLU:HA	2.03	0.59
1:B:499:ASP:OD1	1:B:500:ASP:N	2.37	0.57
1:A:528:ARG:HG2	1:A:586:SER:HB2	1.87	0.57
1:B:581:SER:CB	1:B:583:ASP:HB2	2.34	0.56
1:B:521:LEU:O	1:B:525:THR:OG1	2.17	0.55
1:A:522:TRP:CG	1:A:553:LYS:HE3	2.43	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:443:GLN:HA	1:B:474:ASP:HB3	1.90	0.52
1:A:547:LEU:HD22	1:A:547:LEU:H	1.77	0.49
1:B:526:TYR:O	1:B:530:ILE:HG12	2.14	0.48
1:B:581:SER:HB2	1:B:583:ASP:CB	2.42	0.48
1:B:524:ILE:O	1:B:528:ARG:HG2	2.14	0.48
1:A:425:LYS:HE3	1:A:476:ASN:OD1	2.14	0.48
1:B:545:GLU:O	1:B:549:GLU:HG3	2.14	0.48
1:B:528:ARG:HD3	1:B:586:SER:HB2	1.95	0.47
1:B:591:LEU:O	1:B:595:ARG:HG3	2.13	0.47
2:B:901:PO4:P	3:B:1001:HOH:O	2.73	0.47
1:A:443:GLN:HA	1:A:474:ASP:HB3	1.97	0.46
1:B:421:ILE:HD12	1:B:421:ILE:HA	1.70	0.46
1:A:579:LYS:HB3	1:A:579:LYS:HE3	1.62	0.45
1:A:600:ASP:OD2	1:B:603:GLN:HG2	2.17	0.45
1:A:579:LYS:NZ	1:A:589:SER:HB2	2.32	0.45
1:A:591:LEU:O	1:A:595:ARG:HG3	2.17	0.45
1:B:587:LEU:HD22	1:B:626:LEU:HD21	1.98	0.44
1:B:582:LYS:HG3	1:B:583:ASP:N	2.32	0.44
1:A:421:ILE:HD11	1:A:552:MET:HB3	2.00	0.43
1:A:547:LEU:N	1:A:547:LEU:HD22	2.34	0.43
1:B:582:LYS:HG3	1:B:583:ASP:H	1.83	0.43
1:A:479:ALA:HB2	1:A:558:ARG:HG3	2.02	0.42
1:B:553:LYS:HB3	1:B:553:LYS:HE3	1.71	0.42
1:B:544:ASP:HA	1:B:547:LEU:HD23	2.01	0.42
1:B:478:SER:O	1:B:483:ARG:NH2	2.53	0.42
1:A:612:LYS:HE3	1:A:612:LYS:HB3	1.91	0.42
1:A:543:LEU:HD12	3:A:1021:HOH:O	2.20	0.42
1:A:548:VAL:HG13	1:A:552:MET:SD	2.60	0.41
1:B:516:LEU:HA	1:B:516:LEU:HD23	1.93	0.41
1:A:443:GLN:NE2	3:A:1003:HOH:O	2.32	0.41
1:A:582:LYS:HG3	1:A:583:ASP:H	1.86	0.41
1:A:563:ASN:ND2	1:A:566:ARG:HE	2.20	0.40
1:B:483:ARG:HD2	1:B:569:ASP:OD1	2.21	0.40
1:B:558:ARG:NE	1:B:558:ARG:HA	2.36	0.40
1:A:444:ASN:HA	1:A:447:ILE:HD12	2.02	0.40
1:A:558:ARG:O	1:A:558:ARG:HG3	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:558:ARG:NH1	1:B:500:ASP:OD1[4_555]	1.46	0.74

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/432 (48%)	202 (98%)	5 (2%)	0	100	100
1	B	206/432 (48%)	202 (98%)	4 (2%)	0	100	100
All	All	413/864 (48%)	404 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	201/400 (50%)	197 (98%)	4 (2%)	55	48
1	B	200/400 (50%)	197 (98%)	3 (2%)	65	59
All	All	401/800 (50%)	394 (98%)	7 (2%)	60	53

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	450	LYS
1	A	544	ASP
1	A	589	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	624	LYS
1	B	449	LYS
1	B	555	PHE
1	B	582	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	901	-	4,4,4	0.74	0	6,6,6	0.48	0
2	PO4	B	901	-	4,4,4	0.77	0	6,6,6	0.63	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	901	PO4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	211/432 (48%)	1.04	32 (15%) 2 2	31, 44, 117, 141	0
1	B	210/432 (48%)	0.91	26 (12%) 4 4	33, 54, 117, 131	0
All	All	421/864 (48%)	0.97	58 (13%) 2 3	31, 50, 117, 141	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	628	TYR	11.3
1	B	630	PRO	10.2
1	B	543	LEU	7.2
1	A	628	TYR	7.0
1	A	555	PHE	6.9
1	A	409	THR	6.7
1	B	552	MET	6.7
1	B	526	TYR	6.2
1	A	548	VAL	5.9
1	B	529	VAL	5.8
1	A	544	ASP	5.7
1	A	549	GLU	5.6
1	B	530	ILE	5.6
1	A	543	LEU	5.6
1	B	547	LEU	5.5
1	A	526	TYR	5.3
1	A	630	PRO	5.2
1	A	530	ILE	5.2
1	A	545	GLU	5.1
1	A	520	GLU	5.0
1	A	550	SER	4.8
1	A	558	ARG	4.8
1	B	548	VAL	4.7
1	B	515	ASP	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	629	LYS	4.3
1	A	522	TRP	4.3
1	B	583	ASP	4.0
1	B	551	VAL	3.9
1	A	557	GLN	3.7
1	A	552	MET	3.6
1	A	551	VAL	3.5
1	A	547	LEU	3.5
1	B	627	ASP	3.5
1	B	409	THR	3.5
1	B	555	PHE	3.3
1	A	583	ASP	3.3
1	A	450	LYS	3.2
1	B	516	LEU	3.1
1	B	558	ARG	3.1
1	B	524	ILE	2.8
1	A	519	GLU	2.8
1	A	410	THR	2.7
1	B	522	TRP	2.7
1	A	554	GLY	2.7
1	A	529	VAL	2.4
1	B	582	LYS	2.4
1	B	550	SER	2.3
1	B	546	ASN	2.3
1	A	562	ILE	2.2
1	A	627	ASP	2.2
1	B	580	LEU	2.2
1	A	553	LYS	2.2
1	A	546	ASN	2.1
1	A	523	ASP	2.1
1	A	531	GLN	2.1
1	A	418	ILE	2.1
1	B	519	GLU	2.0
1	B	544	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	PO4	A	901	5/5	0.66	0.20	73,82,98,99	0
2	PO4	B	901	5/5	0.86	0.12	95,95,99,99	0

6.5 Other polymers [i](#)

There are no such residues in this entry.