



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 01:57 am BST

PDB ID : 3U25  
Title : Crystal structure of P. aeruginosa azurin containing a Tyr-His hydrogen bonded pair  
Authors : Warren, J.J.; Winkler, J.R.; Gray, H.B.  
Deposited on : 2011-09-30  
Resolution : 1.18 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.18 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1123 (1.20-1.16)
Clashscore	141614	1182 (1.20-1.16)
Ramachandran outliers	138981	1134 (1.20-1.16)
Sidechain outliers	138945	1134 (1.20-1.16)
RSRZ outliers	127900	1102 (1.20-1.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	127	
1	B	127	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2209 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Azurin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	127	Total	C	N	O	S	0	4	0
			988	618	165	195	10			
1	B	127	Total	C	N	O	S	0	3	0
			983	615	164	194	10			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	HIS	ILE	ENGINEERED MUTATION	UNP P00282
A	48	TYR	TRP	ENGINEERED MUTATION	UNP P00282
A	72	PHE	TYR	ENGINEERED MUTATION	UNP P00282
A	108	PHE	TYR	ENGINEERED MUTATION	UNP P00282
B	20	HIS	ILE	ENGINEERED MUTATION	UNP P00282
B	48	TYR	TRP	ENGINEERED MUTATION	UNP P00282
B	72	PHE	TYR	ENGINEERED MUTATION	UNP P00282
B	108	PHE	TYR	ENGINEERED MUTATION	UNP P00282

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Cu	0	0
			1	1		
2	A	1	Total	Cu	0	0
			1	1		

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			8	4	1	3		
3	B	1	Total	C	N	O	0	0
			8	4	1	3		



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	114	Total	O	0	0
			114	114		
4	B	106	Total	O	0	0
			106	106		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Azurin



#### • Molecule 1: Azurin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.03Å 56.33Å 82.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.18 19.73 – 1.18	Depositor EDS
% Data completeness (in resolution range)	94.9 (20.00-1.18) 94.9 (19.73-1.18)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.80 (at 1.18Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.195 , 0.217 0.192 , 0.214	Depositor DCC
$R_{free}$ test set	3566 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	11.7	Xtriage
Anisotropy	0.113	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 47.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2209	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	12.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.28	3/1017 (0.3%)	1.22	3/1367 (0.2%)
1	B	1.23	1/1009 (0.1%)	1.20	8/1356 (0.6%)
All	All	1.25	4/2026 (0.2%)	1.21	11/2723 (0.4%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	2[A]	GLU	CD-OE1	9.48	1.36	1.25
1	A	2[B]	GLU	CD-OE1	9.48	1.36	1.25
1	A	13	MET	SD-CE	-6.25	1.42	1.77
1	B	105	GLY	N-CA	5.14	1.53	1.46

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	69	ASP	CB-CG-OD1	12.99	129.99	118.30
1	B	11	ASP	CB-CG-OD1	6.73	124.36	118.30
1	B	70	LYS	CD-CE-NZ	6.73	127.18	111.70
1	A	91	GLU	OE1-CD-OE2	-6.43	115.58	123.30
1	B	79	ARG	NE-CZ-NH1	-6.33	117.14	120.30
1	B	71	ASP	CB-CG-OD2	-6.21	112.71	118.30
1	B	69	ASP	CB-CG-OD1	-5.61	113.25	118.30
1	B	91	GLU	OE1-CD-OE2	-5.59	116.59	123.30
1	B	71	ASP	CB-CG-OD1	5.50	123.25	118.30
1	B	127	LEU	CB-CA-C	-5.45	99.85	110.20
1	A	69	ASP	OD1-CG-OD2	-5.03	113.74	123.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	988	0	970	26	2
1	B	983	0	964	13	2
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	8	0	12	2	2
3	B	8	0	12	1	2
4	A	114	0	0	0	0
4	B	106	0	0	1	0
All	All	2209	0	1958	37	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:83:HIS:NE2	3:B:1:TRS:O3	1.80	1.14
1:A:83:HIS:NE2	3:A:129:TRS:O1	1.81	1.11
1:A:24:LYS:HG2	1:A:127:LEU:CD1	1.83	1.06
1:A:24:LYS:HG2	1:A:127:LEU:HD11	1.06	1.04
1:A:24:LYS:CG	1:A:127:LEU:HD11	1.94	0.98
1:A:24:LYS:CG	1:A:127:LEU:CD1	2.48	0.92
1:A:103:LYS:HG3	1:A:105:GLY:H	1.40	0.86
1:A:13:MET:HE2	1:A:120:LEU:HD12	1.57	0.85
1:B:103:LYS:HB3	4:B:266:HOH:O	1.79	0.83
1:B:104:GLU:HA	1:B:127:LEU:HD12	1.68	0.75
1:A:103:LYS:HG3	1:A:105:GLY:N	2.14	0.61
1:A:68[A]:LEU:HD23	1:A:72:PHE:CE2	2.35	0.61
1:A:13:MET:HE2	1:A:120:LEU:CD1	2.31	0.60
1:A:24:LYS:HG3	1:A:127:LEU:HD13	1.84	0.60
1:A:24:LYS:CG	1:A:127:LEU:HD13	2.32	0.59
1:A:18[B]:ASN:ND2	1:B:18:ASN:H	2.04	0.55
1:A:13:MET:CE	1:A:120:LEU:HD12	2.35	0.52
1:A:24:LYS:HG3	1:A:127:LEU:CD1	2.36	0.52
1:A:24:LYS:HZ2	1:A:104:GLU:HG3	1.75	0.51

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:HIS:CE1	3:A:129:TRS:O1	2.61	0.51
1:A:127:LEU:C	1:A:127:LEU:HD13	2.30	0.51
1:A:68[B]:LEU:HD13	1:A:72:PHE:CE2	2.46	0.51
1:B:125:LEU:HD23	1:B:125:LEU:C	2.32	0.50
1:A:125:LEU:C	1:A:125:LEU:HD23	2.32	0.49
1:A:68[A]:LEU:HD23	1:A:72:PHE:HE2	1.77	0.49
1:B:24:LYS:HD3	1:B:104:GLU:CD	2.33	0.49
1:B:102:LEU:O	1:B:103:LYS:HE3	2.13	0.49
1:A:24:LYS:NZ	1:A:104:GLU:HG3	2.29	0.48
1:B:2[B]:GLU:HG3	1:B:3:CYS:SG	2.53	0.48
1:B:106:GLU:HB3	1:B:108:PHE:CZ	2.49	0.47
1:B:24:LYS:HE3	1:B:24:LYS:HB3	1.87	0.45
1:A:18[A]:ASN:HD21	1:B:18:ASN:HB2	1.82	0.43
1:A:68[B]:LEU:CD1	1:A:86:LEU:CD2	2.97	0.43
1:B:106:GLU:HG3	1:B:107:GLN:O	2.21	0.41
1:B:4:SER:HA	1:B:30:THR:O	2.20	0.41
1:A:68[B]:LEU:HD11	1:A:86:LEU:CD2	2.51	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2[B]:GLU:OE1	3:B:1:TRS:O3[4_554]	1.80	0.40
1:A:2[B]:GLU:OE1	3:A:129:TRS:O1[4_455]	1.80	0.40
1:B:2[B]:GLU:OE1	3:B:1:TRS:C3[4_554]	2.05	0.15
1:A:2[B]:GLU:OE1	3:A:129:TRS:C1[4_455]	2.12	0.08

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	128/127 (101%)	124 (97%)	4 (3%)	0	100 100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	127/127 (100%)	123 (97%)	4 (3%)	0	100	100
All	All	255/254 (100%)	247 (97%)	8 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/110 (104%)	113 (99%)	1 (1%)	78	49
1	B	113/110 (103%)	112 (99%)	1 (1%)	78	49
All	All	227/220 (103%)	225 (99%)	2 (1%)	78	49

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	85	LYS
1	B	103	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	8	GLN
1	B	18	ASN
1	B	28	GLN
1	B	107	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	TRS	A	129	-	7,7,7	1.69	1 (14%)	9,9,9	0.77	0
3	TRS	B	1	-	7,7,7	1.51	1 (14%)	9,9,9	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRS	A	129	-	-	3/9/9/9	-
3	TRS	B	1	-	-	3/9/9/9	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	129	TRS	O1-C1	4.41	1.56	1.42
3	B	1	TRS	O3-C3	3.55	1.54	1.42

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	129	TRS	N-C-C1-O1
3	A	129	TRS	C2-C-C1-O1
3	B	1	TRS	N-C-C3-O3
3	A	129	TRS	C3-C-C1-O1
3	B	1	TRS	C1-C-C3-O3
3	B	1	TRS	C2-C-C3-O3

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	129	TRS	2	2
3	B	1	TRS	1	2

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	127/127 (100%)	0.54	7 (5%)	25 25	8, 11, 19, 33	0
1	B	127/127 (100%)	0.58	9 (7%)	16 16	8, 11, 19, 33	0
All	All	254/254 (100%)	0.56	16 (6%)	20 21	8, 11, 19, 33	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	105	GLY	8.0
1	A	104	GLU	5.2
1	B	128	LYS	5.0
1	B	104	GLU	3.8
1	A	120	LEU	3.8
1	A	127	LEU	3.2
1	B	103	LYS	3.0
1	B	102	LEU	3.0
1	B	127	LEU	2.8
1	B	2[A]	GLU	2.4
1	A	2[A]	GLU	2.3
1	A	128	LYS	2.3
1	B	106	GLU	2.2
1	A	105	GLY	2.2
1	A	103	LYS	2.2
1	B	24	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	TRS	A	129	8/8	0.90	0.14	5,14,16,20	0
3	TRS	B	1	8/8	0.92	0.13	5,13,15,15	0
2	CU	B	129	1/1	1.00	0.05	8,8,8,8	0
2	CU	A	1	1/1	1.00	0.05	8,8,8,8	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.