



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 11:39 pm BST

PDB ID : 1U4F  
Title : Crystal Structure of Cytoplasmic Domains of IRK1 (Kir2.1) channel  
Authors : Pegan, S.; Arrabit, C.; Zhou, W.; Kwiatkowski, W.; Slesinger, P.A.; Choe, S.  
Deposited on : 2004-07-24  
Resolution : 2.41 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11




# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4647 (2.44-2.40)
Clashscore	141614	5161 (2.44-2.40)
Ramachandran outliers	138981	5073 (2.44-2.40)
Sidechain outliers	138945	5074 (2.44-2.40)
RSRZ outliers	127900	4543 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	270	<div> <div>3%</div> <div> <div></div> <div>54%</div> <div>18%</div> <div>•</div> <div>27%</div> </div> </div>
1	B	270	<div> <div>3%</div> <div> <div></div> <div>55%</div> <div>16%</div> <div>•</div> <div>28%</div> </div> </div>
1	C	270	<div> <div>3%</div> <div> <div></div> <div>50%</div> <div>21%</div> <div>•</div> <div>28%</div> </div> </div>
1	D	270	<div> <div>3%</div> <div> <div></div> <div>50%</div> <div>22%</div> <div>•</div> <div>27%</div> </div> </div>



## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6533 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inward rectifier potassium channel 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	197	Total	C	N	O	S	0	0	0
			1590	1006	277	299	8			
1	B	195	Total	C	N	O	S	0	0	0
			1570	994	271	297	8			
1	C	195	Total	C	N	O	S	0	0	0
			1570	994	271	297	8			
1	D	197	Total	C	N	O	S	0	0	0
			1590	1012	273	297	8			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	35	PRO	-	CLONING ARTIFACT	UNP P35561
A	36	ARG	-	CLONING ARTIFACT	UNP P35561
A	37	GLY	-	CLONING ARTIFACT	UNP P35561
A	38	SER	-	CLONING ARTIFACT	UNP P35561
A	39	HIS	-	CLONING ARTIFACT	UNP P35561
A	40	GLY	-	CLONING ARTIFACT	UNP P35561
B	35	PRO	-	CLONING ARTIFACT	UNP P35561
B	36	ARG	-	CLONING ARTIFACT	UNP P35561
B	37	GLY	-	CLONING ARTIFACT	UNP P35561
B	38	SER	-	CLONING ARTIFACT	UNP P35561
B	39	HIS	-	CLONING ARTIFACT	UNP P35561
B	40	GLY	-	CLONING ARTIFACT	UNP P35561
C	35	PRO	-	CLONING ARTIFACT	UNP P35561
C	36	ARG	-	CLONING ARTIFACT	UNP P35561
C	37	GLY	-	CLONING ARTIFACT	UNP P35561
C	38	SER	-	CLONING ARTIFACT	UNP P35561
C	39	HIS	-	CLONING ARTIFACT	UNP P35561
C	40	GLY	-	CLONING ARTIFACT	UNP P35561
D	35	PRO	-	CLONING ARTIFACT	UNP P35561
D	36	ARG	-	CLONING ARTIFACT	UNP P35561
D	37	GLY	-	CLONING ARTIFACT	UNP P35561

*Continued on next page...*



*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	38	SER	-	CLONING ARTIFACT	UNP P35561
D	39	HIS	-	CLONING ARTIFACT	UNP P35561
D	40	GLY	-	CLONING ARTIFACT	UNP P35561

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	55	Total	O	0	0
			55	55		
2	B	48	Total	O	0	0
			48	48		
2	C	59	Total	O	0	0
			59	59		
2	D	51	Total	O	0	0
			51	51		




### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Inward rectifier potassium channel 2



#### • Molecule 1: Inward rectifier potassium channel 2



#### • Molecule 1: Inward rectifier potassium channel 2





SER	ASN	ALA	ASN	SER	PHE	CYS	TYR	GLU	ASN	GLU	VAL	ALA	LEU	THR	SER	LYS	GLU	GLU	GLU	ASP	SER	GLU	ASN	GLY	VAL	PRO	GLU	SER	THR	SER	THR	SER	ASP	SER	PRO	PRO	GLY	ILE	ASP	LEU	HIS	ASN	GLN	ALA	SER	VAL	PRO	GLU	PRO	ARG	GLU	SER	GLU	ILE
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

● Molecule 1: Inward rectifier potassium channel 2



PRO	ARG	GLY	SER	HIS	GLY	GLN	GLN	CYS	S45	K49	H53	C54	N55	V56	Q57	F58	V61	G62	E63	K188	ARG	ASN	GLU	T192	L201	L202	M203	R204	D205	L208	C209	V214	G215	N216	L217	R218	K219	S220	V223	E224	R228	L232	K233	S234	T237	S238	E239	G240
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

E241	P244	Q247	V252	G253	T258	L263	I267	T268	I269	E272	L279	T280	D281	L282	S283	K284	I287	F292	V295	T296	I297	L298	E303	R312	S313	S314	H324	R325	Y326	E327	K334	H335	V339	E340	Y341	K346	E349	L355	R359
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

K362	E363	K364	K365	Y366	I367	L368	SER	ASN	ALA	ASN	SER	PHE	CYS	TYR	GLU	ASN	GLU	VAL	ALA	LEU	THR	SER	LYS	GLU	GLU	GLU	GLU	ASP	SER	GLU	ASN	GLY	VAL	PRO	GLU	SER	THR	THR	ASP	SER	PRO	PRO	GLY	ILE	ASP	LEU	HIS	ASN	GLN	ALA	SER	VAL	PRO	LEU	GLU	PRO	ARG
------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

PRO	LEU	ARG	ARG	GLU	SER	GLU	ILE
-----	-----	-----	-----	-----	-----	-----	-----



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.46Å 138.29Å 138.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.44 – 2.41 41.62 – 2.41	Depositor EDS
% Data completeness (in resolution range)	(Not available) (38.44-2.41) 91.8 (41.62-2.41)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.11 (at 2.39Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.225 , 0.281 0.226 , 0.280	Depositor DCC
$R_{free}$ test set	1977 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.8	Xtriage
Anisotropy	0.849	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 45.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6533	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/1622	0.60	0/2192
1	B	0.39	0/1601	0.62	0/2164
1	C	0.41	0/1601	0.63	0/2164
1	D	0.38	0/1622	0.59	0/2192
All	All	0.39	0/6446	0.61	0/8712

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1590	0	1570	47	0
1	B	1570	0	1543	42	0
1	C	1570	0	1543	47	0
1	D	1590	0	1575	51	0
2	A	55	0	0	0	0
2	B	48	0	0	1	0
2	C	59	0	0	2	0
2	D	51	0	0	2	0
All	All	6533	0	6231	172	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.



The worst 5 of 172 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:301:MET:HG2	1:A:308:THR:HG22	1.50	0.93
1:B:237:THR:HG23	1:B:239:GLU:H	1.38	0.87
1:A:188:LYS:HA	1:A:191:GLU:HG2	1.58	0.84
1:B:237:THR:HG22	1:B:241:GLU:H	1.43	0.83
1:A:223:VAL:HG12	1:A:224:GLU:HG3	1.62	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	195/270 (72%)	182 (93%)	9 (5%)	4 (2%)	7	7
1	B	191/270 (71%)	176 (92%)	13 (7%)	2 (1%)	15	22
1	C	191/270 (71%)	179 (94%)	12 (6%)	0	100	100
1	D	193/270 (72%)	182 (94%)	11 (6%)	0	100	100
All	All	770/1080 (71%)	719 (93%)	45 (6%)	6 (1%)	19	27

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	335	HIS
1	A	191	GLU
1	A	218	ARG
1	B	244	PRO
1	B	351	PRO



### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	177/244 (72%)	174 (98%)	3 (2%)	60	77
1	B	175/244 (72%)	169 (97%)	6 (3%)	37	54
1	C	175/244 (72%)	170 (97%)	5 (3%)	42	61
1	D	177/244 (72%)	170 (96%)	7 (4%)	31	48
All	All	704/976 (72%)	683 (97%)	21 (3%)	41	59

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	63	GLU
1	C	263	LEU
1	D	252	VAL
1	B	314	SER
1	D	263	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	221	HIS
1	C	60	ASN
1	D	226	HIS
1	B	226	HIS
1	B	271	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	197/270 (72%)	-0.00	9 (4%) 32 30	28, 48, 84, 105	0
1	B	195/270 (72%)	0.09	9 (4%) 32 30	26, 46, 77, 100	0
1	C	195/270 (72%)	-0.09	8 (4%) 37 35	26, 44, 73, 96	0
1	D	197/270 (72%)	0.04	9 (4%) 32 30	29, 47, 81, 110	0
All	All	784/1080 (72%)	0.01	35 (4%) 33 31	26, 46, 80, 110	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	367	ILE	7.3
1	D	366	TYR	7.3
1	D	368	LEU	5.1
1	A	365	LYS	4.5
1	C	190	ASN	4.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.