



wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 02:38 pm BST

PDB ID : 4U6B
Title : Zg3597, a family 117 glycoside hydrolase, produced by the marine bacterium
Zobellia galactanivorans
Authors : Ficko-Blean, E.
Deposited on : 2014-07-28
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	<div> <div>2%</div> <div> <div></div> <div>61%</div> <div>14%</div> <div>•</div> <div>21%</div> </div> </div>
1	B	433	<div> <div>%</div> <div> <div></div> <div>63%</div> <div>13%</div> <div>•</div> <div>21%</div> </div> </div>
1	C	433	<div> <div>2%</div> <div> <div></div> <div>61%</div> <div>14%</div> <div>• •</div> <div>21%</div> </div> </div>
1	D	433	<div> <div>2%</div> <div> <div></div> <div>62%</div> <div>15%</div> <div>•</div> <div>21%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	ACY	B	503	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 11568 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Conserved hypothetical lipoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	343	Total	C	N	O	S	0	2	0
			2686	1720	447	513	6			
1	B	343	Total	C	N	O	S	0	4	0
			2705	1734	447	518	6			
1	C	342	Total	C	N	O	S	0	3	0
			2693	1729	447	511	6			
1	D	343	Total	C	N	O	S	0	1	0
			2680	1717	442	515	6			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Ca	0	0
			2	2		
2	A	2	Total	Ca	0	0
			2	2		
2	D	2	Total	Ca	0	0
			2	2		
2	C	2	Total	Ca	0	0
			2	2		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Na	0	0
			1	1		

- Molecule 6 is ACETIC ACID (three-letter code: ACY) (formula: C₂H₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	168	Total	O	0	7
			175	175		
7	B	173	Total	O	0	7
			180	180		
7	C	185	Total	O	0	10
			195	195		
7	D	175	Total	O	0	12
			187	187		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Conserved hypothetical lipoprotein




• Molecule 1: Conserved hypothetical lipoprotein




• Molecule 1: Conserved hypothetical lipoprotein





● Molecule 1: Conserved hypothetical lipoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	F 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	187.79Å 223.52Å 225.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.30 47.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.00-2.30) 99.7 (47.00-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.76 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.178 , 0.225 0.178 , 0.225	Depositor DCC
R_{free} test set	5189 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	36.3	Xtriage
Anisotropy	0.441	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.043 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11568	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.02 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2163e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACY, NA, CA, PEG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.93	1/2782 (0.0%)	0.93	9/3805 (0.2%)
1	B	0.93	3/2801 (0.1%)	0.96	8/3829 (0.2%)
1	C	0.93	1/2792 (0.0%)	0.92	8/3813 (0.2%)
1	D	0.92	1/2773 (0.0%)	0.94	8/3793 (0.2%)
All	All	0.93	6/11148 (0.1%)	0.94	33/15240 (0.2%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	152	TRP	CE3-CZ3	7.47	1.51	1.38
1	D	243	ASP	CB-CG	-6.61	1.37	1.51
1	B	243	ASP	CB-CG	-5.57	1.40	1.51
1	B	152	TRP	CE3-CZ3	5.53	1.47	1.38
1	B	225	PRO	N-CD	5.20	1.55	1.47

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	243	ASP	CB-CG-OD1	-13.99	105.71	118.30
1	D	243	ASP	CB-CG-OD1	-10.78	108.60	118.30
1	D	331	MET	CG-SD-CE	-10.15	83.95	100.20
1	B	243	ASP	CB-CG-OD2	9.71	127.04	118.30
1	A	331	MET	CG-SD-CE	-7.46	88.27	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2686	0	2478	61	0
1	B	2705	0	2502	49	0
1	C	2693	0	2515	73	0
1	D	2680	0	2469	54	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	14	0	20	1	0
4	A	4	0	6	3	0
4	B	12	0	18	3	0
4	C	12	0	18	1	0
4	D	12	0	18	4	0
5	A	1	0	0	0	0
6	B	4	0	3	3	0
7	A	175	0	0	8	0
7	B	180	0	0	7	0
7	C	195	0	0	8	0
7	D	187	0	0	2	1
All	All	11568	0	10047	229	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 229 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:308:THR:O	4:A:505:EDO:H12	1.47	1.12
1:A:78:GLU:HG2	1:A:400:LEU:HD23	1.36	1.02
1:D:78:GLU:HG2	1:D:400:LEU:HD23	1.40	1.01
1:A:104:ASP:H	1:A:376:HIS:HD2	1.06	0.96
1:C:314[B]:LYS:HG2	1:C:315:GLU:H	1.30	0.95

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:D:601:HOH:O	7:D:601:HOH:O[2_555]	2.02	0.18

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	341/433 (79%)	322 (94%)	18 (5%)	1 (0%)	41	50
1	B	343/433 (79%)	327 (95%)	15 (4%)	1 (0%)	41	50
1	C	341/433 (79%)	327 (96%)	14 (4%)	0	100	100
1	D	340/433 (78%)	328 (96%)	12 (4%)	0	100	100
All	All	1365/1732 (79%)	1304 (96%)	59 (4%)	2 (0%)	51	64

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	211	TRP
1	B	132	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	274/366 (75%)	256 (93%)	18 (7%)	16	22
1	B	277/366 (76%)	258 (93%)	19 (7%)	15	20
1	C	277/366 (76%)	261 (94%)	16 (6%)	20	27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	274/366 (75%)	254 (93%)	20 (7%)	14	18
All	All	1102/1464 (75%)	1029 (93%)	73 (7%)	16	22

5 of 73 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	364	SER
1	C	126	VAL
1	D	323	LYS
1	C	79	LEU
1	C	135	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 39 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	82	GLN
1	C	199	GLN
1	D	281	HIS
1	C	94	ASN
1	C	95	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 9 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	B	505	-	3,3,3	0.65	0	2,2,2	0.32	0
4	EDO	C	503	-	3,3,3	0.70	0	2,2,2	0.07	0
3	PEG	A	503	-	6,6,6	0.49	0	5,5,5	0.59	0
4	EDO	D	505	-	3,3,3	0.66	0	2,2,2	0.18	0
4	EDO	C	504	-	3,3,3	0.58	0	2,2,2	0.33	0
4	EDO	D	504	-	3,3,3	0.57	0	2,2,2	0.44	0
4	EDO	B	504	-	3,3,3	0.29	0	2,2,2	0.64	0
3	PEG	A	504	-	6,6,6	0.48	0	5,5,5	0.67	0
4	EDO	A	505	-	3,3,3	0.29	0	2,2,2	0.80	0
4	EDO	D	503	-	3,3,3	0.55	0	2,2,2	0.46	0
4	EDO	C	505	-	3,3,3	0.39	0	2,2,2	0.33	0
4	EDO	B	506	-	3,3,3	0.53	0	2,2,2	0.75	0
6	ACY	B	503	-	1,3,3	1.03	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	505	-	-	1/1/1/1	-
4	EDO	C	503	-	-	0/1/1/1	-
3	PEG	A	503	-	-	3/4/4/4	-
4	EDO	D	505	-	-	1/1/1/1	-
4	EDO	C	504	-	-	0/1/1/1	-
4	EDO	D	504	-	-	1/1/1/1	-
4	EDO	B	504	-	-	1/1/1/1	-
3	PEG	A	504	-	-	2/4/4/4	-
4	EDO	A	505	-	-	1/1/1/1	-
4	EDO	D	503	-	-	1/1/1/1	-
4	EDO	C	505	-	-	0/1/1/1	-
4	EDO	B	506	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	503	PEG	O1-C1-C2-O2
3	A	503	PEG	O2-C3-C4-O4
4	B	505	EDO	O1-C1-C2-O2
4	B	504	EDO	O1-C1-C2-O2
4	A	505	EDO	O1-C1-C2-O2

There are no ring outliers.

8 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	505	EDO	1	0
4	B	504	EDO	2	0
3	A	504	PEG	1	0
4	A	505	EDO	3	0
4	D	503	EDO	3	0
4	C	505	EDO	1	0
4	B	506	EDO	1	0
6	B	503	ACY	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	343/433 (79%)	-0.20	9 (2%) 56 63	17, 36, 70, 101	0
1	B	343/433 (79%)	-0.21	6 (1%) 70 76	17, 35, 74, 122	0
1	C	342/433 (78%)	-0.27	10 (2%) 51 58	16, 35, 70, 114	0
1	D	343/433 (79%)	-0.17	7 (2%) 65 71	18, 36, 71, 111	0
All	All	1371/1732 (79%)	-0.21	32 (2%) 60 67	16, 36, 73, 122	0

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	207	LEU	5.3
1	C	208	ARG	4.7
1	D	206	GLY	4.6
1	C	207	LEU	4.3
1	B	135	LYS	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	EDO	B	506	4/4	0.63	0.27	66,73,74,78	0
4	EDO	C	503	4/4	0.68	0.27	66,66,74,83	0
4	EDO	D	505	4/4	0.73	0.28	68,69,70,78	0
4	EDO	C	504	4/4	0.73	0.28	73,76,79,80	0
4	EDO	B	505	4/4	0.77	0.30	54,55,66,71	0
6	ACY	B	503	4/4	0.80	0.20	57,66,69,70	0
3	PEG	A	503	7/7	0.83	0.21	62,72,84,87	0
4	EDO	D	503	4/4	0.83	0.22	70,72,79,80	0
4	EDO	D	504	4/4	0.84	0.23	69,72,77,77	0
3	PEG	A	504	7/7	0.86	0.28	63,81,92,93	0
5	NA	A	506	1/1	0.87	0.20	66,66,66,66	0
4	EDO	C	505	4/4	0.94	0.19	41,43,46,47	0
2	CA	A	501	1/1	0.96	0.03	49,49,49,49	0
4	EDO	A	505	4/4	0.97	0.18	48,50,53,54	0
4	EDO	B	504	4/4	0.98	0.17	38,44,45,52	0
2	CA	D	501	1/1	0.98	0.04	46,46,46,46	0
2	CA	C	501	1/1	0.99	0.03	45,45,45,45	0
2	CA	B	501	1/1	0.99	0.03	40,40,40,40	0
2	CA	B	502	1/1	1.00	0.14	27,27,27,27	0
2	CA	A	502	1/1	1.00	0.12	27,27,27,27	0
2	CA	D	502	1/1	1.00	0.13	29,29,29,29	0
2	CA	C	502	1/1	1.00	0.12	25,25,25,25	0

6.5 Other polymers ⓘ

There are no such residues in this entry.