



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 10, 2020 – 04:14 AM BST

PDB ID : 3U75  
Title : Structure of E230A-fructofuranosidase from Schwanniomyces occidentalis complexed with fructosylmaltose  
Authors : Sainz-Polo, M.A.; Sanz-Aparicio, J.  
Deposited on : 2011-10-13  
Resolution : 2.68 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3863 (2.70-2.66)
Clashscore	141614	4210 (2.70-2.66)
Ramachandran outliers	138981	4141 (2.70-2.66)
Sidechain outliers	138945	4141 (2.70-2.66)
RSRZ outliers	127900	3780 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	535	
1	B	535	
1	C	535	
1	D	535	
2	E	5	
2	F	5	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	G	5	 20% 40% 40%
2	H	5	 40% 60%

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 17209 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fructofuranosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	512	Total	C	N	O	S	0	0	0
			4152	2668	680	798	6			
1	B	512	Total	C	N	O	S	0	0	0
			4152	2668	680	798	6			
1	C	512	Total	C	N	O	S	0	0	0
			4152	2668	680	798	6			
1	D	512	Total	C	N	O	S	0	0	0
			4152	2668	680	798	6			

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-2)-beta-D-fructofuranose-(2-1)-beta-D-fructofuranose-(2-1)-beta-D-fructofuranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	E	5	Total	C	O	0	0	0
			56	30	26			
2	F	5	Total	C	O	0	0	0
			56	30	26			
2	G	5	Total	C	O	0	0	0
			56	30	26			
2	H	5	Total	C	O	0	0	0
			56	30	26			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	82	Total	O	0	0
			82	82		
4	B	69	Total	O	0	0
			69	69		
4	C	71	Total	O	0	0
			71	71		
4	D	57	Total	O	0	0
			57	57		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Fructofuranosidase




#### • Molecule 1: Fructofuranosidase



• Molecule 1: Fructofuranosidase



• Molecule 1: Fructofuranosidase




- Molecule 2: alpha-D-glucopyranose-(1-2)-beta-D-fructofuranose-(2-1)-beta-D-fructofuranose-(2-1)-beta-D-fructofuranose-(2-1)-beta-D-fructofuranose

Chain E: 




- Molecule 2: alpha-D-glucopyranose-(1-2)-beta-D-fructofuranose-(2-1)-beta-D-fructofuranose-(2-1)-beta-D-fructofuranose-(2-1)-beta-D-fructofuranose

Chain F: 



- Molecule 2: alpha-D-glucopyranose-(1-2)-beta-D-fructofuranose-(2-1)-beta-D-fructofuranose-(2-1)-beta-D-fructofuranose-(2-1)-beta-D-fructofuranose

Chain G: 



- Molecule 2: alpha-D-glucopyranose-(1-2)-beta-D-fructofuranose-(2-1)-beta-D-fructofuranose-(2-1)-beta-D-fructofuranose-(2-1)-beta-D-fructofuranose

Chain H: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.21Å 92.66Å 117.75Å 89.91° 82.96° 85.82°	Depositor
Resolution (Å)	50.00 – 2.68 58.43 – 2.68	Depositor EDS
% Data completeness (in resolution range)	96.8 (50.00-2.68) 79.2 (58.43-2.68)	Depositor EDS
$R_{merge}$	0.22	Depositor
$R_{sym}$	0.22	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.76 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, $R_{free}$	0.241 , 0.308 0.241 , 0.308	Depositor DCC
$R_{free}$ test set	3720 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.9	Xtriage
Anisotropy	0.100	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 13.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	17209	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, NAG, FRU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.66	0/4273	0.74	1/5820 (0.0%)
1	B	0.64	0/4273	0.73	1/5820 (0.0%)
1	C	0.73	0/4273	0.80	2/5820 (0.0%)
1	D	0.66	0/4273	0.76	1/5820 (0.0%)
All	All	0.67	0/17092	0.76	5/23280 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	D	0	1
All	All	0	2

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	C	259	LEU	CA-CB-CG	6.60	130.49	115.30
1	A	27	LEU	CA-CB-CG	5.92	128.92	115.30
1	C	27	LEU	CA-CB-CG	5.34	127.58	115.30
1	D	251	LEU	CA-CB-CG	5.32	127.53	115.30
1	B	27	LEU	CA-CB-CG	5.01	126.82	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	197	LYS	Peptide
1	D	197	LYS	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4152	0	3940	115	0
1	B	4152	0	3939	108	0
1	C	4152	0	3939	161	0
1	D	4152	0	3939	187	0
2	E	56	0	52	5	0
2	F	56	0	52	3	0
2	G	56	0	52	4	0
2	H	56	0	52	5	0
3	A	14	0	13	0	0
3	B	28	0	26	1	0
3	C	28	0	26	0	0
3	D	28	0	26	1	0
4	A	82	0	0	4	0
4	B	69	0	0	3	0
4	C	71	0	0	6	0
4	D	57	0	0	5	0
All	All	17209	0	16056	573	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 573 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:36:ARG:O	1:B:483:LYS:HD3	1.56	1.06
4:C:582:HOH:O	2:G:4:FRU:H61	1.55	1.04
1:D:333:ARG:HH11	1:D:353:GLN:NE2	1.56	1.02
1:C:164:TYR:O	1:C:165:GLU:HB2	1.60	1.00
1:A:170:ILE:HD12	1:A:193:MET:HE1	1.45	0.97

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	510/535 (95%)	463 (91%)	35 (7%)	12 (2%)	6	13
1	B	510/535 (95%)	462 (91%)	36 (7%)	12 (2%)	6	13
1	C	510/535 (95%)	448 (88%)	51 (10%)	11 (2%)	6	15
1	D	510/535 (95%)	438 (86%)	58 (11%)	14 (3%)	5	11
All	All	2040/2140 (95%)	1811 (89%)	180 (9%)	49 (2%)	6	13

5 of 49 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	25	ILE
1	A	421	ASN
1	B	26	ASP
1	B	27	LEU
1	B	421	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	463/484 (96%)	443 (96%)	20 (4%)	29	54
1	B	463/484 (96%)	451 (97%)	12 (3%)	46	73
1	C	463/484 (96%)	434 (94%)	29 (6%)	18	37

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	463/484 (96%)	428 (92%)	35 (8%)	13	28
All	All	1852/1936 (96%)	1756 (95%)	96 (5%)	23	46

5 of 96 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	210	ASN
1	C	371	ASN
1	D	420	LEU
1	C	241	ASN
1	C	306	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 69 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	348	GLN
1	C	78	GLN
1	D	353	GLN
1	B	353	GLN
1	C	52	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

20 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FRU	E	1	2	11,11,12	0.52	0	15,15,18	1.43	3 (20%)
2	FRU	E	2	2	11,11,12	0.71	0	15,15,18	1.08	1 (6%)
2	FRU	E	3	2	11,11,12	0.50	0	15,15,18	1.10	1 (6%)
2	FRU	E	4	2	11,11,12	0.34	0	15,15,18	0.71	0
2	GLC	E	5	2	12,12,12	0.45	0	17,17,17	1.64	4 (23%)
2	FRU	F	1	2	11,11,12	0.49	0	15,15,18	1.44	2 (13%)
2	FRU	F	2	2	11,11,12	0.50	0	15,15,18	1.36	3 (20%)
2	FRU	F	3	2	11,11,12	0.55	0	15,15,18	1.08	0
2	FRU	F	4	2	11,11,12	0.42	0	15,15,18	0.76	0
2	GLC	F	5	2	12,12,12	0.69	0	17,17,17	1.32	1 (5%)
2	FRU	G	1	2	11,11,12	0.35	0	15,15,18	1.32	2 (13%)
2	FRU	G	2	2	11,11,12	0.51	0	15,15,18	1.28	2 (13%)
2	FRU	G	3	2	11,11,12	0.65	0	15,15,18	0.94	0
2	FRU	G	4	2	11,11,12	0.43	0	15,15,18	1.22	1 (6%)
2	GLC	G	5	2	12,12,12	0.85	0	17,17,17	1.87	3 (17%)
2	FRU	H	1	2	11,11,12	0.32	0	15,15,18	1.30	2 (13%)
2	FRU	H	2	2	11,11,12	0.75	0	15,15,18	1.24	3 (20%)
2	FRU	H	3	2	11,11,12	0.45	0	15,15,18	1.10	0
2	FRU	H	4	2	11,11,12	0.39	0	15,15,18	1.26	1 (6%)
2	GLC	H	5	2	12,12,12	0.57	0	17,17,17	1.59	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FRU	E	1	2	-	0/4/20/24	0/1/1/1
2	FRU	E	2	2	-	4/4/20/24	0/1/1/1
2	FRU	E	3	2	-	2/4/20/24	0/1/1/1
2	FRU	E	4	2	-	2/4/20/24	0/1/1/1
2	GLC	E	5	2	-	2/2/22/22	0/1/1/1
2	FRU	F	1	2	-	0/4/20/24	0/1/1/1
2	FRU	F	2	2	-	4/4/20/24	0/1/1/1
2	FRU	F	3	2	-	1/4/20/24	0/1/1/1
2	FRU	F	4	2	-	2/4/20/24	0/1/1/1
2	GLC	F	5	2	-	1/2/22/22	0/1/1/1
2	FRU	G	1	2	-	2/4/20/24	0/1/1/1

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FRU	G	2	2	-	4/4/20/24	0/1/1/1
2	FRU	G	3	2	-	2/4/20/24	0/1/1/1
2	FRU	G	4	2	-	0/4/20/24	0/1/1/1
2	GLC	G	5	2	-	0/2/22/22	0/1/1/1
2	FRU	H	1	2	-	0/4/20/24	0/1/1/1
2	FRU	H	2	2	-	4/4/20/24	0/1/1/1
2	FRU	H	3	2	-	0/4/20/24	0/1/1/1
2	FRU	H	4	2	-	2/4/20/24	0/1/1/1
2	GLC	H	5	2	-	2/2/22/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	5	GLC	C1-O5-C5	5.01	123.11	113.66
2	E	5	GLC	O5-C5-C4	3.67	116.36	109.69
2	F	5	GLC	C1-O5-C5	3.63	120.51	113.66
2	G	5	GLC	O5-C5-C4	3.44	115.94	109.69
2	H	5	GLC	C1-O5-C5	3.29	119.87	113.66

There are no chirality outliers.

5 of 34 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	H	2	FRU	O1-C1-C2-C3
2	H	2	FRU	O1-C1-C2-O5
2	F	4	FRU	O5-C5-C6-O6
2	E	4	FRU	O5-C5-C6-O6
2	E	2	FRU	O1-C1-C2-C3

There are no ring outliers.

12 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	3	FRU	2	0
2	F	4	FRU	2	0
2	H	5	GLC	2	0
2	G	4	FRU	1	0
2	G	1	FRU	3	0
2	F	1	FRU	1	0


*Continued on next page...*




*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	5	GLC	1	0
2	H	4	FRU	2	0
2	E	2	FRU	1	0
2	E	3	FRU	1	0
2	H	1	FRU	1	0
2	E	1	FRU	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.




## Oligosaccharide Chain F



## Oligosaccharide Chain G





## 5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	D	1394	1	14,14,15	0.94	0	17,19,21	2.57	7 (41%)
3	NAG	C	1219	1	14,14,15	0.97	0	17,19,21	1.63	4 (23%)
3	NAG	A	1219	1	14,14,15	0.54	0	17,19,21	1.22	3 (17%)
3	NAG	B	1219	1	14,14,15	0.76	0	17,19,21	1.35	3 (17%)
3	NAG	D	1219	1	14,14,15	0.74	0	17,19,21	1.36	3 (17%)
3	NAG	C	1394	1	14,14,15	0.59	0	17,19,21	2.08	5 (29%)
3	NAG	B	1394	1	14,14,15	0.53	0	17,19,21	1.70	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	D	1394	1	-	3/6/23/26	0/1/1/1
3	NAG	C	1219	1	-	0/6/23/26	0/1/1/1
3	NAG	A	1219	1	-	2/6/23/26	0/1/1/1
3	NAG	B	1219	1	-	0/6/23/26	0/1/1/1
3	NAG	D	1219	1	-	1/6/23/26	0/1/1/1
3	NAG	C	1394	1	-	3/6/23/26	0/1/1/1
3	NAG	B	1394	1	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1394	NAG	C2-N2-C7	6.46	132.11	122.90
3	B	1394	NAG	C1-O5-C5	5.64	119.83	112.19
3	C	1394	NAG	C1-O5-C5	4.66	118.50	112.19
3	C	1394	NAG	O5-C1-C2	-4.36	104.40	111.29
3	D	1394	NAG	C1-C2-N2	-3.88	103.85	110.49

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	1394	NAG	C3-C2-N2-C7
3	A	1219	NAG	C4-C5-C6-O6
3	A	1219	NAG	O5-C5-C6-O6
3	C	1394	NAG	O5-C5-C6-O6
3	B	1394	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	1394	NAG	1	0
3	B	1394	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	512/535 (95%)	-0.15	0 100 100	21, 31, 40, 51	0
1	B	512/535 (95%)	-0.09	3 (0%) 89 90	20, 31, 40, 56	0
1	C	512/535 (95%)	-0.21	2 (0%) 92 93	13, 30, 44, 59	0
1	D	512/535 (95%)	0.02	7 (1%) 75 76	18, 38, 52, 63	0
All	All	2048/2140 (95%)	-0.11	12 (0%) 89 90	13, 32, 47, 63	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	25	ILE	3.9
1	D	100	ILE	3.3
1	C	25	ILE	2.8
1	B	25	ILE	2.7
1	D	356	VAL	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GLC	F	5	12/12	0.84	0.21	44,47,48,49	0
2	FRU	H	3	11/12	0.90	0.19	31,32,35,37	0

*Continued on next page...*

*Continued from previous page...*




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GLC	H	5	12/12	0.92	0.12	40,42,42,43	0
2	FRU	H	2	11/12	0.92	0.16	31,35,39,40	0
2	FRU	G	3	11/12	0.92	0.17	26,29,30,31	0
2	GLC	E	5	12/12	0.92	0.17	40,48,50,53	0
2	FRU	E	4	11/12	0.93	0.15	28,35,37,37	0
2	GLC	G	5	12/12	0.93	0.11	33,38,40,41	0
2	FRU	H	4	11/12	0.93	0.14	26,33,35,37	0
2	FRU	F	3	11/12	0.93	0.17	26,29,30,31	0
2	FRU	E	3	11/12	0.93	0.14	23,27,30,30	0
2	FRU	G	4	11/12	0.93	0.13	27,32,33,34	0
2	FRU	G	1	11/12	0.94	0.16	24,27,28,28	0
2	FRU	G	2	11/12	0.95	0.12	26,31,33,36	0
2	FRU	F	4	11/12	0.95	0.13	33,39,40,41	0
2	FRU	H	1	11/12	0.95	0.13	25,29,31,33	0
2	FRU	F	1	11/12	0.95	0.13	17,20,21,23	0
2	FRU	F	2	11/12	0.96	0.13	22,29,33,37	0
2	FRU	E	2	11/12	0.96	0.13	26,31,36,38	0
2	FRU	E	1	11/12	0.97	0.12	22,23,24,25	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around Chain F:**


$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain G:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	D	1394	14/15	0.80	0.26	44,46,53,54	0
3	NAG	B	1394	14/15	0.84	0.16	48,53,56,56	0
3	NAG	B	1219	14/15	0.88	0.16	39,42,44,45	0
3	NAG	A	1219	14/15	0.88	0.17	43,46,48,48	0
3	NAG	D	1219	14/15	0.89	0.16	41,44,47,47	0
3	NAG	C	1219	14/15	0.89	0.14	37,41,47,48	0
3	NAG	C	1394	14/15	0.90	0.13	45,50,53,53	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.